



May 20th, 3:30 PM - 4:00 PM

A Lubricated Immersed Boundary Method in Two Dimensions

Thomas G. Fai

Harvard University, tfai@seas.harvard.edu

Follow this and additional works at: <http://scholarscompass.vcu.edu/bamm>



Part of the [Numerical Analysis and Computation Commons](#)

<http://scholarscompass.vcu.edu/bamm/2016/May20/2>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

A LUBRICATED IMMERSED BOUNDARY METHOD IN TWO DIMENSIONS

Thomas Fai¹, Chris Rycroft²

^{1,2}*Paulson School of Engineering and Applied Sciences, Harvard University, 29 Oxford St., Cambridge, MA 02138*

¹*Email: tfai@seas.harvard.edu*

Abstract

We describe an immersed boundary method that uses elements of lubrication theory to resolve thin fluid layers between immersed boundaries. Such methods are useful for simulating the passage of deformable elastic capsules through narrow constrictions, such as the transit of red blood cells through the narrow slits in the spleen and the intracellular trafficking of vesicles into dendritic spines. Making use of simple two-dimensional flows with known exact solutions, we will show convergence results illustrating the increased accuracy in comparison to the standard immersed boundary method. We further apply this method to the case of an elastic vesicle moving through a fluid-filled channel.