



2013

Quantification of Thoracic Aorta Blood Flow by Magnetic Resonance Imaging During Supine Cycling Exercise of Increasing Intensity

Laura Ellwein

Virginia Commonwealth University, lellwein@vcu.edu

John F. LaDisa

Marquette University

Stacy Leibham

Children's Hospital of Wisconsin

Sheila Schindler-Ivens

Marquette University

Margaret M. Samyn

Marquette University

Follow this and additional works at: http://scholarscompass.vcu.edu/math_pubs

© 2013 Ellwein et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Downloaded from

http://scholarscompass.vcu.edu/math_pubs/11

This Article is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Mathematics and Applied Mathematics Publications by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

POSTER PRESENTATION

Open Access

Quantification of thoracic aorta blood flow by magnetic resonance imaging during supine cycling exercise of increasing intensity

Laura Ellwein^{2,1*}, John F LaDisa^{2,6}, Stacy Leibham⁴, Sheila Schindler-Ivens⁵, Margaret M Samyn^{3,2}

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

Cardiac magnetic resonance imaging (MRI) may be used for the diagnosis and follow-up of diseases affecting the thoracic aorta (TA). Typically, patients are imaged at rest. Quantifying TA blood flow distribution during activity may improve understanding of disease impact by assessing possible factors contributing to exercise intolerance. Blood flow quantification during exercise MRI, though, has heretofore been complicated by prolonged MRI scan times, motion artifact, and gradient field inhomogeneities. We developed a protocol for phase-contrast (PC) MRI flow quantification in the TA and head and neck arteries during three-tiered supine cycling submaximal exercise.

Methods

PC-MRI was acquired in the ascending aorta (AAo) and innominate (IA), left common carotid (LCCA), and left subclavian (LSA) arteries using a 1.5T Siemens MAGNETOM[®] Symphony magnet. Six healthy volunteers (28±2 years) were imaged during rest and supine pedaling of an

MRI-compatible cycle to reach 130%, 150%, and 170% of resting heart rate (HR₁₃₀, HR₁₅₀, HR₁₇₀). Scan parameters balanced image quality with acquisition time, accounting for subject positioning in the magnet and motion from pedaling. Time-resolved volumetric blood flow was calculated from PC-MRI data and estimated in the descending aorta (dAo). Flow quantification included cardiac index from AAo flow, arterial normalized mean flow (NMF, L/min/m²) and flow distribution (FD, % of AAo flow). Significance of flow quantification between HRs and rest was tested post-hoc with Student's t-test.

Results

Cardiac index increased linearly ($r^2=0.99$) from 3.1±0.3 to 5.2±0.3 L/min/m² from rest to HR₁₇₀ ($p<0.05$). HR increased to 111±5.9 bpm and workload to 38±9 watts at HR₁₇₀. In the dAo, NMF increased linearly with respect to HR ($r^2=0.99$) by 46%, 72%, and 93% with significance at all levels (Table 1). The IA showed significant increase in NMF of 39% at HR₁₇₀ compared to rest. Increases in NMF

Table 1 Flow quantification (mean ± SE, n=6) in the descending aorta (dAo), innominate (IA), left common carotid (LCCA), and left subclavian (LSA) arteries. (Statistically significant results are in bold, p<0.05)

	NMF, L/min/m ²				FD, % of AAo flow			
	Rest	HR ₁₃₀	HR ₁₅₀	HR ₁₇₀	Rest	HR ₁₃₀	HR ₁₅₀	HR ₁₇₀
dAo	2.12±0.23	3.00±0.24	3.51±0.22	3.91±0.23	67±1.6%	72±1.5%	76±0.1%	75±1.3%
IA	0.50±0.03	0.59±0.07	0.59±0.07	0.69±0.07	16±0.8%	14±1.0%	13±1.0%	13±1.0%
LCCA	0.23±0.01	0.27±0.02	0.24±0.02	0.26±0.03	7.6±0.7%	6.5±0.4%	5.3±0.6%	5.0±0.5%
LSA	0.28±0.04	0.34±0.05	0.30±0.04	0.38±0.06	8.7±0.9%	7.9±0.7%	6.4±0.6%	7.0±2.0%

NMF: Normalized mean flow; FD: Flow distribution.

²Biomedical Engineering, Marquette University, Milwaukee, WI, USA
Full list of author information is available at the end of the article

in the LCCA and LSA were not significant. FD in the dAo increased significantly by 12% and 11% at HR₁₅₀ and HR₁₇₀ compared to rest, with concomitant decreases in the LCCA (27%, 31%) and LSA (25%, 20%). FD in the IA decreased at each HR level but without significance.

Conclusions

This investigation marks the first known flow quantification in the TA and head and neck arteries during supine exercise cardiac MRI. These pilot data show the changes in cardiac output and preferential redistribution of blood flow during increasing exercise intensity in a group of healthy young adults. Future studies using the protocol introduced here can now be conducted for various populations with TA disease.

Funding

This work was supported by an AHA Postdoctoral Fellowship award 10POST4210030 (to LME) and an NIH grant R15HL096096-01 (to JEL).

Author details

¹Mathematics and Applied Mathematics, Virginia Commonwealth University, Richmond, VA, USA. ²Biomedical Engineering, Marquette University, Milwaukee, WI, USA. ³Pediatrics, Herma Heart Center, Children's Hospital of Wisconsin, Milwaukee, WI, USA. ⁴Radiology, Children's Hospital of Wisconsin, Milwaukee, WI, USA. ⁵Physical Therapy, Marquette University, Milwaukee, WI, USA. ⁶Medicine, Division of Cardiovascular Medicine, Medical College of Wisconsin, Milwaukee, WI, USA.

Published: 30 January 2013

doi:10.1186/1532-429X-15-S1-P239

Cite this article as: Ellwein *et al.*: Quantification of thoracic aorta blood flow by magnetic resonance imaging during supine cycling exercise of increasing intensity. *Journal of Cardiovascular Magnetic Resonance* 2013 15(Suppl 1):P239.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

