



# VCU

Virginia Commonwealth University  
**VCU Scholars Compass**

---

Undergraduate Research Posters

Undergraduate Research Opportunities Program

---

2013

## PERCUSSION AND VIBRATION AIRWAY CLEARANCE THERAPY INCREASES INFLAMMATORY FACTORS

Gil Rubia

*Virginia Commonwealth University*

Mike Davis

*Virginia Commonwealth University*

Alison Montpetit

*Virginia Commonwealth University*

Follow this and additional works at: <https://scholarscompass.vcu.edu/uressposters>

© The Author(s)

---

### Downloaded from

Rubia, Gil; Davis, Mike; and Montpetit, Alison, "PERCUSSION AND VIBRATION AIRWAY CLEARANCE THERAPY INCREASES INFLAMMATORY FACTORS" (2013). *Undergraduate Research Posters*. Poster 12.

<https://scholarscompass.vcu.edu/uressposters/12>

This Article is brought to you for free and open access by the Undergraduate Research Opportunities Program at VCU Scholars Compass. It has been accepted for inclusion in Undergraduate Research Posters by an authorized administrator of VCU Scholars Compass. For more information, please contact [libcompass@vcu.edu](mailto:libcompass@vcu.edu).

# PERCUSSION AND VIBRATION AIRWAY CLEARANCE THERAPY INCREASES INFLAMMATORY FACTORS

Gil C. Rubia, Michael D. Davis, RT, Alison J. Montpetit, Ph.D., RN  
Virginia Commonwealth University School of Nursing, Richmond Va.

## Background



Ventilator associated pneumonia (VAP) is a preventable but common complication of mechanically ventilated (MV) patients due to oral secretions containing bacteria migrating to the lungs via the endotracheal tube. The development of VAP is associated with an increased risk of mortality and substantial increase in hospital stay and healthcare-associated costs. In order to prevent VAP, multiple pulmonary hygiene and airway and secretion clearance therapies (ASCT) have been utilized such as kinetic bed-delivered therapies which include: continuous lateral rotation (CLRT) and percussion/vibration (P/V) therapy.

## Objective

Currently, there is insufficient empirical evidence to support that kinetic bed therapy reduces the onset of nosocomial pneumonia. The purpose of this study is to mimic the clinical scenario of a patient receiving kinetic bed therapy followed by a bacterial exposure in an *in vitro* model through quantification of IL-8, IFN- $\gamma$ , TNF- $\alpha$ , 8-isoprostane and nitric oxide.

## Methods


Human airway 3D normal human bronchial epithelial (NHBE) air-liquid interface (ALI) cell culture model were grown and followed with bacterial challenge through the application of lipopolysaccharide (LPS). Culture was subjected to three different kinetic conditions (1-Acute: 15 min P/V, 2-Intermittent: 15 min P/V per hour for 6 hours, 3-Continuous: 6 hour P/V). Trans-epithelial electric resistance values were taken before and after each condition. In this experiment, NHBE cells underwent submerged culture conditions for two weeks, and then transitioned to ALI for another two weeks. Cell supernatants were collected, stored at -80° C and batch processed via multiplex cytokine bead assay for IL-8, IFN- $\gamma$ , TNF- $\alpha$  and via enzyme immunoassay (EIA) for 8-isoprostane. Supernatant samples were also analyzed for nitric oxide.



**Figure 2 (left):** LPS+ and LPS- stimulated NHBE cells underwent P/V for 6 and 12 hrs. n=3

## Results

- Continuous vibration loosens tight junctions (Figure 1)
- 8-isoprostane secretions increased the longer it was stimulated (Figure 2)



**Figure 1 (top):** Trans epithelial electric resistance decreased considerably from a healthy tissue layer to non-viable tissue layer.

## Acknowledgements

Dr. Bruce K. Rubin – School of Medicine, Department of Pediatrics  
Melissa A. Yopp – Mucus Lab  
Christopher Corday – Mucus Lab  
Shauna Webb-Parker – Mucus Lab  
Dr. Clive M. Baumgarten – School of Medicine, Department of Physiology and Biophysics  
Dr. Paul A. Wetzel – School of Engineering, Department of Biomedical Engineering  
Anathea Pepperl – Wetzel Lab  
\*National Institute of Nursing Research (K99/R00 NR 012016, A. Montpetit, PI), VCU School of Nursing Adult Health and Nursing Systems Department

\*\*Image from hill-rom.com

## Discussion

- Vibration and percussion therapy increases inflammation through the activation of the inflammatory and oxidative stress cascade. Breakdown of tight junctions increases the onset of the inflammation and oxidative stress cascade.
- It is unknown if movement, such as continuous lateral rotation, has a more profound effect in preventing infection in the lower airway.
- Analysis of IL-8, IFN- $\gamma$ , TNF- $\alpha$ , and nitric oxide is pending.

