



2017

Scedosporium prolificans Septic Arthritis

Collier S. Pace

Virginia Commonwealth University, collier.pace@gmail.com

Jessica A. Frankenhoff

Virginia Commonwealth University

Jonathan E. Isaacs

Virginia Commonwealth University

Follow this and additional works at: http://scholarscompass.vcu.edu/surgery_pubs

 Part of the [Surgery Commons](#)

© 2017 Society of Indian Hand & Microsurgeons

Downloaded from

http://scholarscompass.vcu.edu/surgery_pubs/35

This Article is brought to you for free and open access by the Dept. of Surgery at VCU Scholars Compass. It has been accepted for inclusion in Surgery Publications by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Scedosporium prolificans Septic Arthritis

Collier S. Pace¹ Jessica A. Frankenhoff² Jonathan E. Isaacs²

¹Division of Plastic and Reconstructive Surgery, Virginia Commonwealth University Medical Center, Richmond, Virginia, United States

²Department of Orthopedic Surgery, Virginia Commonwealth University Medical Center, Richmond, Virginia, United States

J Hand Microsurg 2017;9:37–38.

Scedosporium prolificans is an emerging fungal pathogen that can cause significant morbidity, and even mortality, in both immunocompromised and immunocompetent patients. Approximately 10% of patients affected by this rare fungal pathogen present with septic osteomyelitis or arthritis. Overall, the rate of mortality is close to 50%, and several patients with orthopedic infections have required amputations.¹

Our patient is a 59-year-old woman, with a history of rheumatoid arthritis requiring immunosuppressants, who presented with a 5-month history of right wrist pain and swelling. She described an oscillating course of swelling, erythema, and pain involving the dorsal aspect of her wrist that did not show sustained response to antibiotics or a radiocarpal joint steroid injection given by other providers. On initial examination, she was afebrile with dorsal wrist swelling. There was no erythema, but wrist mobility was limited. She had tenderness throughout the wrist, but no fluctuance or drainage. All initial laboratory work was normal. Right wrist X-ray showed severe osteopenia with some mild evidence of cortical erosion.

Despite the normal laboratory values, her presenting findings of swelling, radiographic erosions, and immunocompromised state prompted concern for atypical septic arthritis. The patient subsequently underwent wrist exploration for tissue and culture harvest. In addition to extensive wrist synovectomy, the proximal pole of the scaphoid, lunate, triquetrum, capitate, hamate, and metacarpal bases all showed evidence of necrosis that required debridement. One week later, her fluid aspirate and bone cultures unexpectedly grew an unidentified fungal species, and she was taken back to the operating room for further debridement and placement of a voriconazole-impregnated cement spacer.

The culture grew pan-resistant *Scedosporium prolificans*, and based on recommendations from Rheumatology and Infectious Disease, the patient was started on a 6-week course

Address for correspondence Collier S. Pace, MD, Division of Plastic and Reconstructive Surgery, Virginia Commonwealth University Medical Center, 1200 E Broad St., Richmond, VA 23298, United States (e-mail: collier.pace@gmail.com).

of intravenous micafungin, and immunosuppressive medications were stopped. She returned to the operating room for debridement and serial treatments with polyhexamethylene biguanide (PHMB) irrigation until intraoperative tissue cultures showed no growth. PHMB is an antiseptic medication that has been described as a local adjunct to the treatment of resistant fungal infections.^{2–4} After successful eradication of her infection, she underwent wrist reconstruction and fusion with a double-barrel free fibula osteocutaneous flap. She is now 18 months out from her reconstruction, has healed uneventfully, and has a functional, painless upper extremity (→ Fig. 1).



Fig. 1 Radiograph 5 months after reconstruction with signs of bony healing.

received
October 31, 2016
accepted after revision
November 9, 2016
published online
December 7, 2016

© 2017 Society of Indian Hand & Microsurgeons

DOI <http://dx.doi.org/10.1055/s-0036-1597553>.
ISSN 0974-3227.

Due to the rare nature of this infection, all the information and guidance for treatment comes from case reports or case series. The most comprehensive analysis of cases from the literature compiles data on 162 patients with *Scedosporium prolificans* infection.¹ The overall mortality for the 162 patients in the study was 46.9%; however, the vast majority of fatal infections occurred in immunocompromised patients with disseminated disease, and only 10% of patients in this series had osteomyelitis or septic arthritis. All, except one, of the patients with orthopedic infections suffered some form of previous trauma. None of these patients died, but two required amputations for cure.

This case demonstrates the high potential for morbidity associated with *Scedosporium prolificans* septic arthritis and osteomyelitis of the wrist. We describe a successful eradication of this challenging pathogen and a creative solution for reconstruction with a double-barrel free fibula osteocutaneous flap. Hand surgeons should consider this pathogen when evaluating patients for atypical infections, especially if the patient is immunocompromised. Evidence-based treatment guidelines are not currently available, but treatment should generally include adequate debridement, systemic antifungal therapy, and potentially PHMB irrigation.

Note

All identifying details were withheld to protect the privacy of our patient. Informed consent was obtained for photographs and their use for publication.

Funding

None.

Conflict of Interest

None.

References

- 1 Rodriguez-Tudela JL, Berenguer J, Guarro J, et al. Epidemiology and outcome of *Scedosporium prolificans* infection, a review of 162 cases. *Med Mycol* 2009;47(4):359–370
- 2 Hübner NO, Kramer A. Review on the efficacy, safety and clinical applications of polihexanide, a modern wound antiseptic. *Skin Pharmacol Physiol* 2010;23 (Suppl):17–27
- 3 Steinbach WJ, Schell WA, Miller JL, Perfect JR. *Scedosporium prolificans* osteomyelitis in an immunocompetent child treated with voriconazole and caspofungin, as well as locally applied polyhexamethylene biguanide. *J Clin Microbiol* 2003;41(8):3981–3985
- 4 Walls G, Noonan L, Wilson E, Holland D, Briggs S. Successful use of locally applied polyhexamethylene biguanide as an adjunct to the treatment of fungal osteomyelitis. *Can J Infect Dis Med Microbiol* 2013;24(2):109–112