

May 15th, 6:30 PM

Optimal spraying strategies for controlling re-infestation by Chagas disease vectors

Bismark Oduro
oduro@calu.edu

Winfried Just

Mario Grijalva

Follow this and additional works at: <https://scholarscompass.vcu.edu/bamm>

Part of the [Ordinary Differential Equations and Applied Dynamics Commons](#), and the [Other Applied Mathematics Commons](#)

<https://scholarscompass.vcu.edu/bamm/2019/wed/27>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Chagas disease is a major health problem in rural South and Central America where an estimated 8 to 11 million people are infected. It is a vector-borne disease caused by the parasite *Trypanosoma cruzi*, which is transmitted to humans mainly through the bite of insect vectors from several species of so-called "kissing bugs."

One of the control measures to reduce the spread of the disease is insecticide spraying of housing units to prevent infestation by the vectors. However, re-infestation of units by vectors has been shown to occur as early as four to six months after insecticide-based control interventions.

I will present ordinary differential equation models of type SIRS with imperfect treatment that shed light on long-term cost-effectiveness of certain strategies for controlling re-infestation by vectors. The results show that an initially very high spraying rate may push the system into a region of the state space with low endemic levels of infestation that can be maintained in the long run at relatively moderate cost.