



# VCU

Virginia Commonwealth University  
VCU Scholars Compass

---

Theses and Dissertations

Graduate School

---

2004

## PCR Detection of Streptococcus Mutans and Streptococcus Sobrinus in Dental Plaque Samples from Low, Moderate, and High Caries Risk Children

Ajit A. Patel  
*Virginia Commonwealth University*

Follow this and additional works at: <https://scholarscompass.vcu.edu/etd>



Part of the [Pediatric Dentistry and Pedodontics Commons](#)

© The Author

---

Downloaded from

<https://scholarscompass.vcu.edu/etd/1009>

This Thesis is brought to you for free and open access by the Graduate School at VCU Scholars Compass. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of VCU Scholars Compass. For more information, please contact [libcompass@vcu.edu](mailto:libcompass@vcu.edu).

School of Dentistry  
Virginia Commonwealth University

This is to certify that the thesis prepared by Ajit A. Patel entitled “PCR detection of *S. mutans* and *S. sobrinus* in dental plaque samples from low, moderate, and high caries risk children” has been approved by his committee as satisfactory completion of the thesis or dissertation requirement for the degree of Master of Science.

---

Arthur P. Mourino, D.D.S., M.S.D., Thesis Director, Professor Emeritus, Department of Pediatric Dentistry

---

Tegwyn H. Brickhouse, D.D.S., Ph.D., Committee Member, Department of Pediatric Dentistry

---

Francis L. Macrina, Ph.D., Committee Member, Department of Oral and Craniofacial Molecular Biology

---

Michael D. Webb, D.D.S., M.S., Post Graduate Director, Department of Pediatric Dentistry

---

Laurie Carter, D.D.S., Ph.D., Director of Advanced Education Programs, School of Dentistry

---

Dr. F. Douglas Boudinot, Dean of the School of Graduate Studies

October 18, 2004

© Ajit A. Patel, 2004

All Rights Reserved

PCR DETECTION OF *STREPTOCOCCUS MUTANS* AND *STREPTOCOCCUS SOBRINUS* IN DENTAL PLAQUE SAMPLES FROM LOW, MODERATE, AND HIGH CARIES RISK CHILDREN

A Thesis submitted in partial fulfillment of the requirements for the degree of Master of Science at Virginia Commonwealth University.

by

AJIT A. PATEL, D.D.S.  
Virginia Commonwealth University, 2002

Director: ARTHUR P. MOURINO, D.D.S., M.S.D.  
PROFESSOR EMERITUS  
DEPARTMENT OF PEDIATRIC DENTISTRY

Virginia Commonwealth University  
Richmond, Virginia  
October, 2004

### Acknowledgement

The author would like to thank several people. I would like to thank my parents and my sisters for their love and support throughout my life. I would like to extend sincere acknowledgements and thanks to my Thesis Committee Members, Dr. Arthur Mourino, Dr. Tegwyn Brickhouse and Dr. Francis Macrina for their much appreciated assistance with this research project. Your guidance and mentorship has been integral in completing my research successfully. Thank you for all of your help and support. I would like to give a special thanks to Kevin Jones for help with the laboratory aspect of this project. Your hard work was much appreciated. The author would also like to thank Dr. Al Best for calculating the statistical data presented in this study. Most of all, I would like to thank my wife, Manisha Patel, who has been very patient and supportive throughout my professional development.

## Table of Contents

	Page
Acknowledgements.....	ii
List of Tables .....	iv
List of Figures.....	v
Abstract.....	vi
Chapter	
1 Introduction.....	1
2 Methods.....	8
3 Results.....	13
4 Discussion.....	16
References.....	26
Appendix	
A Research subject Information and Consent Form.....	31
B Vita.....	36

## List of Tables

	Page
Table 1: Caries-risk Assessment Tool 1 (CAT 1).....	19
Table 2: Caries-risk Assessment Tool 2 (CAT 2).....	20
Table 3: Description of Patients.....	21
Table 4: Bacterial Prevalence. ....	21
Table 5: Relationship between CAT 1 and CAT 2 (frequency count). ....	22
Table 6: Contingency Table of <i>S. mutans</i> by CAT 1.....	22
Table 7: Contingency Table of <i>S. sobrinus</i> by CAT 1. ....	23
Table 8: Contingency Table of <i>S. mutans</i> by CAT 2.....	23
Table 9: Contingency Table of <i>S. sobrinus</i> by CAT 2. ....	24

## List of Figures

	Page
Figure 1: The Relationship between the number Decayed or Filled (DdFf) and Age according to CAT 1.....	25



## Abstract

### PCR DETECTION OF *STREPTOCOCCUS MUTANS* AND *STREPTOCOCCUS SOBRINUS* IN DENTAL PLAQUE SAMPLES FROM LOW, MODERATE, AND HIGH CARIES RISK CHILDREN

By Ajit A. Patel, D.D.S.

A Thesis submitted in partial fulfillment of the requirements for the degree of Master of Science at Virginia Commonwealth University.

Virginia Commonwealth University, 2004

Major Director: Arthur P. Mourino, D.D.S., M.S.D.  
Professor Emeritus, Department of Pediatric Dentistry

**Purpose:** The purpose of this study was to correlate the presence of *Streptococcus mutans* and *Streptococcus sobrinus* with the incidence of dental caries in children as detected by the polymerase chain reaction (PCR).

**Methods:** Subjects between the ages 2 to 16 years of age were used in this study. After the subjects received a dental examination, dental plaque samples were collected by rubbing a toothpick on the buccal surface of the most posterior tooth in each quadrant of the mouth. Children were grouped into low, moderate, and high caries risk categories by two different Caries Risk Assessment Tools (CAT 1 and CAT 2). PCR was used to

amplify the dextranase (Dex) gene of *S. mutans* and *S. sobrinus*. Dextranase is an enzyme that hydrolyses glucans in a plaque matrix and is believed to be involved in the pathogenesis of dental caries. Identification of the amplified Dex gene sequences was made by gel electrophoresis which in turn was used to determine the presence of *S. mutans* and *S. sobrinus*.

**Results:** *S. mutans* was detected in 57% of patients (65/115), *S. sobrinus* was detected in 10% of patients (12/115) and both organisms were present in 9% of patients (10/115). According to CAT 1, the incidence of *S. mutans* was 64.6%, 9.2%, and 26.2% in low, moderate, and high caries risk, respectively. The incidence of *S. sobrinus* was 66.6%, 16.7%, and 16.7% in low, moderate, and high caries risk, respectively.

**Conclusion:** Presence of *S. mutans* and/or *S. sobrinus* as evaluated in this study had no direct correlation with caries risk level in children.

## Introduction

Dental caries has been a highly prevalent and costly disease in the United States and around the world.<sup>1</sup> Dental caries remains the single most common disease of childhood that is neither self-limiting nor amendable to short-term pharmacological management.<sup>2</sup> United States Surgeon General's report shows that 45% of children ages 5 to 17 have dental caries.<sup>3</sup>

Mutans streptococci (MS) is group of principal micro-organisms involved in the etiology of human dental caries and is divided into 5 species and 8 different serological types (a through h).<sup>4,5</sup> Among the mutans streptococci, *S. mutans* (serotypes c, e and f) and *S. sobrinus* (serotype d and g) are the species closely associated with human dental caries.<sup>4,6,7</sup>

Streptococcal species are the first to colonize the tooth surfaces. In doing so, certain species use specific receptors to attach to the salivary pellicle.<sup>8,9</sup> The pellicle is derived from salivary constituents which are selectively adsorbed onto the tooth surface. Components of the dental pellicle include albumin, lysozyme, amylase, immunoglobulin A, proline-rich proteins and mucins. The formation of pellicle is the first step in plaque formation.<sup>10</sup>

Dental plaque is a soft deposit that accumulates on the teeth. Plaque can be defined as a complex microbial community, with greater than  $10^{10}$  bacteria per milligram. It has

been estimated that as many as 400 distinct bacterial species may be found in plaque. In addition to the bacterial cells, plaque contains a small number of epithelial cells, leukocytes, and macrophages. The cells are contained within an extracellular matrix, which is formed from bacterial products and saliva.<sup>10</sup>

After the initial colonization of the tooth surface, plaque increases by two distinct mechanisms: 1) the multiplication of bacteria already attached to the tooth surface, and 2) the subsequent attachment and multiplication of new bacterial species to cells of bacteria already present in the plaque mass.<sup>10</sup> When provided a suitable metabolic substrate, this infection produces organic acids (primarily lactic acid) that are capable of dissolving the mineral calcium from the crystalline enamel matrix.<sup>1</sup> Continued and repeated dissolution leads to frank cavitation and the subsequent need for repair.<sup>11</sup>

Mutans streptococci (MS) are the group of microorganisms most associated with the dental caries process.<sup>12</sup> MS are believed to contribute to caries because of their ability to adhere to tooth surfaces, produce copious amounts of acid, and survive and continue metabolism at low pH conditions.<sup>7</sup>

While many carbohydrates may be utilized by plaque bacteria to generate acids, sucrose is recognized as being particularly important in the caries process because not only can it be fermented, but it also serves as substrate for extracellular enzymes of plaque bacteria which synthesize sucrose-derived polymers. These polymers are of central importance in adhesive interactions in plaque, where they mediate attachment of bacteria to the tooth surface and to other bacteria thus stabilizing the plaque biofilm, serve as

energy stores aiding the survival of plaque bacteria and modulate the permeability of plaque and hence the level of acid at the enamel surface.<sup>13</sup>

Several epidemiological studies have suggested that subjects harboring both *S. mutans* and *S. sobrinus* have a significantly higher level of past caries experience than those harboring *S. mutans*<sup>14-17</sup> or *S. sobrinus* alone<sup>14</sup>. Other studies have suggested that in a population with high caries prevalence *Strep. mutans* must be the significant organism since it is mostly found in the absence of *Strep. sobrinus*.<sup>13</sup>

However, there have been suggestions that *Strep. sobrinus* is more virulent than *S. mutans*.<sup>18</sup> de Soet *et al.*<sup>19</sup> observed that *S. sobrinus* strains were capable of higher acid production at low pH than *S. mutans* strains. Experimental animal studies have demonstrated that strains of *S. sobrinus* have a high cariogenic potential.<sup>20,21</sup> Studies also show that the prevalence of *S. sobrinus* in saliva was more closely associated with future caries activity, especially with smooth-surface caries increment, than the prevalence of *S. mutans* and that the subjects in the *S. sobrinus* group had both higher caries and higher decayed teeth and surface increments than the *S. mutans* group.<sup>17</sup>

Prevalence of *S. sobrinus* and *S. mutans* in the human oral cavity has been reported by epidemiological studies in which the isolation frequency of *S. mutans* from dental plaque is much higher than that of *S. sobrinus*.<sup>22-24</sup> Studies from around the world have shown that >98% of adults carry *Strep. mutans*, while different studies have found *Strep. sobrinus* in 7-35% of individuals.<sup>12</sup> It should be noted that the primary differentiation of these cariogenic species is usually based on the colonial morphology on mitis-salivarius (MS) or MS-bacitracin (MSB) agar.<sup>24-27</sup> However, this procedure is

sometimes inaccurate, and also time-consuming and laborious.<sup>28</sup> It has been reported that MSB inhibits the growth of *S. sobrinus* to a greater extent than *S. mutans*.<sup>25,29</sup> Therefore, it is conceivable that the actual incidence of *S. sobrinus* may be higher than currently reported.<sup>28</sup> The application of the polymerase chain reaction (PCR) method to human dental plaque showed that the prevalence of *S. sobrinus* (83%) in oral cavities was higher than currently reported (0-50%).<sup>28</sup> It is speculated that the difference depends on the detection method used: the detection method with PCR was performed directly on dental plaque without the culture step and biochemical tests.<sup>28</sup>

The results from previous studies have shown PCR oligonucleotide primers (SD1 and SD2) specific to dextranase gene of *S. mutans*<sup>30</sup> and pair of PCR primers (SOF14 and SOR1623) specific to dextranase gene of *S. sobrinus*<sup>28</sup>. Dextranase is an enzyme that hydrolyses glucans in a plaque matrix and is believed to be involved in the pathogenesis of dental caries.<sup>31,32</sup> Dextranase (Dex) is able to break down glucans to isomaltosaccharides 3-4 glucose units long by cleaving  $\alpha(1-6)$ -linkages within the dextran chain.<sup>33,34</sup> This activity may modify glucans by altering the ratio of  $\alpha(1-6)$ - to  $\alpha(1-3)$ -linked chains, hence influencing solubility and adhesive properties. Dex activity therefore, influences sucrose-dependent adherence by reducing the number of  $\alpha(1-6)$ -linkages in the glucan.<sup>13</sup>

Apart from its effect on the adhesiveness of glucans, the dextranase of *Strep. mutans* breaks down glucans to isomaltosaccharides which may then be transported into the cell via the multiple sugar metabolism (msm) operon,<sup>35,36</sup> where they are further degraded to glucose by a dextran glucosidase (DexB)<sup>31</sup>.

The PCR was able to amplify the DNA fragment from the *S. mutans* sample containing as few as 12 colony-forming units (cfu) prepared by the lysis method<sup>30</sup> and was able to detect as few as 9 cfu of *S. sobrinus*<sup>28</sup>. These results indicate that the primer pairs of *S. sobrinus* and *S. mutans* amplified the species-specific amplicons with different lengths, indicating that the present PCR method is useful for detection and identification of the human cariogenic species, *S. sobrinus* and *S. mutans*.<sup>28</sup>

Besides the precision of identifying the bacterial species present, there are a few other things that need to be considered to obtain an accurate caries risk status. Present literature has used the following factors to assess caries risk.

- 1). Number of teeth present (including their sealant status).
- 2). Previous caries experience: Studies show that “past caries experience is the most significant predictor”<sup>37-42</sup>, but young children may have no cavitations simply due to the fact that the disease has not had time to express itself.<sup>12</sup>
- 3). Smooth surface vs. Pit and fissure caries: Patients that exhibit only pit and fissure caries and not smooth surface caries have less risk of future caries than those who have extension of the disease onto smooth surface.<sup>1</sup>
- 4). Presence of white spot lesions or enamel defect: White spot lesions are the precursors to cavitated lesions. In young children, where the disease has not had the time to express itself, white spot lesions may be the only sign of caries experience. Presence of developmental structural defects in enamel may increase the caries risk in children by increasing plaque retention and acid production, thus increasing the susceptibility to tooth demineralization.<sup>12</sup>

- 5). Presence of visible plaque: Several studies have shown that there is a correlation between visible plaque on primary teeth and caries risk.<sup>43-45</sup> One study found that 91% of the children are correctly classified as to caries risk solely based on the presence or absence of visible plaque.<sup>44</sup>
- 6). Family history: Children who have parents/siblings with high level of caries are likely to have higher chances of getting cavities.
- 7). Sugar consumption: Frequency of carbohydrate consumption and sugar intake through fruit juices and flavored drinks has been implicated to cause caries. Models developed for young children who had little caries history include bacterial levels and sugar use as the strongest predictors.<sup>46-48</sup>
- 8). Socioeconomic level of family: National data shows that caries is highly prevalent in poor and near poor US preschool children.<sup>12</sup>
- 9). Fluoride exposure: If the fluoride content of drinking water is low and systemic fluoride supplementation is given, data from over 20 clinical trials show a caries reduction of 30%-80% in primary teeth from fluoride supplements, provided they are started close to birth and continued for 5 or more years.<sup>49-53</sup>

Recently, tools have been developed by National Institute of Health (NIH) and American Academy of Pediatric Dentistry (AAPD) to assess caries risk in children. The tool developed by NIH (CAT 1)<sup>8</sup> classified children's caries risk as follows: Children with decayed (D/d) or filled (F/f) surfaces less than half their age were classified as low risk. The letters D and F are used to designate decayed and filled permanent teeth. The letters d and f are used to designate decayed and filled primary teeth. Children with DFs more than



half their age were classified as moderate risk group. Finally, children with DFs more than their own age were considered as a high risk group.

The caries risk assessment tool suggested by AAPD (CAT 2)<sup>54</sup> classifies children as follows: Children with no carious teeth in the past 24 months were considered low risk. Children with carious teeth in the past 24 months were considered moderate risk. Lastly, children with carious teeth in the past 12 months or radiographic enamel caries were considered high risk.

In this study, plaque samples were collected from children ages 2 to 16 years of age. Children were grouped into low, moderate, and high caries risk categories by two different Caries Risk Assessment Tools. PCR primer pairs were used to amplify the *Dex* gene for detection and identification of *S. mutans* and *S. sobrinus*. The purpose of this study was to correlate the presence of *Streptococcus mutans* and *Streptococcus sobrinus* with the incidence of dental caries in children.

## Methods

One hundred fifteen children, ages 2-16 years old, who were patients of Virginia Commonwealth University, Department of Pediatric Dentistry, were included in the study. Consent for participation was obtained from at least one parent/guardian before the study. The consent form was approved on 2/9/04 and protocol number 03440 was assigned by VCU IRB (see Appendix A). If the child was between the ages seven and sixteen, an assent was also obtained from the child. The enrolled subjects received a dental examination with appropriate radiographs by a resident or a faculty member of the department. The caries risk level in patients was determined by two different caries risk assessment tools CAT 1 and CAT 2.

CAT 1 (Table 1) classified children with decayed (D/d) or filled (F/f) surfaces less than half their age, no new lesions in 1 year, no white spot lesions, no enamel defects, low titers of mutans streptococci, and high socioeconomic status as low risk group. Children with Dfs more than half their age, one or more lesions in 1 year, infrequent white spot lesions, enamel defects, moderate titers of mutans streptococci, middle socioeconomic status, and visible plaque were classified as moderate risk group. The criteria for high risk group included Dfs more than their own age, two or more lesions in 1 year, numerous white spot lesions, enamel defects, high titer of mutans streptococci, low socioeconomic setting, visible plaque, appliances in mouth and frequent sugar consumption.

Only the age component from CAT 1 is used in this study to determine the caries risk status. Patients with decayed (D/d) or filled (F/f) surfaces less than half their age were classified as low risk. Patients with DFs more than half their age were classified as moderate risk group. Finally, patients with DFs more than their own age were considered as a high risk group.

The revised Caries-risk assessment tool (CAT 2) suggested by AAPD (Table 2) includes three different criteria: clinical conditions, environmental characteristics, and general health conditions, to assess caries risk. Children with no carious teeth in the past 24 months, no enamel demineralization (“white spot lesions”), no visible plaque, no gingivitis, optimal systemic and topical fluoride exposure, consumption of simple sugars primarily at mealtimes, high caregiver socioeconomic status, and regular use of dental services are considered to be at low risk. Children with carious teeth in the past 24 months, one area of enamel demineralization, presence of gingivitis, suboptimal systemic fluoride exposure with optimal topical exposure, 1-2 between meal exposures to simple sugars, midlevel caregiver socioeconomic status, and irregular use of dental services are considered to be at moderate risk. Children with carious teeth in the past 12 months, more than one area of enamel demineralization, visible plaque on anterior teeth, radiographic enamel caries, high titers of mutans streptococci, wearing dental or orthodontic appliances, enamel hypoplasia, suboptimal topical fluoride exposure, 3 or more between meal exposures to simple sugars, low-level caregiver socioeconomic status, no usual source of dental care, and active caries present in mother are considered to be at high caries risk.

Children with special health care needs and children suffering from conditions impairing saliva composition/flow are also considered to be in high caries risk group.

CAT 2 component used in this study is as follows: Patients with no carious teeth in the past 24 months were considered low risk. Patients with carious teeth in the past 24 months were considered moderate risk. Lastly, patients with carious teeth in the past 12 months or radiographic enamel caries were considered high risk.

The letters D and F are used to designate decayed and filled permanent teeth, respectively. The letters d and f are used to designate decayed and filled primary teeth, respectively. For the purposes of this study, no distinction was made between primary and permanent teeth when recording the decayed and filled teeth.

#### *Patient Examination*

Multiple people (faculty and residents) were used to perform the examination. The data was then collected from patient charting. Examiner reliability was not done due to high number of faculty and residents.

#### *Plaque sampling*

Dental plaque samples were collected by rubbing a sterile toothpick on the buccal surface of the most posterior tooth in each quadrant of the mouth. The plaque sample was then placed into the GeneMate PCR strip-tubes with caps (0.2 ml size, cat.# T-3014-1) made by ISC-Bioexpress, Kaysville, UT. The PCR tubes contained 25  $\mu$ l of TB buffer. The samples were transported and stored at -4°C.

### *Extraction of Genomic DNA*

Frozen plaque samples were thawed by using a water bath at 55°C. 10 µl of each sample was transported on to FTA<sup>®</sup> (Flinders Technology Associates, Clifton, New Jersey) Cards by Whatman<sup>®</sup> and labeled appropriately. The cards were allowed to dry completely. Once dried, a 2.0mm disc was removed from each sample card, which contained a dried plaque sample, and placed into a separate microcentrifuge tubes (PCR tubes). The punch was washed for 5 min. by adding 200 µl of FTA<sup>®</sup> Purification Reagent to the tube and gently pipetting the reagent up and down twice. The reagent was discarded completely and the process was repeated two more times. The punch was washed by adding 200 µl of sterile TE<sup>-1</sup> buffer (10 mM Tris, 0.1 mM EDTA, pH 8.0) to the tube and gently pipetting the buffer up and down twice. The buffer was discarded completely and the process was repeated one more time. The discs were allowed to dry completely in the tubes.

### *Polymerase Chain Reaction (PCR)*

*S. mutans* (ATCC 25175) and *S. sobrinus* (ATCC 33478) were used as positive controls. The strains were obtained from ATCC (American Type Culture Collection), Manassas, VA. PCR detection of these species was performed using oligonucleotide primers designed to amplify a 1272-bp and 1610-bp fragment on the dextranase genes of *S. mutans* and *S. sobrinus*, respectively. SD1 and SD2 primers were used to specifically amplify a 1272-bp fragment in *S. mutans*. The sequence of SD1 and SD2 were 5'-TAT GCT GCT ATT GGA GGT TC-3' (positions 973 to 992) and 5'-AAG GTT GAG CAA

TTG AAT CG-3' (positions 2225 to 2244), respectively.<sup>10</sup> SOF14 and SOR 1623 primers were used to specifically amplify a 1610-bp fragment in *S. sobrinus*. The nucleotide sequence of SOF14 and SOR1623 were 5'-TGC TAT CTT TCC CTA GCA TG-3' (position 134-153) and 5'-GGT ATT CGG TTT GAC TGC-3' (positions 1743-1726), respectively.<sup>12</sup>

PCR reaction mixture (50 µl) was added into the tubes before starting the PCR. PCR reaction mixture was comprised of 1 µl of each of the two primers and 2 µl of Magnesium Chloride added to Invitrogen® PCR Supermix.

PCR was performed starting with the denaturation of the reaction mixture at 95°C for 3 min followed by a series of amplification: denaturation at 94°C for 30 sec, annealing at 52°C for 12 sec and extension at 72°C for 45 sec. The series was repeated for 28 cycles. The final cycle comprised of 94°C for 30 sec, 52°C for 12 sec, and 72°C for 7 min. The PCR products were then stored at 4°C for later use.

The PCR products were subjected to electrophoresis on a 0.8% agarose gel and stained with ethidium bromide.

#### *Statistical methods*

Groups were compared using a chi-square test of independence. All tests were performed at alpha = .05.

## Results

A total of  $n = 115$  subjects received a dental examination and forty-seven percent of these subjects were male (54/115). Table 3 shows a description of patients. Their average age was 7.3 years and they ranged from 2 years 1 month to 15 years 7 months. The number of decayed teeth on a patient ranged from 0 to 16, with the average of 3.19 decayed teeth per person. Each child had anywhere from 13 to 28 teeth with the average of 21.97 teeth per person. (Table 3) The subjects were measured for the presence of *S. mutans* and *S. sobrinus*. Caries risk was classified using only the dental caries indicators of CAT 1 and CAT 2. The number decayed (D, d) and filled (F, f) teeth were categorized low, medium, or high.

CAT 1: Children with DFs less than half their age were classified as low risk. Children with DFs more than their age were classified as high risk. The remaining children were classified as medium risk.

CAT 2: Children with no carious teeth in the past 24 months were considered low risk. Children with carious teeth in the past 24 months were considered moderate risk. Lastly, children with carious teeth or radiographic enamel caries in the past 12 months were considered high risk.

The CAT 1 method was used in Figure 1. The high risk children are above the upper line and the low risk children are below the lower line. The figure also shows that

there were no high risk caries children found above 11 years of age. There was only one child categorized as moderate caries risk above the age of 12 years. However, there was no statistical correlation with age.

*S. mutans* was detected in 57% of patients (65/115), *S. sobrinus* was detected in 10% of patients (12/115) and both organisms were present in 9% of patients (10/115, Table 4). The mean number of decayed teeth found in patients positive for both, *S. mutans* and *S. sobrinus*, was 2.8 teeth per patient. The highest number of decayed teeth (3.9) was found in patients negative for both *S. mutans* and *S. sobrinus* (Table 4).

The relationship between the CAT 1 and CAT 2 caries risk score is shown in Table 5. There were 40 patients “low” on both risk scores, 7 patients who were “low” caries risk according to CAT 1 and “moderate” caries risk according to CAT 2. Table 5 also shows the total number to low caries risk patients according to CAT 1 (n=67) and CAT 2 (n=41). There is a notable difference in the caries risk categorization between CAT 1 and CAT 2 as shown in Table 5. The majority of patients classified by CAT 1 are low caries risk patients (n=67), while the majority of patients classified by CAT 2 are high caries risk patients (n=63).

The relationship between *S. mutans* prevalence and caries risk (CAT 1) is shown in the contingency Table 6. This table shows that from the low risk group, 62.7% of children were positive for *S. mutans* and 37.3% were negative for *S. mutans*. The table also shows that from the children who were positive for *S. mutans*, 64.6% = low risk, 9.2% = moderate risk and 26.2% = high risk group. Pearson chi-square indicated no significant relationship (chi-square = 3.04, p-value = 0.2188).



The relationship between *S. sobrinus* prevalence and caries risk (CAT 1) is shown in the contingency Table 7. Pearson chi-square indicated no significant any relationship (chi-square = 0.97, p-value = 0.6148).

The relationship between *S. mutans* prevalence and the Caries Risk Assessment tool (CAT 2) is shown in the contingency Table 8. There was a trend towards a relationship (chi-square = 5.29, p-value = 0.0710).

The relationship between *S. sobrinus* prevalence and the CAT 2 is shown in the contingency Table 9. There was no significant relationship (chi-square = 1.4, p-value = 0.4847).

## Discussion

This study did not find a correlation between the presence of *S. mutans* and/or *S. sobrinus* in dental plaque in either CAT 1 or CAT 2 methods of risk assessment.

Identifying “risk” is more difficult in chronic diseases that are caused by multiple factors and events over a longer period of time in individuals. In the case of dental caries, the profession knows that bacteria capable of producing acids as a metabolic by product are necessary to produce the clinical manifestations of the disease. However, they are not in and of themselves “sufficient” to produce a cavitated tooth. Also needed are a susceptible host and a metabolic substrate on which these bacteria can act over time.<sup>1</sup>

This study was able to incorporate the new CAT 1 and CAT 2 methods that have been reported in the literature. Due to the recent creation of CAT 1 and CAT 2, there are no published studies making comparisons or examine the validity and reproducibility of these CAT methods. Due to time and limitations of funding, this study was not able to incorporate all of the criteria used in CAT 1 and CAT 2. The study was also based on a disproportionate number of young children. The population seen in the clinic is comprised of high number of young children. This study serves as a first step in the examination of caries risk assessment tools and the relationship to plaque by genomic plaque sampling methods.

Due to the nature of the environment, multiple people (faculty and residents) were used to perform the examination. The data on decayed and filled teeth was collected from the patient's charts. Examiner reliability was not assessed due to the number of faculty and residents. Inter-examiner reliability is a limitation of this study that could result in the misclassification of subjects according to CAT 1 and CAT 2.

According to CAT 1, the incidence of *S. mutans* was 64.6%, 9.2%, and 26.2% in low, moderate, and high caries risk, respectively. The incidence of *S. sobrinus* was 66.6%, 16.7%, and 16.7% in low, moderate, and high caries risk, respectively. CAT 2 also showed high number of *S. mutans* and *S. sobrinus* in low caries risk population. These results indicate a possible flaw in plaque sampling technique and/or misclassification of risk category according to caries risk assessment tools. Due to limited time and funding, all of the components of CAT 1 and CAT 2 were not used to determine the caries risk status. The results may have been different if all or more components of CAT 1 and CAT 2 were included in the study.

Future study should also include the use of real-time PCR in order to determine the amount of *S. mutans* and *S. sobrinus* in a given plaque sample and its relation to caries risk. Conventional PCR method used in this study revealed the presence or absence of the target template. Real-time PCR monitors the fluorescence emitted during the reaction as an indicator of amplicon production during each PCR cycle. This signal increases in direct proportion to the amount of PCR product in a reaction. By recording the amount of fluorescence emission at each cycle, it is possible to monitor the PCR reaction during exponential phase where the first significant increase in the amount of PCR product

correlates to the initial amount of target template. The higher the starting copy number of the nucleic acid target, the sooner a significant increase in fluorescence is observed. A significant increase in fluorescence above the baseline value measured during the 3-15 cycles indicates the detection of accumulated PCR product.<sup>55</sup>

This study was able to show a quick, accurate, and inexpensive diagnostic method of detecting the presence of *S. mutans* and *S. sobrinus*. With this new tool (PCR using FTA<sup>®</sup> cards), it is possible to diagnose within hours with a very small number of bacteria. This tool will aid in the identification of one of the variables used to determine caries risk and allow early preventive interventions. It will eliminate any inconsistency encountered by identifying the bacteria according to their colony morphology.

Table 1. Caries-risk Assessment Tool 1 (CAT 1)

Table 1. Caries-risk Assessment Tool 1 (CAT 1)		
Caries-risk indicators		
<b>Low risk</b>	<b>Moderate risk</b>	<b>High risk</b>
Dfs<half the child's age	Dfs>half the child's age	Dfs>child's age
No new lesions in 1 year	1 or more lesions in 1 year	2 or more lesions in 1 year
No white spot lesions	Infrequent white spot lesions	Numerous white spot lesions
No enamel defects	Enamel defects	Enamel defects
Low titers of mutans streptococci	Moderate titers of mutans streptococci	High titers of mutans streptococci
High socioeconomic setting	Middle socioeconomic setting	Low socioeconomic setting
	Visible plaque	Visible plaque
		Appliances in mouth
		Frequent sugar consumption

Adapted from NIH Consensus Development Conference on the Diagnosis and Management of Dental Caries. *J Dent Ed.* 2001;65:1133-1142.

Table 2. Caries-risk Assessment Tool 2 (CAT 2)

<b>Table 2. AAPD Caries-risk Assessment Tool (CAT 2)</b>		
<b>Caries-risk indicator</b>		
<b>Low risk</b>	<b>Moderate risk</b>	<b>High risk</b>
<b><i>Clinical conditions</i></b>		
No carious teeth in past 24 months	Carious teeth in the past 24 months	Carious teeth in the past 12 months
No enamel demineralization (enamel caries "white-spot lesions")	1 area of enamel demineralization (enamel caries "white-spot lesions")	More than 1 area of enamel demineralization (enamel caries "white-spot lesions")
No visible plaque; no gingivitis	Gingivitis	Visible plaque on anterior (front) teeth
		Radiographic enamel caries
		High titers of mutans streptococci
		Wearing dental or orthodontic appliances
		Enamel hypoplasia
<b><i>Environmental characteristics</i></b>		
Optimal systemic and topical fluoride exposure	Suboptimal systemic fluoride exposure with optimal topical exposure	Suboptimal topical fluoride exposure
Consumption of simple sugars or foods strongly associated with caries initiation- primarily at mealtimes	Occasional (ie, 1-2) between-meal exposures to simple sugars or foods strongly associated with caries	Frequent (ie, 3 or more) between-meal exposures to simple sugars or foods strongly associated with caries
High caregiver socioeconomic status	Midlevel caregiver socioeconomic status (ie, eligible for school lunch program or SCHIP)	Low-level caregiver socioeconomic status (ie, eligible for Medicaid)
Regular use of dental care in an established dental home	Irregular use of dental services	No usual source of dental care
		Active caries present in the mother
<b><i>General health conditions</i></b>		
		Children with special health care needs
		Conditions impairing saliva composition/flow

Table 3 Description of Patients

Table 3: Description of Patients				
Characteristic	Mean	Std. Dev.	Range	
Age (yrs)	7.33	3.44	2y 1mo	15y 7mo
Number of teeth:				
Decayed	3.19	4.28	0	16
Filled	0.81	2.14	0	10
Missing	0.15	0.68	0	4
Total teeth:	21.97	3.02	13	28

Table 4. Bacterial Prevalence

Table 4. Bacterial Prevalence					
			Mean		
Mutans	Sobrinus	Children (n)	Decayed	Missing	Filled
+	+	8.70%(10)	2.8	0.0	0.1
+	-	47.83%(55)	2.6	0.3	1.2
-	+	1.74%(2)	2.5	0.0	0.0
-	-	41.74%(48)	3.9	0.0	0.6

Table 5. Relationship between CAT 1 and CAT 2 (frequency count)

Table 5. Relationship between CAT 1 and CAT 2 (Frequency Count)				
	CAT 2 %(n)			
CAT 1 %(n)	Low	Moderate	High	Total
Low	34.78(40)	6.09(7)	17.39(20)	58.26(67)
Moderate	0(0)	0.87(1)	12.17(14)	13.04(15)
High	0.87(1)	2.61(3)	25.22(29)	28.7(33)
Total	35.65(41)	9.57(11)	54.78(63)	100(115)

Table 6. Contingency Table of S Mutans By CAT 1

Table 6. Contingency Table of <i>S. mutans</i> By CAT 1					
		Low	Moderate	High	Total
+	Count	42	6	17	65
	Column %	62.7	40.0	51.5	
	Row %	64.6	9.2	26.2	
-	Count	25	9	16	50
	Column %	37.3	60.0	48.5	
	Row %	50.0	18.0	32.0	
	Total	67	15	33	115



Table 7. Contingency Table of S Sobrinus By CAT 1

Table 7. Contingency Table of <i>S. sobrinus</i> By CAT 1					
		Low	Moderate	High	Total
+	Count	8	2	2	12
	Column %	11.9	13.3	6.1	
	Row %	66.6	16.7	16.7	
-	Count	59	13	31	103
	Column %	88.1	86.7	93.9	
	Row %	57.3	12.6	30.1	
Total		67	15	33	115


Table 8. Contingency Table of S Mutans By CAT 2

Table 8. Contingency Table of <i>S. mutans</i> By CAT 2					
		Low	Moderate	High	Total
+	Count	29	5	31	65
	Column %	70.7	45.5	49.2	
	Row %	44.6	7.7	47.7	
-	Count	12	6	32	50
	Column %	29.3	54.6	50.8	
	Row %	24.0	12.0	64.0	
Total		41	11	63	115

Table 9. Contingency Table of S Mutans By CAT 2

Table 9. Contingency Table of <i>S. sobrinus</i> By CAT 2					
		Low	Moderate	High	Total
+	Count	5	0	7	12
	Column %	12.2	0.0	11.1	
	Row %	41.7	0.0	58.3	
-	Count	36	11	56	103
	Column %	87.8	100.0	88.9	
	Row %	34.9	10.7	54.4	
	Total	41	11	63	115

Figure 1. The Relationship between the number Decayed or Filled (DdFf) and Age according to CAT 1.



Literature Cited

## Literature Cited

1. Anderson, M. Risk assessment and epidemiology of dental caries: review of the literature. *Pediatric Dentistry*. 2002; 24:5, 377-385.
2. Vargas CM, Crall JJ, Schneider DA. Sociodemographic Distribution of Pediatric Dental Caries: NHANES III, 1988-1994. *JADA*. 1998; 129: 1229-1238.
3. Allukian, M Jr. The neglected epidemic and the Surgeon General's report: a call to action for better oral health. *Am J Public Health*. 2000; 90: 843-345.
4. Hamada S, Slade HD. Biology, immunology, and cariogenicity of *Streptococcus mutans*. *Microbiol Rev*. 1980; 44: 331-384.
5. Whiley RA, Russell RRB, Hardie JM, Beighton D. *Streptococcus downei* sp. nov. for strains previously described as *Streptococcus mutans* serotype h. *Int J Syst Bacteriol*. 1988; 38: 25-29.
6. Batoni G, Marchetti F, Ota F et al. First characterization in Italy of clinical isolates of *mutans streptococci* by using specific monoclonal antibodies. *Eur J Epidemiol*. 1993; 9: 483-488.
7. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev*. 1986; 50: 353-383.
8. Jenkinson, H. F. Cell surface protein receptors in oral streptococci. *FEMS Microbiol Lett*. 1994; 121: 133-140.
9. Kolenbrander, P. E. and J. London. Adhere today, here tomorrow: oral bacterial adherence. *J. Bacteriol*. 1993; 175: 3247-3252.
10. Haake, SK. Microbiology of Dental Plaque.  
<http://www.dent.ucla.edu/pic/members/microbio/mdphome.html>. (Accessed May, 2004)
11. Featherstone JD. The science and practice of caries prevention. *JADA*. 2000; 131: 887-899.
12. Tinanoff N, Kanellis MJ, Vargas CM. Current understanding of the epidemiology, mechanisms, and prevention of dental caries in preschool children. *Pediatric Dentistry*. 2002; 24: 6,543-551.
13. Colby SM, Russell RRB. Sugar metabolism by *mutans streptococci*. *J of Applied Microbiol Symposium Supplement*. 1997; 83: 80S-88S.
14. Hirose H, Hirose K, Isogai E, Miura H, Ueda I. Close association between *Streptococcus sobrinus* in the saliva of young children and smooth-surface caries increment. *Caries Res*. 1993; 27: 292-297.
15. Köhler B, Bjarnaons S. *Mutans streptococci*, lactobacilli and caries prevalence in 11- and 12-year-old Icelandic children. *Community Dent Oral Epidemiol*. 1987; 15: 332-335.

16. Fujiwara T, Sasada E, Mima N, Ooshima T. Caries prevalence and salivary mutans streptococci in 0-2 year-old children of Japan. *Community Dent Oral Epidemiol.* 1991; 19: 151-154.
17. Okada M, Soda Y, Hayashi F, Doi T, Suzuki J, Miura K, Kozai K. PCR detection of *Streptococcus mutans* and *S. sobrinus* in dental plaque samples from Japanese pre-school children. *J Med Microbiol.* 2002; 51: 443-447.
18. de Soet JJ, Holbrook WP, Magnusdottir MO, De Graaff J. *Streptococcus sobrinus* and *Streptococcus mutans* in a longitudinal study of dental caries. *Microbial Ecology in Health and Disease.* 1993; 6: 237-243.
19. de Soet JJ, Toors FA, de Graaff J. Acidogenesis by oral streptococci at different pH values. *Caries Res.* 1989; 23(1):14-7.
20. de Soet JJ, van Loveren C, Lammens AJ, Pavicic MJ, Homburg CH, ten Cate JM, de Graaff J. Differences in cariogenicity between fresh isolates of *Streptococcus sobrinus* and *Streptococcus mutans*. *Caries Res.* 1991; 25(2):116-22.
21. Ooshima T, Sobue S, Hamada S, Kotani S. Susceptibility of rats, hamsters, and mice to carious infection by *Streptococcus mutans* serotype c and d organisms. *J Dent Res.* 1981 Apr; 60(4):855-9.
22. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev.* 1986 Dec; 50(4):353-80.
23. Carlsson P, Gandour IA, Olsson B, Rickardsson B, Abbas K. High prevalence of mutans streptococci in a population with extremely low prevalence of dental caries. *Oral Microbiol Immunol.* 1987 Sep; 2(3):121-4.
24. Van Palenstein Helderman WH, Ijsseldijk M, Huis in 't Veld JH. A selective medium for the two major subgroups of the bacterium *Streptococcus mutans* isolated from human dental plaque and saliva. *Arch Oral Biol.* 1983; 28(7):599-603.
25. Gold OG, Jordan HV, Van Houte J. A selective medium for *Streptococcus mutans*. *Arch Oral Biol.* 1973 Nov; 18(11):1357-64.
26. Svanberg M, Krasse B. Comparative recovery of mutans streptococci on two selective media. *Caries Res.* 1990; 24(1):36-8.
27. Wade WG, Aldred MJ, Walker DM. An improved medium for isolation of *Streptococcus mutans*. *J Med Microbiol.* 1986 Dec; 22(4):319-23.
28. Igarashi T, Yamamoto A, Goto N. PCR for detection and identification of *Streptococcus sobrinus*. *J Med Microbiol.* 2000 Dec; 49(12):1069-74.
29. de Soet JJ, van Dalen PJ, Pavicic MJ, de Graaff J. Enumeration of mutans streptococci in clinical samples by using monoclonal antibodies. *J Clin Microbiol.* 1990 Nov; 28(11):2467-72.
30. Igarashi T, Yamamoto A, Goto N. Direct detection of *Streptococcus mutans* in human dental plaque by polymerase chain reaction. *Oral Microbiol Immunol.* 1996 Oct; 11(5):294-8.
31. Colby SM, Russell RR. Sugar metabolism by mutans streptococci. *Soc Appl Bacteriol Symp Ser.* 1997; 26:80S-88S.
32. Pulkownik A, Walker GJ. Purification and substrate specificity of an endo-dextranase of *Streptococcus mutans* K1-R. *Carbohydrate Research.* 1977; 54:237-251.

33. Colby SM, Whiting GC, Tao L, Russell RR. Insertional inactivation of the *Streptococcus mutans* dexA (dextransase) gene results in altered adherence and dextran catabolism. *Microbiology*. 1995 Nov; 141 ( Pt 11):2929-36.
34. Walker GJ, Pulkownik A, Morrey-Jones JG. Metabolism of the polysaccharides of human dental plaque: release of dextransase in batch cultures of *Streptococcus mutans*. *J Gen Microbiol*. 1981 Nov; 127, pt 1:201-8.
35. Russell RR, Aduse-Opoku J, Sutcliffe IC, Tao L, Ferretti JJ. A binding protein-dependent transport system in *Streptococcus mutans* responsible for multiple sugar metabolism. *J Biol Chem*. 1992 Mar 5;267(7):4631-7.
36. Tao L, Sutcliffe IC, Russell RR, Ferretti JJ. Transport of sugars, including sucrose, by the msm transport system of *Streptococcus mutans*. *J Dent Res*. 1993 Oct; 72(10):1386-90.
37. Powell LV. Caries prediction: a review of the literature. *Community Dent Oral Epidemiol*. 1998 Dec;26(6):361-71.
38. Reisine S, Litt M, Tinanoff N. A biopsychosocial model to predict caries in preschool children. *Pediatr Dent*. 1994 Nov-Dec;16(6):413-8.
39. Birkeland JM, Broch L, Jorkjend L. Caries experience as predictor for caries incidence. *Community Dent Oral Epidemiol*. 1997;4:66.
40. demers M, Brodeur JM, Mouton C, et al. A multivariate model to predict caries increment in Montreal children age 5 years. *Community Dent Health*. 1992;9:373.
41. ter Pelkwijk A, van Palenstein Helderma WH, van Dijk JWE. Caries experience in the deciduous dentition as predictor for caries in the permanent dentition. *Caries Res*. 1990;24:65-71.
42. Steiner M, Helfenstein U, Marthaler TM. Dental Predictors of high Caries increment in children. *J Dent Res*. 1992;71:1926-1933.
43. Beck JD, Weintraub JA, Disney JA, Graves RC, Stamm JW, Kaste LM, Bohannon HM. University of North Carolina Caries Risk Assessment Study: comparisons of high risk prediction, any risk prediction, and any risk etiologic models. *Community Dent Oral Epidemiol*. 1992 Dec;20(6):313-21.
44. Alaluusua S, Malmivirta R. Early plaque accumulation--a sign for caries risk in young children. *Community Dent Oral Epidemiol*. 1994 Oct;22(5 Pt 1):273-6.
45. Roeters J, Burgersdijk R, Truin GJ, van 't Hof M. Dental caries and its determinants in 2-to-5-year-old children. *ASDC J Dent Child*. 1995 Nov-Dec;62(6):401-8.
46. Stewart PW, Stamm JW. Classification tree prediction models for dental caries from clinical, microbiological, and interview data. *J Dent Res*. 1991 Sep;70(9):1239-51.
47. Holbrook WP, de Soet JJ, de Graaff J. Prediction of dental caries in pre-school children. *Caries Res*. 1993;27(5):424-30.
48. Grindefjord M, Dahlöf G, Nilsson B, Modéer T. Stepwise prediction of dental caries in children up to 3.5 years of age. *Caries Res*. 1996;30:256-66.
49. Driscoll WA. The use of fluoride tablets for the prevention of dental caries. In: Forrester DJ, Schulz, EM Jr, eds. *International Workshop on Fluorides and Dental Caries Reduction*. Baltimore: University of Maryland; 1974:25-96.
50. Mellberg JR, Ripa LW. Dietary Fluoride supplementation. In: *Fluoride in Preventive*

- Dentistry-Theory and Clinical Applications*. Chicago, Ill: Quintessence;1983; 123-149.
51. Nikiforuk G. Alternatives to water fluoridation. In: *Understanding Dental Caries. Vol 2. Prevention, Basic and Clinical Aspects*. Basel, Switzerland: Karger; 1985:29-41.
  52. Murray JJ, Naylor MN. Fluorides and dental caries. In: Murray JJ, ed. *Prevention of Oral Disease*. Oxford, England: Oxford University Press; 1996.
  53. Wei SHY. Fluoride supplements. In: Newbrun E, ed. *Fluorides and Dental Caries-Contemporary Concepts for Practitioners and Students*. 3<sup>rd</sup> ed. Springfield, Ill: Charles C. Thomas; 1986:33-49.
  54. AAPD Caries Risk Assessment Tool. *Pediatric Dentistry*. Vol. 24, No. 7: 17
  55. Dorak, MT. Real-Time PCR. <http://dorakmt.tripod.com/genetics/realtime.html>  
(Accessed Sept. 2004)



APPENDIX A

**RESEARCH SUBJECT INFORMATION AND CONSENT FORM**  
**Bacteriological Samples**

**Title of Research:** Detection of *Streptococcus mutans* and *Streptococcus sobrinus* in dental plaque samples from low, moderate, and high caries risk children.

**Investigator:** Arthur P. Mourino, D.D.S.  
Virginia Commonwealth University  
School of Dentistry  
521 N. 11<sup>th</sup> Street  
Richmond, VA 23298  
(804) 828-1790 (daytime)

**Subject #** \_\_\_\_\_ **Name** \_\_\_\_\_

This consent form may contain words that you do not understand. Please ask the study doctor or the study staff to explain any words or information that you do not clearly understand. You may take home an unsigned copy of this consent form to think about or discuss with family or friends before making your decision.

This consent form is for use in a research study that involves subjects who do not have the legal capacity to consent for their participation. Accordingly, when the subject cannot legally consent to participate, the pronouns “you” and “your” should be read as referring to the subject rather than the legally authorized representative who is signing the form to give consent for the subject.

Introduction

We are trying to determine the presence of two specific types of bacteria associated with dental caries (cavities). In order to do so, we are studying persons with dental cavities. The study is directed by Dr. Arthur P. Mourino, Professor Emeritus, Department of Pediatric Dentistry in the School of Dentistry.

### Procedures

Dental caries (cavities) is caused by the bacteria (dental plaque) that grow on the surface of teeth. Dental plaque is composed almost completely of bacteria and their by-products. We wish to retrieve samples of your plaque in order to determine the presence of two different types of bacteria associated with cavities. Dental plaque will be collected by rubbing a toothpick along the outer surface of the very last tooth on each quadrant of the mouth. This will cause the plaque to cling to the toothpick. The bacteria from the plaque sample will be grown and examined in a laboratory.

### Risks, Inconveniences, and Discomforts

There is minimal discomfort associated with this procedure. It is similar to that experienced with normal tooth brushing or getting your teeth cleaned at the dental office. There are no additional risks associated with the procedure.

### Benefits

The tests are experimental (investigational) and will not be of immediate benefit to you. However, they may help us find out more about bacteria in dental plaque that cause dental cavities.

### Payment for Participation

There will not be any compensation for your time and participation in our study.

### Alternative Treatments

This is not a treatment study. You do not have to participate.

### Research-Related Injury

Virginia Commonwealth University and the VCU Health System (formerly known as Medical College of Virginia Hospitals) have no plan for providing long-term care or compensation in the event that you suffer injury as a result of your participation in this research study.

If you are injured as a result of your participation in this study, contact your study doctor immediately. Your study doctor will arrange for short-term emergency care or referral if it is necessary.

Fees for such treatment may be billed to you or to appropriate third party insurance. Your health insurance company may or may not pay for treatment of injuries as a result of your participation in this study.

### Confidentiality of Records

Confidentiality of personal information about you – including your medical records and personal research data gathered in connection with this study – will be maintained in a manner consistent with federal and state laws and regulations.

Information obtained for this study will solely be used for this study. Medical records which identify you and the consent form signed by you, may be inspected and/or copied by:

- Department of Health and Human Services (DHHS) agencies;
- Virginia Commonwealth University (VCU)

Because of the need to release information to these parties, absolute confidentiality cannot be guaranteed. Results of this study may be published, but your identity will not be revealed. Once the samples are ‘coded’, the documents which can trace the sample to the subject will be destroyed.

### Voluntary Participation/Withdrawal

Participation in this study is voluntary. You are free to withdraw at any time. If you decide to withdraw from this study, you should contact Dr. Mourino. A decision not to participate, or to withdraw, will in no way affect or jeopardize the quality of care you receive now or in the future at this institution. Dr. Mourino or other investigators involved in this study may also withdraw you without your consent for medical, dental, or administrative reasons. Any significant new findings which develop during the course of the research study which in the opinion of the study doctor may affect your willingness to continue to participate will be provided to you as soon as possible.

### Questions

If you have any questions concerning your participation in this study, contact:

Dr. Mourino or Dr. Patel at (804) 828-1790 or (804) 828-9095.

For information about research and subjects’ rights, you may contact:

Office of Research Subjects Protection  
 Virginia Commonwealth University  
 1101 E. Marshall St., Room 1-023  
 P.O. Box 980568  
 Richmond, VA 23298  
 Telephone: 804-828-0868

Do not sign this consent form unless you have had a chance to ask questions and have received satisfactory answers to all of your questions.

Consent

\_\_\_\_\_

Date

\_\_\_\_\_

Subject's Name

Signing this paper means that you have read this or had it read to you and that you want to be in the study. If you don't want to be in the study, don't sign the paper. Remember, being in the study is up to you, and no one will be mad if you don't sign this paper or even if you change your mind later.

-----

If you want to be in our study, sign your name here:

Name: \_\_\_\_\_ Date: \_\_\_\_\_

Subject's Signature (if greater than 7 and less than 18 years of age)

I have been provided with an opportunity to read this consent form carefully. All my questions about this study and my participation in it have been answered to my satisfaction.

By signing this consent form, I have not waived any of the legal rights or benefits, to which I or my child otherwise would be entitled. My signature indicates that I freely give consent to my child's voluntary participation in this research study.

\_\_\_\_\_  
Date

\_\_\_\_\_  
Legally Authorized Representative's Signature

\_\_\_\_\_  
Date

\_\_\_\_\_  
Witness' Signature

\_\_\_\_\_  
Date

\_\_\_\_\_  
Signature of person Conducting Informed Consent Discussion

\_\_\_\_\_  
Date

\_\_\_\_\_  
Investigator's Signature (if different from above)

APPENDIX BVITA

Ajit A. Patel was born on July 22, 1973 in Bombay, India and is an American citizen. He received his Bachelor of Science from University of Pittsburgh in December 1995, and his Doctor of Dental Surgery from Virginia Commonwealth University in May 2002. He is currently a postgraduate resident in the Pediatric Dentistry Program at Medical College of Virginia, Virginia Commonwealth University and upon graduation will practice in Ashburn, Virginia.