

2012

# Effect of Local Anesthesia on Postoperative Pain with General Anesthesia

Belinda Campbell

*Virginia Commonwealth University*

Follow this and additional works at: <https://scholarscompass.vcu.edu/etd>

Part of the [Dentistry Commons](#)

© The Author

---

Downloaded from

<https://scholarscompass.vcu.edu/etd/2699>

This Thesis is brought to you for free and open access by the Graduate School at VCU Scholars Compass. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of VCU Scholars Compass. For more information, please contact [libcompass@vcu.edu](mailto:libcompass@vcu.edu).

© Belinda L. Campbell 2012  
All Rights Reserve

EFFECT OF LOCAL ANESTHESIA ON POSTOPERATIVE PAIN  
WITH GENERAL ANESTHESIA

A thesis submitted in partial fulfillment of the requirements for the degree of Masters of Science  
in Dentistry at Virginia Commonwealth University.

by

BELINDA LANORE CAMPBELL, D.D.S.,  
University of Southern California School of Dentistry, 2005

Director: WILLIAM P. PISCITELLI, D.D.S  
ASSOCIATE PROFESSOR, DEPARTMENT OF PEDIATRIC DENTISTRY

Virginia Commonwealth University  
Richmond, VA  
May 2012

## Acknowledgments

Special thanks go out to Dr Malinda Husson for her innovative ideas and dedication to this project. I would like to acknowledge my thesis advisor Dr William Piscitelli for taking over this project when I needed someone. To Dr Tegwyn Brickhouse who is always there as a mentor and support for all; Dr Al Best for his patience and guidance in statistics; Drs Kordis, Berry, and Wunsch for their encouragement, support and mentorship; research assistant Sina Sadeghi for his countless follow-up calls; and to all my fellow residents who helped implement this research and collected data.

To my partner Keith and my parents Robert and Willow Campbell who constantly reminded me it could be accomplished and provided endless encouragement; and to my beautiful daughter Kerrigan Susan whose arrival was the greatest and most welcome distraction.

## Table of Contents

Abstract .....	vi
Introduction.....	1
Methods.....	7
Results.....	12
Discussion.....	24
Conclusion .....	28
References.....	29
Vita.....	32

## List of Tables

Table 1: Pain assessment across evaluators .....	13
Table 2: Correlation of raters .....	13
Table 3: Agreement between parents and patients .....	14
Table 4: Agreement between residents and patients .....	15
Table 5: Baseline comparisons.....	16
Table 6: ANCOVA results-Primary outcome: Patient post-op pain.....	17
Table 7: Mean patient post-op pain, depending upon local anesthesia and extraction .....	18
Table 8: ANCOVA follow-up results-Primary outcome: Patient post-op pain .....	18
Table 9: ANCOVA results-Secondary outcome: Parent post-op pain.....	19
Table 10: Mean parent post-op pain, depending upon local anesthesia and extraction .....	20
Table 11: ANCOVA results-Secondary outcome: Nurse post-op pain.....	20
Table 12: Mean nurse post-op pain, depending upon local anesthesia and extraction .....	21
Table 13: ANCOVA results-Secondary outcome: Patient in-home pain.....	21
Table 14: Mean patient post-discharge pain, depending upon local anesthesia and medications ..	22
Table 15: ANCOVA results-Secondary outcome: Parent in-home pain .....	22
Table 16: Mean parent post-discharge pain, depending upon local anesthesia and medications ..	23

## List of Figures

Figure 1: Wong Baker Faces Pain Scale: Visual Scale for numerical pain intensity evaluation. ....8

Figure 2: Agreement between parents and patients .....14

## Abstract

### EFFECT OF LOCAL ANESTHESIA ON POSTOPERATIVE PAIN WITH GENERAL ANESTHESIA

By Belinda Lanore Campbell, D.D.S

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in Dentistry at Virginia Commonwealth University.

Virginia Commonwealth University, 2012

Major Director: William P. Piscitelli, D.D.S.  
Associate Professor, Department of Pediatric Dentistry

**Purpose:** The aim of this study was to determine if the use of local anesthesia with general anesthesia results in less postoperative pain. The alternative hypothesis is that children will experience less postoperative discomfort when utilizing intraligamental local anesthetic during the intra-operative time period.

**Methods:** Patients were recruited for this single blind, randomized, prospective cohort study with the following inclusion criteria: English speaking children age 3-6 years, ASA I/II requiring general anesthesia for dental treatment. Randomization was done to place patients in groups of no local anesthetic vs. local anesthetic administration. A Wong-Baker Faces Pain Scale was utilized to evaluate pre-operative and postoperative pain. Data were compared using a two way mixed model ANCOVA controlling for sex, ethnicity, pre-op pain, and intra-op meds given.

**Results:** Data was collected and evaluated on 90 patients. There was a statistically significant



difference in postoperative pain for patients who received extractions without local anesthesia vs. those with local anesthetic. There was no statistically significant difference in pain outcomes based solely on whether local anesthetic was administered regardless of treatment type.

**Conclusions:** The outcome of this study shows evidence for provision of local anesthetic during general anesthesia in patients receiving extractions to reduce postoperative pain.

## **Introduction**

The International Association for the Study of Pain's widely used definition states, "Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage". Pain is described as a very subjective feeling related to experience in early life and is influenced by such factors as age, fear, personality, circumstances and culture.<sup>1</sup> A well documented phenomenon in medicine is the under-treatment of pain in children.<sup>2</sup> The American Academy of Pediatrics and the American Pain Society issued joint recommendations in 2001 regarding the role of the pediatrician to ensure effective treatment of pain in infants, children, and adolescents but does not comment on pediatric dentists and their role in alleviating discomfort for their patients.<sup>3</sup>

The Centers for Disease Control and Prevention reports that caries is the most prevalent infectious disease in our nation's children. More than 40% of children have caries by the time they reach kindergarten. In contrast to declining prevalence of dental caries in older age groups, the prevalence of caries in poor US children under the age of 5 is increasing.<sup>4</sup> This increase in caries has led to full-mouth dental rehabilitation under general anesthesia (GA) becoming an increasingly necessary and accepted modality of treatment for comprehensive pediatric dental treatment.<sup>5</sup> GA may be indicated for children with the following: extreme anxiety, extensive needs, very young age, and/or physical/mental disabilities.<sup>6</sup> The chief advantage of GA is that it facilitates completion of all necessary dental care in a single visit with minimal duress to the

patient. Dental treatment under GA is usually the last resort due to expense, risk-benefit considerations, and acceptability to parent. Minimizing morbidity is a necessity to ensure acceptability of this treatment modality.

Morbidity related to GA is less of a problem for patients than morbidity related to dentistry.<sup>7</sup> Postoperative dental pain is a common occurrence in patients undergoing general anesthesia for dental procedures.<sup>1,7-9</sup> Studies have shown reports of postoperative pain ranging from 57.5-95% of patients.<sup>1,9</sup> Some pain is common within the first few hours after surgery and thought to be secondary to the trauma to hard and soft tissues. Uncontrolled pain can delay discharge from the hospital. Studies investigating the use of oral preoperative analgesics, i.e. ibuprofen and paracetamol, suggest that these medications provide no greater benefit to postoperative pain than placebo.<sup>10</sup> Improvements in pain control have the potential to significantly reduce reported morbidity following dental treatment under GA.

The concomitant use of local anesthetics is one aspect of outpatient dental treatment under GA that has not received specific recommendations in terms of efficacy, purpose and safety and is still not always part of routine clinical practice.<sup>5</sup> Currently, the American Society of Anesthesiologists and American Dental Association have no recommendations regarding the use of local anesthetics during dental rehabilitation under GA.<sup>11,12</sup> The American Academy of Pediatric Dentistry guidelines state that local anesthesia “may be used” to reduce pain in the postoperative recovery period after GA; and reduce the dosage of inhalation anesthetics required during GA. No directive statements exist in the AAPD guidelines regarding the use of local anesthesia (LA).<sup>13</sup> As evidence-based dentistry has become the standard of care, examination of

the impact of local and general anesthetics on pediatric pain is not only clinically relevant but also necessary in order for acceptable treatment to be rendered.

Local anesthesia has been used in conjunction with general anesthesia to reduce postoperative pain in a variety of other surgical procedures, i.e. joint replacements. Some operators use local anesthetic as an adjunct to GA for its associated vasoconstriction. While this could reduce bleeding, it could also increase post-op distress due to associated facial numbness. Prior research on effectiveness of LA in young children has been inconclusive. Gazal et al. investigated bupivacaine soaked swabs as an alternative topical method of pain control to avoid postoperative facial numbness. The results indicated that they were ineffective at reducing pain.<sup>14</sup> Numerous other studies regarding the use of injected local anesthetic have shown no differences in postoperative pain when a of local anesthetic was used in conjunction with GA.<sup>2,8,15</sup> A study of intraligamental local anesthetic by Sammons et al. showed less pain initially after recovery, but that the difference was not sustained over the first hour after dental extraction.<sup>16</sup> There is evidence for the use of local anesthetic for reduction in intra-operative hemorrhage.<sup>15,17</sup>

The intraoral infiltration of a local anesthetic causes profound alteration of orofacial sensation, particularly affecting the lips and cheeks. Older children can be counseled preoperatively to expect numbness of the lips or gums when they wake up from general anesthesia, but younger children are often incapable of understanding this. In many cases, general anesthesia is indicated in children because of anxiety, behavioral issues, or extent of treatment required. Many of these children do not attend regular dental appointments and therefore may not have previous experience with the altered sensation of the oral cavity associated with a local anesthetic. Many

providers note that younger children sometimes appear to be as distressed by the feeling of numbness as by the postoperative pain of the procedure.

The use of local anesthesia may also lead to inadvertent lip biting.<sup>2,18</sup> A prospective study published in 2000 found that 13% of children aged 2 to 18 experienced soft tissue trauma after unilateral or bilateral mandibular nerve block anesthesia.<sup>19</sup> Predictably, the evidence of soft tissue trauma was highest among the youngest age groups- 18% among children less than 4 years, 16% in children aged 4 to 7, 13% in 8 to 11 year old children, and 7% in children 12 year and older.<sup>19</sup> Children may bite their lower lip, either out of curiosity associated with the unfamiliar sensation of being numb or inadvertently because no pain is felt. Inadvertent lip biting can also occur during the immediate postoperative period when eating or sleeping.

One method to overcome this distress and possible trauma is to abandon the use of local anesthesia altogether. Alternatively, LA could be administered using intraligamental injection as there is less associated soft tissue numbness.<sup>17,20</sup> Intraligamental administration does not produce anesthesia of the lip or tongue and, therefore, eliminates the risk of self-damage in children during the immediate postoperative period of anesthesia. Leong et al. showed that intraligamental injection resulted in significantly lower pain scores during the first night postoperative compared to local infiltration of LA. The researchers reasoned that the after-effect of less soft tissue numbness initially could be better tolerated with reduced perceived pain/discomfort and thus, showed more favorable outcomes later.<sup>17</sup>

Numerous formulations of anesthetic are available for dental use. Most of the anesthetic

solutions commonly used contain a vasoconstrictor in order to prolong the anesthetic effect. Vasoconstrictors have been shown to interfere with pulpal blood flow.<sup>21</sup> It has been found that infiltration injection of 2% lidocaine with 1:100,000 epinephrine reduces pulpal blood flow to as low as 28% and the recovery of flow may take up to 75 minutes.<sup>22</sup> During this time period, the dental pulp may experience decreased oxygenation which may contribute to an irreversible injury. Other studies have shown that at higher doses, local injections of epinephrine causes pulpal blood vessels to collapse, producing total ischemia of the pulp.<sup>21</sup> The individual effect of epinephrine in attaining anesthesia was investigated by Handler and Albers in their study of intraligamental anesthesia. They found no significant differences between 2% lidocaine, 2% lidocaine with epinephrine 1:50,000, 2% lidocaine with epinephrine 1:100,000 with regard to occurrence or duration of pulpal anesthesia.<sup>23</sup> In a clinical trial, Edwards et al. successfully anesthetized and removed 79% of permanent teeth with use of 2% lidocaine administered by intraligamental injection.<sup>24</sup>

Since a primary goal of intraoperative local anesthesia is the reduction of postoperative distress, it is important to determine if local anesthesia actually accomplishes this or whether it would be better to omit this practice in young children. Observational studies in the community dental clinic highlight that local anesthetic injections appear superior to systemic analgesia, and patients who received local anesthetic injections seem more settled in recovery.<sup>25</sup> Ashkenazi et al. evaluated postoperative dental pain and determined that root canal treatment and preformed stainless steel crowns, with or without pulpotomy, induced a significantly higher incidence of postoperative dental pain compared to extraction, restorations, and sealant.<sup>26</sup> Other studies have indicated that extractions are the most invasive procedure and that children undergoing

extractions were 7 times more likely to report pain after returning home.<sup>9</sup>

The aim of this study was to determine if the use of local anesthesia utilizing the intraligamental injection results in less postoperative pain in patients undergoing GA. Intraligamental injections concurrently eliminate soft tissue anesthesia and the possible confounding factor of unusual sensation which may lead to distress. The correlation between pain scores provided by patients and the evaluation of their pain by providers and parents to determine if outsiders can accurately evaluate pain in a young child was also evaluated. If the hypothesis is correct, this study will give evidence-based recommendations for use of local anesthesia when providing full mouth dental rehabilitation under GA in terms of reduction of postoperative pain.

## **Methods**

All patients included in this single blind, randomized, prospective cohort pilot study were seen on an outpatient basis at Virginia Commonwealth University Ambulatory Care Center. The VCU Institutional Review Board (IRB) for Humans Subjects Protection approved this study. Informed consent was obtained from the parent on the day of surgery during the pre-surgical consultation at the Ambulatory Care Center. Patients seen at the Ambulatory Care Center were scheduled for care under general anesthesia based on the pre-cooperative/uncooperative behavior and/or amount of treatment needed. The inclusion criteria for this study were as follows: English speaking ASA 1 and 2 children with restorative needs in primary dentition only, 3-6 years of age, predetermined to require general anesthesia care for dental treatment and/or extractions. A prior pilot study determined that 84 patients would be required to have 80% power to determine a difference between groups. A total of 90 patients were recruited and data collected.

Prior to the commencement of this study, Post Anesthesia Care Unit (PACU) nurses and pediatric dental residents were formally calibrated prior to their participation in this study. All instructions, informed consent, and IRB paperwork were reviewed prior to this study allowing practitioners and nurses to participate. There were two groups with a total of n=90 patients. N=46 patients were randomized to receive local anesthesia and n=44 were in the no local anesthetic control group. Each of these groups were then treated either with or without local anesthesia using intraligamental lidocaine, as previously randomized using the Statistical Analysis System (SAS)



randomization technique prior to starting the study. During data analysis the groups were further divided into a first group of children receiving dental restorations only for primary teeth, and a second group of children receiving the combination of dental restorations and dental extractions of primary teeth. Each number from 1-90 was assigned a random value of local anesthesia or no local anesthesia. Each child participating received one of the pre-numbered and randomized packets. The children not receiving local anesthetic served as the control group for this study.

After receiving informed consent, the child, parent and the resident in the preoperative assessment area evaluated and rated the patient's preoperative pain utilizing the Wong-Baker Faces Pain Scale (Figure 1), prior to the start of anesthesia care. The Wong-Baker Faces pain scale consists of six cartoon faces with varying expressions ranging from very happy to very sad.<sup>27</sup> The six different faces with associated numbers are on an ordinal continuous value scale ranging from 0 (no hurt) to 10 (hurts worst). Three preoperative baseline pain scores were recorded at the pre-operative assessment time.



*Figure 1: Wong Baker Faces Pain Scale: Visual Scale for numerical pain intensity evaluation.*

The study used a standardized anesthetic regimen, as deemed appropriate by the consulting pediatric anesthesiologist. The anesthesia protocol included, pre-operative oral midazolam at 0.5

milligrams (mg) per kilogram (kg) up to 20 mg total, mask induction with sevoflurane/oxygen/nitrous oxide, induction medications such as fentanyl (narcotic) 0.5-1.0 micrograms per kilogram and propofol at 2 mg/kg. It was requested that no additional pain medications (narcotics) be administered throughout the intra-operative time period unless found to be medically necessary by the anesthesia team (interventions were recorded).

Subjects were randomly assigned to either receive LA or no LA. The pediatric dental resident opened the pre-randomized sealed envelope with the corresponding number and value of LA or no LA after consent had been obtained. Subjects assigned to the LA group received a standardized LA protocol as follows: 2% plain lidocaine administered in the first quadrant to be treated, after placement of a gauze throat pack, and prior to the start of the procedure. Operators used intraligamental injections of the 2% lidocaine plain with a 3mL syringe and a 30 gauge extra-short needle. Previous literature shows evidence of pulpal ischemia with use of epinephrine containing anesthetics.<sup>21,22</sup> To eliminate any possible confounding pain due to ischemia by use of a vasoconstrictor, a plain formulation of lidocaine was selected. The local anesthetic was administered in two locations for each single rooted tooth (buccal and lingual), and four locations for each multi-rooted tooth treated (mesial buccal, distal buccal, mesial lingual and distal lingual). The operators did not exceed doses of 4.4mg/kg total of 2 % plain lidocaine. The total amount and time of administration of the local anesthesia was recorded in the anesthesia record. Treatment in each quadrant was completed in the following order: sealants, anesthesia administration, extractions, composite resins, pulpotomies, and stainless steel crowns. Treatment was completed by quadrants with administration of local anesthetic occurring just prior to beginning treatment in the subsequent quadrants.

Following completion of the dental treatment and general anesthesia care, the patients were escorted to the PACU. The PACU nurses, patients, and their parents were blinded as to whether or not the child had received a local anesthetic. Three pain scores were obtained as follows: patients, PACU nurse and the parent subjectively graded the child's pain intensity in the immediate postoperative time period, using the visual Wong-Baker pain scale.. Additional pain medications administered in the PACU were recorded if needed in the immediate postoperative time period prior to discharge. The patients were contacted 6-8 hours following their procedure. The parents evaluated their pain at this time, utilizing the Wong-Baker Faces Pain Scale which was sent home along with routine postoperative instructions. The research assistant, who was blinded as to whether or not the patient received LA, recorded the pain measurement for future review. If parents reported that pain medications were administered at home, then that amount was recorded.

A total of eight pain scores were recorded for each patient: three preoperative (patient, parent, and pediatric dental resident obtaining consent), three in the immediate postoperative time period (patient, PACU nurse, and parent), and two (patient and parent) 6-8 hours postoperative.

Outcome variables were self reported patient postoperative pain scores, and the control variables included pre-operative pain score, treatment type, and the need for intra-operative interventions.

ANCOVA controlling for pre-operative reported pain scores, treatment completed, and the need for intra-operative medications was used for data analysis. Primary independent variable comparison was made between the local anesthetic and no local anesthetic experimental groups.

Agreement between raters was assessed by the calculation of Cohen's Kappa, a change-corrected measure of agreement.

## **Results**

The overview of the results section is as follows: The first portion of results describes the level of agreement between the different individual's rating of pain. The second section addresses the primary aim, whether local anesthesia use has an effect on the patient rating of post-op pain. The final portions address the secondary analyses that consider other factors that may be related to pain, other raters of pain, and the in-home follow-up rating.

### **Pain ratings: Descriptive**

There were N = 674 pain ratings given by the four evaluators (parent, patient, resident, and nurse), across the three occasions (pre-op, post-op, and home). Residents only gave pain assessments at pre-op and nurses only gave assessments at post-op. About two-thirds of the time, pain was rated at 0= "no hurt" (Table 1) and pain at the 10= "hurts worst" did occur.

*Table 1: Pain assessment across evaluators*

Pain	Frequency				%
	Parent	Patient	Resident	Nurse	
0	153	151	80	61	66.0
1	5	3	2		1.5
2	33	37	6	7	12.3
3	6	2		2	1.5
4	22	18	2	5	7.0
5	5	4			1.3
6	5	9		5	2.8
7	2	1		1	0.6
8	6	9		4	2.8
9	2				0.3
10	8	13		5	3.9
n	247	247	90	90	100
Mean	1.59	1.77	0.24	1.77	
SD	2.61	2.87	0.77	3.05	

Abbreviation: n = frequency, SD = standard deviation

Since multiple raters assessed pain intensity at the same time, the level of agreement between raters may be described. Overall there is good agreement between the parent and patient ratings ( $r = 0.67$ , Table 2).

*Table 2: Correlation of raters*

Rater	Rater		
	Patient*	Resident	Nurse
Parent	0.67	0.45	0.55
Patient		0.49	0.34

Correlations constrained to have a zero intercept. \* n = 247 paired ratings. All other paired ratings are n = 90.

### **Agreement between Parents and Patients**

There were n = 247 occasions where both parents and patients rated the level of pain and on 128 of those occasions, both raters gave a rating of 0 (Table 3). There was no mean difference between the mean levels of pain (paired t-test  $P = 0.1705$ ). In 66% of all cases, both raters agreed (shown in bold in the table; chance corrected Kappa = 0.42). In 89% of all cases the ratings differed by 3 or less. This level of agreement may also be seen in Figure 2, where the sizes of the

circles are proportional to the frequency of occurrence. There were few cases ( $n = 15$ ) where raters disagreed by 5 or more and the maximum disagreement was 8.

Table 3: Agreement between parents and patients

Patient	Parent										
	0	1	2	3	4	5	6	7	8	9	10
0	<b>128</b>	2	12	1	5			1	2		151
1		<b>2</b>	1								3
2	17	1	<b>11</b>	4	1		2			1	37
3			1	<b>1</b>							2
4	3		3		<b>11</b>	1					18
5	1				1	<b>1</b>			1		4
6	3				2	2	<b>1</b>			1	9
7								<b>1</b>			1
8	1		2		1	1	1		<b>1</b>		2
9											0
10			3		1		1		2		<b>6</b>
	153	5	33	6	22	5	5	2	6	2	8
											247




Figure 2: Agreement between parents and patients

### Agreement between Residents and Patients

During pre-op there were  $n = 90$  occasions where both residents and patients rated the level of pain. For 58 patients, both raters gave a rating of 0 (Table 4). There was a mean difference between the mean levels of pain (paired t-test  $P < .0001$ ). The patients recorded significantly

more pain than did the residents (mean = 1.23 versus 0.244). In 66% of all cases, both raters agreed (shown in bold in the table; chance corrected Kappa = 0.11; which was not beyond a chance level of agreement,  $P = 0.0757$ ). In 87% of all cases the ratings differed by 2 or less. There were few cases ( $n = 5$ ) where raters disagreed by 5 or more and the maximum disagreement was 8.

*Table 4: Agreement between residents and patients*

Patient	Resident										
	0	1	2	3	4	5	6	7	8	9	10
0	<b>58</b>	1	2								61
1											
2	14	1	<b>0</b>								15
3											
4	4		1		<b>1</b>						6
5	1		0		0						1
6	1		2		0						3
7											
8	2				1						3
9											
10			1								1
	80	2	6		2						90

### **Agreement between Nurses and Patients**

During post-op, there were  $n = 90$  occasions where both the recovery nurse and the patient rated the level of pain.

### **Baseline comparisons**

At pre-op there was no difference in the patient's reported pain depending upon race, resident year, whether an extraction was to be performed, whether a restoration was to be performed, or the age of the patient (Table 5). The average age of the patient at the time of surgery was 4.5 years ( $SD = 1.02$ , range = 3.0 to 6.9 years). There also was no pre-op pain difference depending upon the randomly assigned local anesthesia groups or whether an intra-op pain medication was



used.

Table 5: Baseline comparisons

Characteristics	Patient Pre-Op Pain			
	n	Mean	SE	p-value
<b>Race/Ethnicity</b>				0.759
Caucasian	43	1.233	0.352	
African American	36	1.389	0.384	
Hispanic	4	0.500	1.153	
<b>Resident year</b>				0.607
1	33	1.394	0.390	
2	57	1.140	0.297	
<b>Extraction</b>				0.620
No	30	1.067	0.410	
Yes	60	1.317	0.290	
<b>Restoration</b>				0.582
No	1	0.000	2.243	
Yes	89	1.247	0.238	
<b>Age (years)</b>				0.515
			r =	
	90	1.233	0.07	
<b>Local anesthesia</b>				0.335
Yes	46	1.457	0.329	
No	44	1.000	0.337	
<b>Intra-op pain medication</b>				0.316
No	33	1.545	0.389	
Yes	57	1.053	0.296	

### Outcome analysis

Patients were randomly allocated to local anesthesia groups (Yes = 46, No = 44) and then during the surgical process it was determined whether an extraction would be done. It was also documented whether intra-op pain medication was administered. In the local anesthesia groups, there were an approximately equal number of patients with extractions (34/46 in the local anesthesia group, and 26/44 in the no local anesthesia group). There was also an approximately equal number of patients receiving intra-op pain medication (27/46 in the local anesthesia group, and 30/44 in the no local anesthesia group). The results of the primary outcome analysis are

shown in Table 6. There was weak evidence for a relationship between the use of intra-op pain medications and patient post-op pain ( $P < 0.07$ ) and weak evidence for a correlation between pre-op pain and post-op pain ( $P < 0.08$ ). The interaction test indicated that the effect of local anesthesia was not consistent across the two extraction groups ( $P = 0.019$ ). So, an interpretation of the main-effects of local anesthesia and of extraction is not appropriate.

*Table 6: ANCOVA results-Primary outcome: Patient post-op pain*

Source	df	F	p-value
Intra-op pain meds	1	3.413	0.068
Pre-op pain	1	3.334	0.071
Extraction	1	0.682	0.411
Local anesthesia	1	0.349	0.557
Extract*Local	1	5.723	0.019
Error	84		

N = 90, R<sup>2</sup> = 15%

The effect of local anesthesia on patient post-op pain is seen in Table 7. Within the no extraction group, there was no significant difference depending upon the use of local anesthesia ( $P > 0.2$ ) but in the extraction group, there was a significant difference in pain ( $P < 0.02$ ) with the local anesthesia group showing less pain (mean = 1.6 vs. 3.9). The 95% CI on the difference indicates that when there is an extraction, the use of local anesthesia reduces pain by between 0.6 and 4.0 units.

*Table 7: Mean patient post-op pain, depending upon local anesthesia and extraction*

Local	Post-op Pain				
	LS Mean	SE	95% CI	p-value	
No extraction					
Yes	2.819	0.970	0.889	4.748	
No	1.432	0.789	-0.136	3.000	
difference	1.387	1.249	-1.062	3.836	0.270
Extraction					
Yes	1.609	0.574	0.468	2.750	
No	3.895	0.683	2.537	5.252	
difference	-2.286	0.882	-4.014	-0.558	0.011

Least-squares mean estimates and p-values calculated from an ANCOVA with the following

effects: Intra-op pain meds (yes/no), pre-op pain, extraction (yes/no), local anesthesia (yes/no), extraction\*local interaction.

The follow-up analysis of this primary outcome also explored the effect of age (in months), number of interventions, restoration, patient race, and resident year. Since race was not reported by 7 patients and since the preliminary analysis indicated no evidence for an effect of race ( $P = 0.529$ ), race was not included in subsequent analyses. The result of this follow-up analysis is shown in Table 8. As may be seen, none of the additional factors were related to patient post-op pain and the significance of the factors in the primary analysis did not change materially.

*Table 8: ANCOVA follow-up results-Primary outcome: Patient post-op pain*

Source	df	F	p-value
Intra-op pain meds	1	2.032	0.158
Pre-op pain	1	3.472	0.066
Extraction	1	0.552	0.460
Local anesthesia	1	0.328	0.569
Extract*Local	1	6.128	0.015
Patient age	1	0.062	0.804
#interventions	1	0.248	0.620
Restoration	1	0.066	0.797
Resident Year	1	2.443	0.122
Error	80		

N = 90,  $R^2 = 17.6\%$

### **Parent’s pain rating post-op,**

In these follow-up analyses, since none of the exploratory factors was significant, the analysis model will be the same as that used for the primary outcome. The results of the analysis of the parent’s pain rating immediately post-op is shown in Table 9. The pattern of results is similar to that found when analyzing the patient pain score in that there was a non-significant interaction effect ( $P < 0.06$ ).

Table 9: ANCOVA results-Secondary outcome: Parent post-op pain

Source	df	F	p-value
Intra-op pain meds	1	5.65	0.020
Pre-op pain	1	0.23	0.631
Extraction	1	0.36	0.550
Local anesthesia	1	0.05	0.821
Extract*Local	1	3.69	0.058
Error	84		

N = 90, R<sup>2</sup> = 10%

The average parent post-op pain score, after controlling the effects of intra-op pain med, and pre-op pain is shown in Table 10. Within the no extraction group, there was no significant difference depending upon the use of local anesthesia ( $P > 0.3$ ) and in the extraction group, there was a non-significant difference in pain ( $P > 0.06$ ) with the local anesthesia group showing nominally less pain (mean = 2.3 vs. 4.0). The 95% CI on the difference indicates that when there is an extraction, the use of local anesthesia reduces pain by between  $-0.06$  and  $+3.5$  units.

Table 10: Mean parent post-op pain, depending upon local anesthesia and extraction

Local	Post-op Pain				p-value
	LS Mean	SE	95% CI		
	No extraction				
Yes	3.327	1.019	1.301	5.353	
No	1.973	0.804	0.375	3.571	
difference	1.354	1.299	-1.192	3.900	0.300
	Extraction				
Yes	2.266	0.586	1.101	3.432	
No	3.976	0.698	2.588	5.364	
difference	-1.710	0.901	-3.476	0.057	0.061

Estimates and p-values calculated from an ANCOVA with the following effects: Intra-op pain meds (yes/no), parent rating of pre-op pain, extraction (yes/no), local anesthesia (yes/no), extraction\*local interaction.

### Nurse's pain rating post-op,

The analysis of the nurse's pain rating used the same model as for the primary analysis, including

the patient's pre-op pain rating (since the nurse did not rate pain pre-op). The results are shown in Table 11. Again, there was a marginally non-significant interaction ( $P = 0.053$ ) indicating that the effect of a local anesthesia may depend upon whether an extraction is performed.

*Table 11: ANCOVA results-Secondary outcome: Nurse post-op pain*

Source	df	F	p-value
Intra-op pain meds	1	0.00	0.959
Pre-op pain	1	3.67	0.059
Extraction	1	3.02	0.086
Local anesthesia	1	1.02	0.315
Extract*Local	1	3.84	0.053
Error	84		

N = 90, R<sup>2</sup> = 14%

The table of the mean nurse rating of post-op pain (Table 12) shows a similar pattern except that in this instance, in the extraction group the effect of a local anesthetic is significant ( $P < .02$ ).

*Table 12: Mean nurse post-op pain, depending upon local anesthesia and extraction*

Local	Post-op Pain				p-value
	LS Mean	SE	95% CI		
	No extraction				
Yes	1.434	0.850	-0.255	3.124	
No	0.791	0.690	-0.582	2.164	
difference	0.643	1.094	-1.501	2.787	0.558
	Extraction				
Yes	1.270	0.502	0.271	2.269	
No	3.261	0.598	2.073	4.450	
difference	-1.992	0.772	-3.504	-0.479	0.012

Estimates and p-values calculated from an ANCOVA with the following effects: Intra-op pain meds (yes/no), patient rating of pre-op pain, extraction (yes/no), local anesthesia (yes/no), extraction\*local interaction.

### **Patient's pain rating 6-8 hours post-op**

Not all of the subjects provided in-home 6-8 hour post-op pain ratings. N = 67 subjects did, n = 33 in the local=yes group and n = 34 in the local=no group. An additional factor to consider in the analysis of these data is the use of post-discharge pain medications. These medications were used in 54% of the cases (n = 36). Additionally, the analysis also added the immediate post-op

pain rating as a covariate instead of the pre-op pain rating. The results of the analysis are shown in Table 13. The only significant difference was for post-op pain meds ( $P < .03$ ).

*Table 13: ANCOVA results-Secondary outcome: Patient in-home pain*

Source	df	F	p-value
Intra-op pain meds	1	0.04	0.838
Post-op pain	1	0.39	0.536
Extraction	1	1.65	0.204
Local anesthesia	1	1.18	0.282
Extract*Local	1	0.08	0.773
post-op pain meds	1	5.02	0.029
Error	84		

N = 67, R<sup>2</sup> = 15%

The differences associated with the use of post-discharge pain meds is shown in the upper portion of Table 14. Those using pain meds reported more pain ( $P < .02$ ). Additionally, the table also shows the non significant difference depending upon the use of local anesthesia.

*Table 14: Mean patient post-discharge pain, depending upon local anesthesia and medications*

groups	Post-discharge Pain				p-value
	LS Mean	SE	95% CI		
	Post-discharge pain meds				
Yes	2.022	0.333	1.356	2.687	
No	0.928	0.342	0.243	1.613	
difference	1.094	0.448	0.198	1.989	0.018
	Local anesthesia				
Yes	1.290	0.541	0.207	2.372	
No	2.045	0.452	1.141	2.949	
difference	-0.756	0.697	-2.121	0.610	0.282

Estimates and p-values calculated from an ANCOVA with the following effects: Intra-op pain meds (yes/no), patient rating of post-op pain, extraction (yes/no), local anesthesia (yes/no), extraction\*local interaction, and the use of post-op pain meds (yes/no).

### **Parent's pain rating 6-8 hours post-op**

Using the same model as the patient's in-home pain rating, the results are shown in Table 15. The only significant effect was for post-discharge pain meds ( $P < .02$ ).

Table 15: ANCOVA results-Secondary outcome: Parent in-home pain

Source	df	F	p-value
Intra-op pain meds	1	0.00	0.977
Post-op pain	1	0.05	0.816
Extraction	1	1.01	0.319
Local anesthesia	1	1.03	0.315
Extract*Local	1	0.95	0.335
post-op pain meds	1	5.97	0.018
Error	84		

N = 67, R<sup>2</sup> = 17%

Table 16 shows the mean post-discharge (in-home) pain rating as given by the parent depending upon the use of post-discharge pain meds. Those who did use pain meds had significantly more pain ( $P < .02$ ). Additionally, the results of the primary research question—the effect of local anesthesia—is shown in the bottom portion of the table. There was no significant difference ( $P > 0.3$ ).

Table 16: Mean parent post-discharge pain, depending upon local anesthesia and medications

groups	Post-discharge Pain				p-value
	LS Mean	SE	95% CI		
	Post-discharge pain meds				
Yes	2.022	0.333	1.356	2.687	
No	0.928	0.342	0.243	1.613	
difference	1.094	0.448	0.198	1.989	0.018
	Local anesthesia				
Yes	1.225	0.383	0.458	1.992	
No	1.725	0.320	1.084	2.366	
difference	-0.500	0.494	-1.467	0.468	0.315

Estimates and p-values calculated from an ANCOVA with the following effects: Intra-op pain meds (yes/no), parent rating of post-op pain, extraction (yes/no), local anesthesia (yes/no), extraction\*local interaction, and the use of post-op pain meds (yes/no).

## Discussion

Our ability to manage postoperative pain is a vital component of successful dental care for pediatric patients. Managing pain first requires accurately assessing the degree of pain experienced. Based on data collected during this study it can be concluded that parental account of pain intensity for their child can be fairly reliable. There was overall good agreement between pain scores reported by patients and those of their parents ( $r=0.67$ ). On the contrary, residents and nurses rated patient pain to be lower than that described by the patients themselves. Residents recorded significantly lower pain than patients ( $p<0.0001$ ). This may be due to providers extensive exposure to patients and therefore loss of sensitivity to signals that express pain. It would be appropriate to assume a parent would be in touch with signs and signals that their child is experiencing discomfort that outsiders may not be tuned into.

The primary aim of this study was to determine if administration of LA utilizing the intraligamental technique resulted in lower postoperative pain. Analysis of the data reveals that LA administration does not result in lower postoperative pain. However, when evaluation of treatment type occurred there effect of local anesthesia was not consistent across extraction groups. Patient who received LA for extractions had significantly less pain than patients who did not receive LA for extractions ( $p<0.02$ ). In other words, administration of LA for treatment that did not include extractions resulted in non-significant postoperative pain differences ( $p>0.5$ ). Concurrently, if a patient received any extractions as part of their treatment plan, the



administration of LA decreased postoperative pain by between 0.6 and 4.0 units.

Secondary analyses were completed to look at the effect of pre-operative pain levels and use of intra-operative pain medications on postoperative pain. There is weak evidence that patients with higher pre-operative pain levels subsequently have higher postoperative pain levels ( $p < 0.08$ ). The anesthesia providers were asked to not administer additional narcotics during the intra-operative period unless deemed necessary. Narcotic administration past the point of induction was termed an intervention. There appears to be a possible relationship between necessity for additional intra-operative narcotic administration and postoperative pain ( $p < 0.07$ ). Patients who received medication had higher levels of postoperative pain.

Patients were contacted at home the evening of their surgery to determine pain levels and whether the parent at had administered pain medication at home. Patients who had been given pain medications at home had reported significantly higher pain levels ( $p < .03$ ). There was no difference in at home pain levels between the group that received LA and the no LA group. The local anesthetic administered intra-operatively would no longer be present 6-8 hours postoperative when the patients were contacted at home and therefore should not have an effect on levels of pain.

There have been numerous studies on effects of local anesthetic on pain outcomes. It is important to be able to evaluate critically the evidence that is presented to determine clinical relevance to practitioners. A weakness to the current study was the use of plain lidocaine. The majority of pediatric dentists use 2% lidocaine with 1:100,000 epinephrine or 4% septocaine with

1:100,000 epinephrine. The addition of a vasoconstrictor prolongs the period of pulpal anesthesia and therefore may increase effectiveness. Even without the presence of a vasoconstrictor there was significant pain reduction in patients undergoing extractions in this study. The anesthetic solution was administered using standard anesthetic syringes instead of a specially designed intraligamental injection syringe. The standard syringe is readily available for all providers but does not allow for pressurized injection into the ligamental space and therefore does not deliver a standardized dose of anesthetic.

There are many strengths to the current research including the delivery of anesthetic to 4 locations in posterior teeth to ensure thorough anesthesia. Past research administered intraligamental anesthetic into the mesial-buccal root only<sup>17</sup> or did not indicate location of anesthetic administration.<sup>16</sup> There was a uniform order to the procedures completed after administration of local anesthetic. Extractions were completed immediately after administration of local anesthetic. Pulpal anesthesia should have been present for extractions but may have worn off prior to completion of all other restorative treatment leading to the non-significant effect of local anesthesia for restorative work alone.

Future studies in the area of local anesthesia usage during full mouth dental rehabilitation with general anesthesia should evaluate commonly used anesthetic formulations including 2% lidocaine with 1:100,000 epinephrine and/or 4% septocaine with 1:100,000 epinephrine delivered by infiltration and/or nerve blocks to determine if overall postoperative pain can be reduced. The current study was able to show evidence for reducing postoperative pain in patients undergoing extractions but does not give recommendations for local anesthetic usage with

treatment that does not include extractions.

## **Conclusion**

1. Use of intraligamental 2% lidocaine can significantly decrease postoperative pain with general anesthesia when extractions are performed.
2. Use of local anesthesia does not significantly reduce pain with treatment not including extractions when utilizing the intraligamental technique.
3. Patients with higher preoperative pain had a tendency for higher postoperative pain.
4. Parents give more accurate accounts of the children's pain than other providers.
5. Patients who receive pain medications at home had higher pain levels.

The use of the intraligamental technique of anesthetic administration provides an effective means of anesthetizing for extractions during general anesthesia care without the unwanted soft tissue numbness that coincides with other techniques.

## References

1. Fung DE, Cooper DJ, Barnard KM, Smith PB. Pain reported by children after dental extractions under general anesthesia. A pilot study. *Int J Pediatr Dent* 1993; 3:23-8.
2. Townsend JA, Ganzberg S, Thikkurissy S. The effect of local anesthetic on quality of recovery characteristics following dental rehabilitation under general anesthesia in children. *Anesth Prog* 2009; 56:115-122.
3. American Pain Society. The assessment and management of acute pain in infants, children, and adolescents. Available at: "<http://www.ampainsoc.org/advocacy/pediatric2.htm>". Accessed November 27, 2010.
4. US Dept of Health and Human Services. Oral Health in American: A report of the Surgeon General. Rockville, MD: US Dept of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health; 2000.
5. Watts A, Thikkurissy S, Smiley M, McTigue D, Smith T. Local anesthesia effects physiologic parameters and reduces anesthesiologist intervention in children undergoing general anesthesia for dental rehabilitation. *Pediatr Dent* 2009; 31:414-9.
6. Ersin NK, Oncag O, Cogulu D, Cicek S, Balcioglu ST, Cokmez B. Postoperative morbidities following dental care under day-stay general anesthesia in intellectually disabled children. *Journal of Oral and Maxillofacial Surgery* 2005 December; 63(12):1731-36.
7. Atan S, Ashley P, Gilthorpe MS, Scheer B, Mason C, Roberts G. Morbidity following dental treatment of children under intubation general anesthesia in a day-stay unit. *Int J Paediatr Dent* 2004; 14:9-16.
8. Coulthard P, Rolfe S, Mackie IC, Gazal G, Morton M, Jackson-Leech D. Intraoperative local anesthesia for paediatric postoperative oral surgery pain-a randomized controlled trial. *Int. J. Oral Maxillofac. Surg.* 2006; 35: 1114-19.
9. Needleman HL, Harpavat S, Wu S, Allred, EN, Berde C. Postoperative Pain and Other Sequelae of Dental Rehabilitations Performed on Children Under General Anesthesia. *Pediatric Dentistry* 2008 Mar-Apr; 30(2):111-21.
10. Primosch R, Nichols D, Courts D. Comparison of pre-operative ibuprofen, acetaminophen and placebo administration on parental report of postextraction pain in children. *Pediatr Dent* 1995; 17:187-191.

11. American Society of Anesthesiologists. Guidelines for Ambulatory Anesthesia and Surgery from American Society of Anesthesiologists. Available at: "<https://www.asahq.org/For-Members/Clinical-Information/~media/For%2520Members/documents/Standards%2520Guidelines%2520Smts/Ambulatory%2520Anesthesia%2520and%2520Surgery.ashx>" Accessed November 27, 2010.
12. ADA. Policy statement: The use of sedation and general anesthesia by dentists. Available at: "<http://www.ada.org/prof/resources/positions/statements/useof.aep>". Accessed November 27, 2010.
13. American Academy of Pediatric Dentistry. Guideline on use of local anesthesia for pediatric dental patients. Originating council. *Pediatr Dent* 2009; 31(6):141-7.
14. Gazal G, Bowman R, Worthington HV, Mackie IC. A double-blind randomized controlled trial investigating the effectiveness of topical bupivacaine in reducing distress in children following extractions under general anesthesia. *Int Journal of Paediatric Dentistry* 2004; 14:425-431.
15. McWilliams P, Rutherford J. Assessment of early postoperative pain or distress in children after exodontia under general anesthesia. *Evid Based Dent* 2007; 8(2):45-6.
16. Sammons HM, Unsworth V, Gray C, Choonara I, Cherrill J, Quirke W. Randomized controlled trial of the intraligamental use of a local anaesthetic (lignocaine 2%) versus controls in paediatric tooth extraction. *Int Journal of Paediatric Dentistry* 2007; 17(4):297-303.
17. Leong KJ, Roberts GJ, Ashley PF. Perioperative local anaesthetic in young paediatric patients undergoing extractions under outpatient 'short case' general anaesthesia. A double-blind randomized controlled trial. *British Dental Journal* 2007 Sep 22; 203(6):E11; discussion 334-5. *Epub* 2007 Aug 10.
18. Chi D, Kanellis M, Himadi E, Asselin M. Lip biting in a pediatric dental patient after dental local anesthesia: a case report. *Journal of Pediatric Nursing* 2008; 23(6):490-3.
19. College C, Feigal R, Wandera A, & Strange M. Bilateral versus unilateral mandibular block anesthesia in a pediatric population. *Pediatr Dent* 2000; 22:453-457.
20. Endo T, Gabka J, Taubenheim L. Intraligamentary anesthesia: benefits and limitations. *Quintessence Int* 2007; 37(88): 15-25.
21. Simard-Savoie S, Lemay H, Taleb L. The effect of epinephrine on pulpal microcirculation. *J Dent Res* 1979; 58(11): 2074-2079.
22. Caviedes-Bucheli J, Rojas P, Escalona M. The effect of different vasoconstrictors and

local anesthetic solutions on substance P expression in human dental pulp. *J Endod* 2009; 35:631-633.

23. Handler L, Albers D. The effects of the vasoconstrictor epinephrine on the duration of pulpal anesthesia using the intraligamentary injection. *J Am Dent Assoc* 1987; 114:807-809.
24. Edwards RW, Head TW. A clinical trial of intraligamentary anesthesia. *J Dent Res* 1989; 68(7): 1210-1214.
25. Jurgens S, Warwick R, Inglehearn P, Gooneratne D. Pain relief for paediatric dental chair anaesthesia: current practice in a community dental clinic. *Int J Paediatr Dent* 2003; 13:93-97.
26. Ashkenazi M, Blumer S, Eli I. Post-operative pain and use of analgesic agents in children following intrasulcular anaesthesia and various operative procedures. *British Dent J* 2007; 202: E13.
27. Wong DL, Baker CM. Pain in children: comparison of assessment scales. *Pediatr Nurs* 1988; 14:9-16.

## **Vita**

Dr. Belinda Lanore Campbell was born on February 9, 1979, in Wrangell, Alaska. She graduated from Stanwood High School, Stanwood, Washington in 1997. She received her Bachelor of Science in Psychology from the University of Washington, Seattle, Washington in 2000. She received her Doctorate of Dental Surgery from University of Southern California in 2005. During her time at USC she was awarded a National Health Service Corp scholarship and subsequently practiced dentistry for 5 years prior to returning to a residency in Pediatric Dentistry. She will graduate from Virginia Commonwealth University with a Master of Science in Dentistry and a Certificate in Pediatric Dentistry in June of 2012.