



VCU

Virginia Commonwealth University
VCU Scholars Compass

Theses and Dissertations

Graduate School

2012

Heart Failure Symptom Clusters and Functional Status

Janet Herr

Virginia Commonwealth University

Follow this and additional works at: <https://scholarscompass.vcu.edu/etd>

 Part of the [Nursing Commons](#)

© The Author

Downloaded from

<https://scholarscompass.vcu.edu/etd/2936>

This Dissertation is brought to you for free and open access by the Graduate School at VCU Scholars Compass. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Heart Failure Symptom Clusters and Functional Status

A dissertation submitted in partial fulfillment of the requirements for the degree of Doctor of Philosophy at Virginia Commonwealth University

by

Janet Kay Herr

Bachelor of Science in Nursing, University of Virginia, 1973
Master of Science, Virginia Commonwealth University, 1993

Director: Dr. Jeanne Salyer, Ph.D., R.N., F.N.A.P.
Associate Professor, Adult Health & Nursing Systems

Virginia Commonwealth University
Richmond, Virginia
November, 2012

Acknowledgment

The author wishes to thank several people. I would like to thank Dr. Jeanne Salyer Ph.D., RN, F.N.A.P, chairman of my dissertation committee, for her patience, support and guidance during my program of study at the School of Nursing, Virginia Commonwealth University. I would like to thank the members of my dissertation committee, Dr. Dolores Clement, Dr. P.H. FACHE, Professor, Dr. Lauren Goodloe, Ph.D., RN, Clinical Associate Professor, and Dr. Christine Schubert Kabban, Ph.D., MBA, Assistant Professor for their assistance and direction with the completion of this project. I would like to thank my family for their patience and support as I completed my program of study.

TABLE OF CONTENTS

List of Tables	v
List of Figures.....	vi
Heart Failure Symptom Clusters: A Systematic Review.....	
Abstract.....	2
Background.....	4
Methods.....	5
Data Synthesis.....	6
Relationship.....	7
Stability.....	11
Summary and Conclusions.....	13
References.....	16
Summary and Implications.....	24
Heart Failure Symptom Clusters and Functional Status	
Abstract.....	43
Background.....	45
Conceptual Framework.....	47
Methods.....	47
Results.....	52
Discussion	57

Limitations.....	60
Implications.....	61
Summary and Conclusions.....	62
References.....	63
What's New.....	71
Vita.....	79

LIST OF TABLES

Heart Failure Symptom Clusters: A Systematic Review

Table 1. Statistical Findings.....	25
------------------------------------	----

Heart Failure Symptom Clusters and Functional Status

Table 1. Sample Characteristics.....	72
--------------------------------------	----

Table 2. Symptom Prevalence.....	73
----------------------------------	----

Table 3. Symptom Factor Loadings.....	74
---------------------------------------	----

Table 4. Model Variables	75
--------------------------------	----

Table 5 Descriptive Statistics of Symptom Clusters.....	76
---	----

Table 6 Regression Analysis Functional Limitations and Mobility.....	77
--	----

LIST OF FIGURES

Heart Failure Symptom Clusters: A Systematic Review

Figure 1. Search and Screening Results..... 41

Heart Failure Symptom Clusters and Functional Status

Figure 1. Conceptual Framework..... 78

Abstract

HEART FAILURE SYMPTOM CLUSTERS AND FUNCTIONAL STATUS

Janet Kay Herr Ph.D

A dissertation submitted in partial fulfillment of the requirements for the degree of Doctor of Philosophy at Virginia Commonwealth University

Virginia Commonwealth University 2012

Major Director: Dr. Jeanne Salyer, Ph.D., R.N., F.N.A.P
Associate Professor, Adult Health & Nursing Systems

Clinical assessment of heart failure includes symptom identification and the evaluation of the relationship of symptoms to functional status. Symptom clusters are groups of at least 2 or 3 co-occurring symptoms that are related but are independent of other groups of symptoms.

The objectives of this study are to: (1) examine relationships among symptoms commonly experienced by individuals with heart failure, (2) identify symptoms that form clusters, and (3) evaluate the impact of heart failure symptom clusters on attributes of functional status: limitations and mobility

The Theory of Unpleasant Symptoms guided the conduct of this study. Heart failure symptoms and the outcome variables functional limitations and mobility were evaluated in a convenience sample of individuals (n = 117) with a confirmed diagnosis of heart failure recruited

from an academic medical center. Principle components analysis was used to extract symptom clusters and regression analysis was used to evaluate the relationship between the symptom clusters, their interaction terms, the demographic variables, age and co-morbidity, and functional status.

Three symptom clusters, sickness behavior, discomforts of illness, and GI distress were extracted. Predictors of functional limitations ($F = 35.96$, $p = 0.0005$, $R^2 = 0.578$) included sickness behavior ($\beta = -.681$, $p = 0.0005$), discomforts of illness ($\beta = -.765$, $p = 0.0005$) and the interaction term between these two symptom clusters ($\beta = .649$, $p = 0.014$). This model predicted 59% of the variance in functional limitations. Predictors of limited mobility ($F = 20.68$, $p = 0.0005$, $R^2 = 0.275$) included sickness behavior ($\beta = -0.441$, $p = 0.0005$) and co-morbidity ($\beta = -.200$, $p = 0.019$). This model predicted 28% of the variance in mobility.

Relationship between clusters of heart failure symptoms and functional limitations or mobility was observed. The interaction between discomforts of illness and sickness behaviors implies that not only do functional limitations increase as discomforts of illness increase, but increases at a faster rate when sickness behaviors are increased. Changes in sickness behaviors has the potential to improve mobility from being bed or chair bound everyday or most days to being bed or chair bound only some days.

Heart Failure Symptom Clusters: A Systematic Review

Janet K. Herr, MS, RN,

Jeanne Salyer, Ph.D., RN, Associate Professor

Debra E. Lyon, Ph.D., RN, FNP-BC, FNAP, FAAN, Professor

Lauren Goodloe, Ph.D., RN, Clinical Associate Professor

School of Nursing, Virginia Commonwealth University

Christine Schubert Kabban, Ph.D., MBA, Assistant Professor

Department of Mathematics and Statistics/ENC

Air Force Institute of Technology

Dolores G. Clement, Dr.P.H. FACHE, Professor

Department of Health Administration, School of Allied Health Professions

Virginia Commonwealth University, Richmond, Virginia

Corresponding author: Janet K. Herr, MS, RN

School of Nursing, Virginia Commonwealth University, Richmond, Virginia

1100 East Leigh Street

Richmond, Virginia 23298-0567

Phone: 804-323-8281 Fax: 1-866-891-1105 E-mail: jkherr@vcu.edu

Acknowledgements: Support for this study was provided by The Center of Excellence for Biobehavioral Approaches to Symptom Management (P30 NRO11403), Mary Jo Grap, Ph.D., RN, ACNP, FAAN, Principle Investigator, Department of Adult Health and Nursing Systems, School of Nursing, Virginia Commonwealth University

There are no conflicts of interest

Funding was not received for the conduct of this study.

1 Table

1 Figure

Abstract

Background: Heart failure is a prevalent chronic health condition in the United States.

Individuals who have heart failure experience as many as 2 to 9 symptoms. Symptom clusters are groups of 2 to 3 symptoms that co-occur in a relationship with each other.

Objective: The purpose of this systematic review is to examine which heart failure symptoms are related and to examine heart failure symptom relationships using different symptom measurement instruments.

Methods: The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed in the conduct of this systematic review. Pub Med, Psych Info, Cumulative Index of Nursing and Allied Health Literature (CINAHL) and the Cochrane Database were searched using the search terms heart failure and a pair of symptoms.

Results: Thirty-four studies out of a total of 1316 studies identified from database searches are included in this systematic review. A moderate level of correlation was found by more than one investigator between depression and fatigue, depression and anxiety, depression and sleep, depression and pain, anxiety and fatigue, and dyspnea and fatigue. Dyspnea was included in a symptom cluster identified by 6 studies. Three investigators identified a symptom cluster composed of psychosocial or emotional symptoms.

Conclusions: The findings of this systematic review provide support for the presence of heart failure symptom clusters. Some symptom relationships such as the relationships between fatigue and anxiety or sleep or pain were dependent on the symptom characteristics studied. Instrument selection affected the persistence of relationships among heart failure symptoms.

These findings suggest that studies defining the phenotype of individual heart failure symptoms may be a beneficial step in the study of heart failure symptom clusters.

Background

Heart failure is a clinical syndrome that affects 5.7 million adults age 20 and older. The incidence of heart failure is approximately 10 per 1000 population after 65 years of age.¹ Heart failure is the primary diagnosis for 41.6 % of visits to primary care offices and 21% of hospital emergency department visits.² Symptoms, subjective reports of abnormal physical, emotional, or cognitive conditions, trigger an individual to enter the healthcare system.³ Dyspnea and fatigue are frequently reported symptoms by individuals seeking emergency care for heart failure⁴ and in patients followed in cardiology clinics⁵ A growing public health concern, heart failure is a progressive and symptomatic illness with published reports suggesting that, on average, patients with heart failure experience anywhere from 2 to 9 symptoms.⁵⁻⁷

Symptom clusters are groups of symptoms that co-occur, are related, and yet, are independent of other groups of symptoms.⁸ Symptom clusters are composed of at least 2 or 3 symptoms.^{8,9} Although their recognition and use in the diagnosis and treatment of disease is not new,^{8,10} symptom cluster research is considered a new frontier in the study of symptoms.¹¹ Attributes of a symptom cluster include common etiology, concurrence, relationship, and stability.⁸ Heart failure, for example, provides a common etiology for symptoms such as dyspnea, fatigue, and edema which occur together in an individual who has heart failure. Relationship is an attribute that denotes the co-occurrence of symptoms, correlation between symptoms, and shared or overlapping variance. Stability is the reliability of the symptom cluster across subjects and over time.

The purpose of this systematic review is to summarize what is known about heart failure symptoms, a step in the identification of heart failure symptom clusters. Specific aims include examining which symptoms are related and examining heart failure symptom relationships using

different symptom measurement instruments. In the context of this review, a symptom cluster will be considered to be two or more symptoms occurring together and that are related to each other. A relationship among symptoms is indicated by a statistical association.

Methods

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines, which outline a methodology for performing systematic reviews in health care, were followed in the conduct of this systematic review.¹² Pub Med, Psych Info, Cumulative Index of Nursing and Allied Health Literature (CINAHL) and the Cochrane Database were searched for relevant articles. Search terms were heart failure in combination with a pair of symptoms. Specific symptoms used in the literature searches were dyspnea, fatigue, anxiety, depression, edema, pain, loss of appetite, sleep disturbance, and cognitive impairment. Dyspnea, fatigue, anxiety, depression, edema, pain, loss of appetite, sleep disturbance and cognitive impairment were selected as the focal point of this review because of their prevalence and distressing nature in heart failure patients.^{5,13,14} Reported to be associated with loss of gray matter volume assessed by MRI,¹⁵ Sloan and Pressler¹⁶ observed that individuals with heart failure relate awareness of heart failure-related cognitive impairments such as difficulty with memory and concentration. Studies were included in the review if they were published in the English language between 2000 and 2010 and reported a statistically significant relationship among 2 or more heart failure symptoms occurring together in subjects 18 years of age and older. A total of 1316 studies were identified from database and ancestry searches. Studies were excluded because they included conditions in addition to heart failure, evaluated the efficacy of an intervention for the treatment of heart failure or did not examine relationships among individual heart failure symptoms.

Seventy-nine studies were screened. Thirty-four studies meeting the inclusion criteria were included in this systematic review. Search and screening results are reported in Figure 1.

The methodological quality of the studies was evaluated using the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) criteria.¹⁷ The STROBE criteria consist of 22 items that should be included in descriptive observational studies and in studies investigating associations between exposures and health outcomes. The specific criteria refer to the title and abstract, introduction, methods, reporting of the results, and discussion sections of the report and to the reporting of study funding. A score of 1 was given for each of the criteria on the checklist that was present in the study report. Possible methodological quality scores range from 0 to 22. A score of 22 indicates a high level of clarity and reporting of the details about the conduct of the study. Studies included in this systematic review and the scientific quality score assigned using the STROBE criteria may be found in Table 1. Scientific quality scores for the studies included in this systematic review ranged from a low score of 14 to a high score of 22. The median quality score was 18.

Data Synthesis

A total of 34 studies were included in the systematic review. Relationship and stability are symptom cluster attributes considered in this analysis of heart failure symptom associations. Miaskowski, Aouizerat, Dodd, and Cooper¹⁸ advise that before a group of symptoms can be classified as a symptom cluster, illustration of a relationship among the symptoms is required. Kim et al.⁸ characterizes the nature of symptom relationships as an associative rather than a causal connection.

Stability refers to the replicability of a symptom cluster at different times and in different samples.⁸ Stability may be assumed in symptom clusters that are established for diagnostic

purposes.⁸ In identifying stable symptom clusters Miaskowski, Dodd, and Lee¹¹ suggest evaluations of symptom clusters should focus on heterogeneous patient populations to permit comparisons across patients with respect to prevalence, severity, and the effect of the symptom cluster on patient outcomes. If a symptom cluster does not occur across different patient groups the data may reflect a different underlying mechanism.

Relationship

Clinically, symptom clusters may be identified using observed relationships and concurrence. On the other hand symptom clusters may be identified using mathematical procedures. Barsevick¹⁹ suggests mathematical procedures such as bivariate correlation, factor analysis, or cluster analysis may be useful in examining symptom relationships. Investigators also used regression analysis to uncover symptom relationships in the studies included in this systematic review.

The relationship between depression and fatigue was frequently studied by investigators. Chen et al.,²⁰ Evangelista, et al.,²¹ Fink, et al.,²² Redeker,²³ Sullivan, et al.,²⁴ and Tang, et al.²⁵ reported relationships between depression and general measures of fatigue such as the Multidimensional Assessment of Fatigue Scale, the Fatigue Symptom Inventory, or the Profile of Mood States Fatigue Scale ($r = 0.34 - 0.77$, $p < 0.001 - 0.01$). Smith et al.²⁶ observed relationships between depression and general fatigue ($r = 0.42$, $p < 0.001$) and depression and exertion fatigue ($r = 0.33$, $p < 0.001$). Falk, et al.²⁷ reported relationships between depression and general fatigue ($R^2 = .145$; $F = 21.2$; $\beta = .30$, $p < .01$), depression and reduced activity ($R^2 = .178$; $F = 23.0$; $\beta = .37$; $p < .001$), and depression with reduced motivation ($R^2 = .110$; $F = 13.7$; $\beta = .40$; $p < .001$). However Falk, et al.²⁷ did not observe a statistically significant relationship between depression and physical fatigue or between depression and mental fatigue.

The relationship between depression and anxiety ($r = 0.52 - 0.74$, $p = 0.01 - 0.001$) persists across studies and irrespective of the instrument used to measure these symptoms.²⁸⁻³² Goebel, et al.⁶ observed a relationship between depression and pain ($r = 0.32$, $p = 0.002$). Sullivan et al.²⁴ found depression was related to chest pain ($r = 0.38$ to 0.43 , $p < .001$) and inversely related to bodily pain ($r = -0.37$ to -0.47 , $p < .001$). Depression was related to the sleep characteristics, difficulty initiating and maintaining sleep (OR = 5.09 95% CI 2.41 - 10.75),³³ sleep quality ($r = 0.40 - 0.71$, $p = 0.001$)³⁴⁻³⁵ and sleep disordered breathing ($r = 0.88$, $p = 0.001$)³⁴ but not to severity of sleep disordered breathing.³⁶ Depression was also reported to be related to appetite loss (OR 2.28, $p = 0.001$)³⁷ and to vitality ($r = -0.40$ to -0.44 , $p < 0.001$).²⁴

Relationships between fatigue and anxiety, sleep, or pain were contingent on particular characteristics of fatigue. Anxiety was related to general measures of fatigue ($r = 0.42 - 0.49$, $p < 0.001$)^{20,23} and to mental fatigue ($R^2 = 0.254$; $F = 36.2$; $\beta = .39$; $p < .001$).²⁷ Yu et al.³⁸ detected a relationship between fatigue and the construct of anxiety/depression assessed using the Hospital Anxiety and Depression Scale (HADS) ($r = -0.50$, $p < 0.001$). Anxiety was not related to general fatigue, physical fatigue, reduced activity or reduced motivation.²⁷ Falk, Patel, et al.²⁷ observed that general fatigue ($\beta = 0.22$; $p < 0.05$) and physical fatigue ($\beta = 0.22$, $p < 0.05$) were related to pain distress. Smith et al.²⁶ found a relationship between cardiac pain and general fatigue ($r = 0.44$, $p < 0.001$) as well as cardiac pain and exertion fatigue ($r = 0.42$, $p < 0.001$). Difficulty with sleep was not related to fatigue intensity ($r = 0.20$, $p > .05$),³⁹ but was related to general fatigue ($r = 0.42$, $p < .001$) and to exertion fatigue ($r = 0.29$, $p = .001$).²⁶ Redeker et al.³⁶ noted a non-significant relationship between global fatigue and severity of sleep disordered breathing. Stephen³⁹ observed a relationship between fatigue intensity and nocturia ($r = 0.36$, $p < 0.01$). Falk et al.²⁷ reported that physical fatigue was related to appetite ($\beta = 0.26$; $p < 0.01$).

while nausea was related to general fatigue ($\beta= 0.18$; $p<0.05$) and reduced activity ($\beta=0.23$; $p < 0.05$).

Dyspnea, a cardinal heart failure symptom, was related to several other heart failure symptoms. Dyspnea was observed by Ramasamy et al.⁴⁰ ($r = 0.40$ df 46 $p<0.01$) and Sullivan et al.²⁴ ($r = 0.43 - 0.56$, $p < 0.001$) to be related to depression. Bekelman et al.⁵ found 5:1 odds that an individual with probable depression would be quite a bit or very much distressed by shortness of breath compared to study participants who did not have depression (OR 5.28, $p = .05$). Redeker²³ found a relationship between dyspnea and anxiety ($r = 0.22 - 0.35$, $p = 0.001 - 0.02$), however, Ramasamy et al.⁴⁰ did not find a relationship between dyspnea and anxiety. Yu et al.³⁸ observed an inverse correlation between dyspnea and the construct of anxiety/depression ($r = -0.17$, $p < 0.05$). Weibel et al.⁴¹ observed moderate daily correlations between dyspnea and edema ($r = 0.45 - 0.63$, $p \leq 0.001$). Dyspnea was related to fatigue including the characteristics general fatigue,²⁶⁻²⁷ exertion fatigue²⁶ and fatigue intensity.³⁹ Paroxysmal nocturnal dyspnea was related to the sleep characteristics sleepiness ($r = 0.161$, $p 0.02$) and insomnia ($r = 0.279$, $p = 0.004$).⁴² Dyspnea had a non-significant relationship with sleep disturbance.⁴⁰ The relationship between sleep disordered breathing and sleep quality or excessive daytime sleepiness was also non-significant.³⁶

Akomolafe, et al.⁴³ and Sauve et al.⁴⁴ did not detect a relationship between cognitive function and depression. Sauve et al.⁴⁴ did not find that cognitive function was related to anxiety or pain. Some investigators document that individuals with heart failure have difficulty detecting symptoms of heart failure.⁴⁵⁻⁴⁷ Older individuals may require assistance with the detection and interpretation of their heart failure symptoms.⁴⁸ Cameron, Worrall-Carter, Page, Riegel, Lo, and

Stewart⁴⁹ observed that mild cognitive dysfunction is a predictor of the ability to independently recognize symptoms changes.

Using statistical procedures such as principal components analysis,⁵⁰⁻⁵² confirmatory factor analysis,³³ hierarchical cluster analysis,^{7,53,54} and structural equation modeling⁵⁵ investigators developed the following relationships among 3 or more symptoms. Dyspnea was included in a cluster of symptoms in 6 studies.^{7,50,51,53-55} The characteristics of dyspnea that were observed in a cluster included orthopnea, paroxysmal nocturnal dyspnea, dyspnea on exertion, shortness of breath at rest, and shortness of breath at night. Two studies reported a symptom cluster composed entirely of characteristics of dyspnea.^{7,50} Five of the symptom clusters that included dyspnea also included fatigue^{50,51,53} or the dimensions of fatigue, increased need to rest⁵⁴ and vital exhaustion.⁵² Sleep disturbance, sleep difficulty or sleep problems were included with dyspnea in 3 symptom clusters.^{51,53,54} Edema described as weight gain,⁵⁰ leg edema,⁵⁵ or bloating and swelling⁵³ was also included in a symptom cluster. However, three investigators reported that edema, formed a cluster that was separate from other heart failure symptoms.^{7,50,54} Hertzog, et al.⁵³ found that forgetfulness was included in a cluster with difficulty sleeping, fatigue, and four dimensions of shortness of breath. Hertzog, et al.⁵³ also reported that dizziness was associated with shortness of breath with activity, shortness of breath at rest, bloating and swelling, fatigue, and depression. Jurgens et al⁵⁰ identified a group of symptoms composed of feeling one's heart beat become faster, feeling chest pain, having an upset stomach, and having a cough.

Three investigators identified heart failure symptom clusters primarily consisting of psychosocial or emotional symptoms.^{51, 54, 55} Depression was a component of all 3 of the heart failure symptom clusters. Worry was included in 2 of these symptom clusters.^{51, 54} Fatigue and

vital exhaustion were each included in a symptom cluster that included sleep disturbance and depression.^{52,55} Jurgens et al.⁵¹ and Smith et al.⁵² observed that difficulty concentrating, was included in a heart failure symptom cluster with psychosocial or emotional symptoms.

In summary this systematic review points out relationships between symptoms experienced by individuals with heart failure that include depression, fatigue and dyspnea. Additionally, relationships between symptom characteristics were also observed. Cognitive impairment does not appear to have a relationship with the heart failure symptoms of depression, anxiety or pain. However, forgetfulness and difficulty concentrating appear in symptom clusters along with shortness of breath, fatigue, difficulty sleeping, worry and feeling depressed.

Stability

Stability may be indicated by the reproducibility of a symptom cluster across multiple patient samples.¹⁸ At this point in the study of heart failure symptom clusters, it is early to comment on the stability of a heart failure symptom cluster. However, there are several observations regarding the persistence of co-occurring heart failure symptom relationships. It is notable that there is similarity in the composition of specific heart failure symptoms in pairs and heart failure symptom clusters reported by several investigators. For example, depression and fatigue,²⁰⁻²⁵ depression and anxiety,²⁸⁻³² and dyspnea and fatigue along with sleep disturbance^{51,53,54} were validated by several investigators. Observation of a symptom cluster in different settings is indicative of stability or the replicability of a symptom relationship. Although the studies included in this systematic review were conducted primarily in the outpatient setting*, some of the studies were conducted in the inpatient setting†. Time, which refers to the duration of association among the symptoms in a cluster,⁸ is a component of the stability of a symptom

*5,6,21 – 24, 26, 29, 34, 35, 41, 39, 43, 44, 50, 52, 53, 55.

†7,25, 27, 30, 37, 38, 51.

cluster. The symptom pairs, pain and depression, anxiety and depression, and depression and fatigue were validated by studies conducted over the time period 2004 through 2010.

Instrument selection also appears to have an influence on the stability of relationships among heart failure symptoms and the persistence of the heart failure symptom clusters. For example, depression evaluated using the Short Form Geriatric Depression Scale, Beck Depression Inventory, the Center for Epidemiologic Studies Depression Scale (CESD) or the Profile of Mood States Depression/Dejection scale correlated with sleep ($r = 0.402, p < 0.001$),³⁵ sleep disturbance ($r = 0.39 - 0.53, p < 0.001$),²³ or fatigue ($r = 0.34 - 0.64, p 0.001 - 0.01$).^{22,23} However, investigators reported that depression assessed using the Hospital Anxiety and Depression Scale (HADS) depression subscale was not related to sleep disturbance ($r = 0.19 - 0.26, p 0.06 - < 0.11$)^{23,40} or excessive daytime sleepiness ($r = -0.02, p 0.855$).²³ Chen, et al.²⁰ observed a relationship between depression assessed using the HADS depression scale and fatigue assessed using the Piper Fatigue Scale ($r = 0.5, p < 0.001$). Ramasamy et al.⁴⁰ and Redeker²³ did not observe a relationship between depression, assessed using the HADS depression scale, and fatigue.

Seven of the 34 studies reported non-significant relationships between some or all of the symptoms studied. Four of these studies had sample sizes between 50 and 70 subjects and used Pearson's correlation to detect symptom relationships. Two studies with sample sizes of 100 or more observed a low proportion of the symptoms of interest. For example, Akomolafe et al.⁴³ who studied depression and cognitive impairment in a sample of 100 heart failure patients, reported that 23% of the sample had depression and 10% of the sample had cognitive impairment. This investigator did not detect a relationship between depression and cognitive impairment. Falk et al.²⁷ who studied relationships between fatigue, anxiety and depression,

reported the prevalence of depression in the sample of 112 individuals with heart failure was 18% and the prevalence of anxiety was 10%. Nonsignificant relationships were observed between anxiety and the fatigue characteristics, general fatigue, physical fatigue, reduced activity and reduced motivation. A statistically significant relationship was detected between anxiety and mental fatigue. Falk et al.²⁷ also detected non-significant relationships between depression and the fatigue characteristics, physical fatigue and mental fatigue. However, relationships were detected between depression and the fatigue characteristics, general fatigue, reduced activity, and reduced motivation. Redeker et al.³⁶ did not detect relationships among symptoms studied in a sample of 170 individuals with heart failure in which the prevalence of the symptoms of interest was between 46% and 85%. The 3 studies with samples of 100 or more study participants employed regression analysis to examine symptom relationships.^{27, 36, 43} While there were statistically significant findings in these studies, a greater prevalence of a symptom in the sample would add credibility to the conclusions reached by the investigators.

As work on identification of heart failure symptom clusters progresses, it will be important to be aware of the symptom characteristics measured by the assessment tool selected to measure the symptoms of interest as well as the prevalence of the symptom in the sample. Perhaps findings of variation in symptom relationships realized by the use of different instruments to evaluate a symptom is leading symptom assessment for heart failure symptom cluster development toward an emphasis on symptom characteristics rather than overall symptom construct.

Summary and Conclusions

This systematic review of studies reporting about heart failure symptom relationships provides evidence of the existence of heart failure symptom clusters. Depression and anxiety,

depression and fatigue, depression and pain, fatigue and pain, dyspnea and depression, and dyspnea and sleep are some symptoms for which relationships were established by investigators. A notable finding of this systematic review is differences in relationships when symptom characteristics are considered. For example, anxiety was related to mental fatigue but not to general fatigue, physical fatigue, reduced activity or reduced motivation. There was an association between depression and difficulty initiating and maintaining sleep, sleep quality and sleep disordered breathing, however, depression was not related to severity of sleep disordered breathing. Relationships between depression and sleep or fatigue did not persist when the HADS depression scale was used to assess depression as opposed to when the Beck Depression Inventory, CESD, or POMS depression/dejection scale were used to evaluate depression. These findings suggests that as work on heart failure symptom cluster identification continues, consideration should be given to examining relationships among the various symptom characteristics in addition to relationships among general measures of the symptom. Studies that define the phenotype of individual heart failure symptoms may be needed prior to the identification of heart failure symptom clusters.

Symptom relationships between cognitive function and depression, anxiety or pain were not detected by the investigators included in this systematic review. As work on isolation of heart failure symptom clusters progresses investigators would be well advised to remember that symptom detection may be difficult for individuals with heart failure and to consider the use of instruments for symptom assessment in impaired individuals such as those available for pain assessment in non-verbal individuals.

This systematic review adds to the body of work on the identification of heart failure symptom clusters by providing some insight into the composition of heart failure symptom

clusters. Additional study is needed before the composition of heart failure symptom clusters can be defined. This study, however, provides some initial findings about heart failure symptom assessment for use in identification of heart failure symptom clusters. Advancing the science of heart failure symptom phenotyping and heart failure symptom cluster identification has the potential to lead to more effective interventions that target a group of symptoms rather than a single symptom. Decreasing the number of interventions a patient must employ to manage heart failure symptoms is one way to facilitate heart failure symptom self-management.

References

1. Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics – 2011 update: A report from the American Heart Association. *Circulation*. 2011;123(4): e18 -e209.
doi:10.1161/CIR.0b013e3182009701
2. Schappert SM, Rechtsteiner EA. Ambulatory medical care utilization estimates for 2007 National Center for Health Statistics. *Vital Health Stat* 13. 2011;13(169):23.
http://www.cdc.gov/nchs/data/series/sr_13/sr13_169.pdf. Accessed August 8, 2011.
3. Wilson, IB, Cleary, PD. Linking clinical variables with health-related quality of life, a conceptual model of patient outcomes. *JAMA*. 1995;273(1):59 – 65.
4. Patel H, Shafazand M, Schaufelberger M, Ekman, I. Reasons for seeking acute care in chronic heart failure. *Eur J Heart Fail*. 2007;9(6-7):702-708.
doi:10.1016/j.ejheart.2006.11.002
5. Bekelman DB, Havranek EP, Becker DM, et al. Symptoms, depression, and quality of life in patients with heart failure. *J Card Fail*. 2007;13(8): 643-648.
doi:10.1016/j.cardfail.2007.05.005
6. Goebel JR, Doering LV, Shugarman LR, et al. Heart failure: The hidden problem of pain. *J Pain Symptom Manage*. 2009;38(5): 698-707. doi: 10.1016/j.jpainsymman.2009.04.022.
7. Song EK, Moser DK, Rayens MK, Lennie TA. Symptom clusters predict event-free survival in patients with heart failure. *J Cardiovasc Nurs*. 2010; 25(4): 284-291.
doi:10.1097/JCN.0b013e318cfbcbb.
8. Kim HJ, McGuire DB, Tulman L, Barsevick AM. Symptom clusters, concept analysis and clinical implications for cancer nursing. *Cancer Nurs*. 2005; 28(4):270-282.
<http://ovidsp.tx.ovid.com.proxy.library.vcu.edu/sp-3.5.1a/ovidweb.cgi?QS2>. Accessed April 21, 2011.

9. Dodd, M.J., Miaskowski, C., & Paul, S.M. Symptom clusters and their effect on the functional status of patients with cancer. *Oncol Nurs Forum*. 2001;28(3): 465 – 470.
<http://web.ebscohost.com.proxy.library.vcu.edu/ehost/pdfviewer/pdfviewer?sid=b271ab0b-faae-430a-9568-99976fc9ab95%40sessionmgr12&vid=2&hid=25>. Published July 13, 2001.
Updated December 12, 2009. Accessed April 21, 2011.
10. Barsevick A. The concept of symptom cluster. *Semin Oncol Nurs*. 2007; 23(2):89 – 98.
doi:10.1016/j.soncn.2007.01.009
11. Miaskowski C, Dodd M, Lee K. Symptom clusters: The new frontier in symptom management research. *J Natl Cancer Inst Monogr*. 2004;2004(32):17-21.
doi:10.1093/jncimonographs/lgm003
12. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS medicine*.2009; 6(7): 1 – 26. doi:10.1371/journal.pmed.1000100
13. Blinderman CD, Homel P, Billings JA, Portenoy RK, Tennstedt SL (2008). Symptom distress and quality of life in patients with advanced congestive heart failure. *J Pain Symptom Manage*.2008;35(6): 594 – 603. doi:10.1016/j.jpainsymman.2007.06.007
14. Zambroski CH, Moser DK, Bhat G, Ziegler C. Impact of symptom prevalence and symptom burden on quality of life in patients with heart failure. *Eur J Cardiovasc Nurs*, 2005; 4(3):198-206. doi:10.1016/j.ejcnurse.2005.03.010
15. Almeida OP, Garrido GJ, Beer, C, Lautenschlager NT, Arnolda L, Flicker L. Cognitive and brain changes associated with ischaemic heart disease and heart failure. *Eur Heart J*. 2012; 33(14): 1769-1776. doi:10.1093/eurheartj/ehr467

16. Sloan RS, Pressler SJ. Cognitive deficits in heart failure, re-cognition of vulnerability as a strange new world. *J Cardiovasc Nurs.* 2009;24(3):241-248.
doi:10.1097/JCN.0b013e3181a00284
17. Vanderbroucke, J.P., von Elm, E., Altman, D.G., et al. Strengthening the reporting of observational studies in epidemiology (STROBE): explanation and elaboration. *Ann Intern Med.* 2007;147(8):W-163-W-194. <http://annals.org/content/147/8/W-163-full.pdf+html>.
Accessed November 20, 2011
18. Miaskowski C, Aouizerat BE, Dodd M, Cooper B. Conceptual issues in symptom clusters research and their implications for quality-of-life assessment in patients with cancer. *J Natl Cancer Inst Monogr.* 2007;2007(37):39-46. doi: 10.1093/jncimonographs/lgm003
19. Barsevick AM. The elusive concept of the symptom cluster. *Oncol Nurs Forum.* 2007;34(5):971 – 980. doi:10.1188/07.ONF.971-980
20. Chen LH, Li C Y, Shieh SM, Yin, WH, Chiou AF Predictors of fatigue in patients with heart failure. *J Clin Nurs.*2010;19(11-12):1588-1596. doi:10.1111/j.1365-2702.2010.03218.x
21. Evangelista LS, Moser DK, Westlake C, Pike N, Ter-Galstanyan A, Dracup K. Correlates of fatigue in patients with heart failure. *Prog Cardiovasc Nurs.* 2008; 23(1): 12-17.
<http://web.ebscohost.com.proxy.library.vcu.edu/ehost/pdfviewer/pdfviewer?sid=45a802e2-e9c3-4dae-85cd-0be4389c6159%40sessionmgr4&vid=2&hid=25>. Published May 2, 2008.
Updated December 31, 2010. Accessed 8/14/2011.
22. Fink AM, Sullivan SL, Zerwic JJ, Piano MR. Fatigue with systolic heart failure. *J Cardiovasc Nurs.* 2009; 24(5): 410 – 417. doi:10.1097/JCN.0b013e3181ae1e84.
23. Redeker NS. Somatic symptoms explain differences in psychological distress in heart failure patients vs. a comparison group. *Prog Cardiovasc Nurs.* 2006; 21(4): 182-189.

<http://web.ebscohost.com.proxy.library.vcu.edu/ehost/pdfviewer/pdfviewer?sid=dea525e7-5db1-4ef3-9812-3da3034e981e%40sessionmgr11&vid=4&hid=8>. Accessed April 19, 2011.

24. Sullivan M, Levy WC, Russo JE, Spertus JA. Depression and health status in patients with advanced heart failure: A prospective study in tertiary care. *J Card Fail.* 2004; 10(5):390-396. doi: 10.1016/j.cardfail.2004.01.011
25. Tang WR, Yu CY, Yeh SJ. Fatigue and its related factors in patients with chronic heart failure. *J Clin Nurs.* 2010; 19(1-2): 69-78. doi:10.1111/j.1365-2702.2009.02959.x
26. Smith ORF, Michielsen HJ, Pelle AJ, Schiffer AA, Winter JB, Denollet J. Symptoms of fatigue in chronic heart failure patients: clinical and psychological predictors. *Eur J Heart Fail.* 2007; 9(9): 922-927. doi:10.1016/j.ejheart.2007.05.016
27. Falk K, Patel H, Swedberg K, Ekman I. Fatigue in patients with chronic heart failure – a burden associated with emotional and symptom distress. *Euro J Cardiovasc Nurs.* 2009; 8(2): 91-96. doi: 10.1016/j.ejcnurse.2008.07.002.
28. Bean MK, Gibson D, Flattery M, Duncan A, Hess, M. Psychosocial factors, quality of life, and psychological distress: ethnic differences in patients with heart failure. *Prog Cardiovasc Nurs.* 2009;24(4): 131-140. doi:10.1111/j.1751-7117.2009.00051x
29. Chung ML, Moser DK, Lennie TA, Rayens MK. The effects of depressive symptoms and anxiety on quality of life in patients with heart failure and their spouses: Testing dyadic dynamics using Actor-Partner Interdependence Model. *J Psychosom Res.* 2009; 67(1): 29-35. doi:10.1016/j.jpsychores.2009.01.009
30. Jiang W, Kuchibhatala M, Cuffe MS, et al. Prognostic value of anxiety and depression in patients with chronic heart failure. *Circulation.* 2004; 110(22): 3452 – 3456. doi: 101161/01.CIR.0000148138.25157.F9.

31. Luyster FS, Hughes JW, Gunstad J. Depression and anxiety symptoms are associated with reduced dietary adherence in heart failure patients treated with an implantable cardioverter defibrillator. *J Cardiovasc Nurs*. 2009; 24(1): 10-17. doi 10.1097/01.JCN.0000317469.63886.24
32. Paukert AL, LeMaire A, Cully JA. Predictors of depressive symptoms in older veterans with heart failure. *Aging Ment Health*. 2009; 13(4): 601-610. doi: 10.1080/13607860802459823.
33. Redeker NS, Jeon S, Muench U, Campbell D, Walsleben J, Rapoport DM. Insomnia symptoms and daytime function in stable heart failure. *Sleep*. 2010; 33(9): 1210-1216. <http://www.ncbi.nlm.nih.gov.proxy.library.vcu.edu/pmc/articles/PMC2938862/pdf/aasm.33.9.1210.pdf>. Accessed April 19, 2011.
34. Skobel E, Norra C, Sinha A, Breuer C, Hanrath P, Stellbrink C. Impact of sleep-related breathing disorders on health-related quality of life in patient with chronic heart failure. *Eur J Heart Fail*. 2005; 7(4): 505-511. doi:10.1016/j.eheart.2004.07.020.
35. Wang TJ, Lee SC, Tsay SL, Tung HH. Factors influencing heart failure patients' sleep quality. *J Adv Nurs*. 2010; 66(8):1730-1740. doi: 10.1111/j.1365-2648.2010.05342.x.
36. Redeker NS, Muench U, Zucker MJ, et al. Sleep disordered breathing, daytime symptoms, and functional performance in stable heart failure. *Sleep*. 2010; 33(4): 551-560. <http://www.ncbi.nlm.nih.gov.proxy.library.vcu.edu/pmc/articles/PMC2849795/pdf/aasm.33.4.551.pdf>. Accessed April 13, 2011.
37. Lesman-Leegte I, Jaarsma T, Sanderman R, Linssen G, van Veldhuisen DJ. Depressive symptoms are prominent among elderly hospitalized heart failure patients. *Eur J Heart Fail*. 2006; 8(6): 634-640. doi:10.1016/j.ejheart.2005.11.010

38. Yu DSF, Lee DTF, Woo J, Thompson DR. Correlates of psychological distress in elderly patients with congestive heart failure. *J Psychosom Res.* 2004; 57(6): 573-581.
doi:10.1016/j.jpsychores.2004.04.368
39. Stephen SA. Fatigue in older adults with stable heart failure. *Heart Lung.* 2008; 37(2): 122 - 131. doi:10.1016/j.hrtlng.2007.03.006.
40. Ramasamy R, Hildebrandt T, O’Hea E, et al. Psychological and social factors that correlate with dyspnea in heart failure. *Psychosomatic.*2006; 47(5): 430-434. doi:
10.1176/appl.psy.47.5.430
41. Webel AR, Frazier SK, Moser DK, Lennie TA. Daily variability in dyspnea, edema and body weight in heart failure patients. *Eur J Cardiovasc Nur.* 2007; 6(1):60-65.
doi:10.1016/j.ejcnurse.2006.04.003
42. Principe-Rodriguez K, Strohl KP, Hadziefendic S, Piña IL. Sleep symptoms and clinical markers of illness in patients with heart failure. *Sleep Breath.* 2005; 7(3): 127-133. doi:
10.1007/s11325-005-0023-0.
43. Akomolafe A, Quarshie A, Jackson P, et al. The prevalence of cognitive impairment among African-American patients with congestive heart failure. *J Natl Med Assoc.* 2005; 97(5):689-694.
44. Sauvé MJ, Lewis WR, Blankenbiller M, Rickabaugh B, Pressler SJ. Cognitive impairments in chronic heart failure: a case controlled study. *J Card Fail.* 2009; 15(1): 1-10.
doi:10.1016/j.cardfail.2008.08.007.
45. Friedman MM, Quinn JR. Heart failure patients’ time, symptoms, and actions before a hospital admission. *J Cardiovasc Nurs.* 2008;23(6):506 – 512.
doi:10.1097/01.JCN.0000338928.51093.40

46. Horowitz CR, Rein SB, Leventhal H. A story of maladies, misconceptions and mishaps: effective management of heart failure. *Soc Sci Med.* 2004;58(3):631-643.
doi:10.1016/S0277-9536(03)00232-6
47. Jurgens CY. Somatic awareness, uncertainty, and delay in care-seeking in acute heart failure. *Res Nurs Health.* 2006;29(2):74-86. doi:10.1002/nur.20118
48. Riegel B, Dickson VV, Cameron J., et al. Symptom recognition in elders with heart failure. *J Nurs Scholarsh.* 2010;42(1):92-100. doi: 10.1111/j.1547-5069.2010.01333.x
49. Cameron J, Worrall-Carter L, Page K, Riegel B, Lo SK, Stewart S. Does cognitive impairment predict poor self-care in patients with heart failure? *Eur J Heart Fail.* 2010;12(5):508-515. doi:10.1093/eurjhf/hfq042
50. Jurgens, C.Y., Fain, J.A., & Riegel, B. Psychometric testing of the heart failure somatic awareness scale. *J Cardiovasc Nurs.* 2006;21(2): 95-102.
<http://ovidsp.tx.ovid.com.proxy.library.vcu.edu/sp-3.5.1a/ovidweb.cgi?QS2>. Accessed August 8, 2011.
51. Jurgens CY, Moser DK, Armola R, et al. Symptom clusters of heart failure. *Res Nurs Health.* 2009; 32(5), 551 – 560. doi:10.1002/nur.20343
52. Smith ORF, Gidron Y, Kupper N, Winter JB, Denollet J. Vital exhaustion in chronic heart failure: symptom profiles and clinical outcome. *J Psychosom Res.* 2009; 66(3):195-201.
doi:10.1016/j.jpsychores.2008.10.021
53. Hertzog, MA, Pozehl, B, Duncan K. Cluster analysis of symptom occurrence to identify subgroups of heart failure patients, a pilot study. *J Cardiovasc Nurs.* 2010; 25(4): 273-283.
doi:10.1097/JCN.0b013e318cfbb6c.

54. Lee KS, Song EK, Lennie TA, et al. Symptom clusters in men and women with heart failure and their impact on cardiac event-free survival. *J Cardiovasc Nurs.* 2010; 25(4): 263-272.
doi: 10.1097/JCN.0b013e3181cfbb88.
55. Johansson P, Alehagen U, Svensson E, Svanborg E, Dahlström U, Broström A. Determinants of global perceived health in Community-dwelling elderly screened for heart failure and sleep-disordered breathing. *J Cardiovasc Nurs.* 2010; 25(5), E16 – E26.
doi:10.1097/JCN.0b013e3181d6de6f.

Summary and Implications

- This systematic review observed relationships among several heart failure symptom pairs, for example, depression and anxiety, depression and fatigue, depression and pain, fatigue and pain, dyspnea and depression, and dyspnea and sleep.
- This analysis reveals relationships among heart failure symptom characteristics, such as mental, physical, or exertion fatigue, suggesting that identification of the phenotype of individual heart failure symptoms may be informative.
- Symptom assessment in cognitively impaired individuals with heart failure may present a challenge to investigators' attempts to identify heart failure symptom clusters.
- Instrument selection may influence the persistence of relationships among heart failure symptoms.

Table 1

Statistical Findings

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
Akomolafe et al ⁴³ (2005)	Cross-sectional design 100 heart failure patients who were consecutively recruited from an inner-city physician practice, rural physician practice and an urban physician practice for a single-blinded, randomized, controlled trial to evaluate the effectiveness of a telemonitoring intervention. Sample age <64 years - 39% 65-74 years -39% > 75 years - 22% Gender - 64% female	multiple logistic regression	Mini-Mental Status Examination (not reported) Geriatric Depression Scale (not reported)	Depression and cognitive impairment OR 0.78, 95%, p=0.8.0	18
Bean et al ²⁸ (2009)	Cross-sectional, correlational design 97 heart failure patients who were inpatients on the cardiology unit or an outpatient in the heart failure clinic of	Pearson's correlation	Hospital Anxiety and Depression Scale Anxiety scale $\alpha = .85$ Depression scale $\alpha = .83$	Anxiety and depression $r = 0.74$ $p < 0.01$	19

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	an urban health center Mean age 53 years 33% female				
Bekelman et al. ⁵ (2007)	Cross-sectional design 60 participants recruited from outpatient cardiology clinics at an academic – affiliated community hospital and a tertiary care academic referral hospital. Median age 75 years 37% female	Logistic regression model	Memorial Symptom Assessment Scale – Short Form (not reported) Geriatric Depression Scale – Short Form (not reported)	Depression and shortness of breath – patients with probable depression were more likely to be quite a bit or very much distressed by shortness of breath Adjusted odds ratio 5.28 p = .05	17
Chen et al. ²⁰ (2010)	Cross-sectional correlational design Convenience sample of 105 patients with diagnoses of heart failure recruited from outpatient clinics or cardiovascular units of 3 hospitals Mean age 65.2 years Gender – 35.2% female	Pearson product moment correlation coefficients	Piper fatigue scale ($\alpha = 0.85$) Hospital anxiety and depression scale ($\alpha 0.78$ for the global scale, $\alpha 0.73$ for the anxiety scale, $\alpha 0.95$ for the depression scale)	Depression and fatigue – $r = 0.5$ p< 0.001 Anxiety and fatigue – $r = 0.49$ p< 0.001	21
Chung et al. ²⁹ (2009)	Cross-sectional descriptive design 58 patient-spouse dyads recruited from cardiac	Pearson Product Moment Correlation	Brief Symptom Inventory - depressive symptom subscale (for patients $\alpha = .92$, for	Depression and anxiety $r = .72$ p<.001	18

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	outpatient clinics at an academic center Mean age 61.7 years Gender – 26% women		spouses $\alpha = .83$,) Anxiety subscale (for patients $\alpha = .86$, for spouses $\alpha = .87$)		
Evangelista et al ²¹ (2008)	Cross-sectional correlational design 150 patients with HF recruited from a single outpatient heart failure clinic within a tertiary, university affiliated medical center Mean age 55 years Gender – 27.3% women	Correlation	Profile of Mood States – Fatigue subscale ($\alpha = 0.88$) Beck Depression Inventory (not reported)	Depression and fatigue $r = .576$ $p < .001$	15
Falk et al ²⁷ (2009)	(no study design reported) [cross-sectional] 112 patients hospitalized in a university hospital with CHF 112 hospitalized patients with CHF Mean age 77 years Gender – 40% women	Stepwise forward multiple regression	Multidimensional Fatigue Inventory – Swedish version (α ranged from 0.60 – 0.78) Hospital Anxiety and Depression Scale Anxiety $\alpha = 0.77$ Depression $\alpha = 0.78$ Symptom Distress Scale $\alpha = 0.79$	General Fatigue Anxiety - $R^2 .049$ F 6.1; $\beta -.03$ (ns) Depression $R^2 .145$ F 21.2; $\beta .30$ ($p < .01$) Physical Fatigue Anxiety – $R^2 .016$ F 1.8; $\beta -.07$ (ns) Depression – $R^2 .056$ F 6.8; $\beta .14$ (ns) Reduced Activity Anxiety $R^2 .010$ F 0.9; $\beta -.16$ (ns) Depression $R^2 .178$ F 23.0; $\beta .37$ ($p < .001$) Reduced Motivation Anxiety – $R^2 .044$ F 4.9 $\beta .10$ (ns)	18

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
				Depression – R^2 .110 F 13.7; β .40 (p<.001) Mental Fatigue Anxiety – R^2 .254 F 36.2; β .39 (p<.001) Depression – R^2 .007 F 1.0 β -.01 (ns) Appetite and physical fatigue β 2.6 t 2.7 (p<.01) Nausea and general fatigue β .18 t 2.0(p<.05) Nausea and reduced activity β .23 t 2.3 (p<.05) Pain and general fatigue β .22 t 2.4 (p<.05) Pain and physical fatigue β .22 t 2.3 (p<.05) Breathing and general fatigue β .30 t 3.4 p<.001)	
Fink et al ²² (2009)	Cross-sectional research design Convenience sample of 87 heart failure patients recruited from 2 urban medical center outpatient Heart Failure clinics. Mean age 57 years Gender – 56% female	Pearson correlations	Fatigue Symptom Inventory – Interference Scale α = .93 Profile of Mood States Fatigue scale α = .90 Depression scale α .93	Depressed mood and fatigue Fatigue Symptom Inventory Interference Scale r=0.43 p = 0.01 Profile of Moods States – Fatigue Scale (POMS – F) r=0.64 p = 0.01	15

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
Goebel et al ⁶ (2009)	Secondary analysis 96 veterans who were outpatients at two hospitals and six affiliated community sites Mean age 67.2 years Gender – 4.2% female	Pearson coefficient correlations	Brief Pain Inventory – pain severity scale α 0.91 Patient Health Questionnaire – 2 (not reported) Generalized Anxiety Disorder – 2 (not reported)	Depression and pain $r = 0.32$ $p = 0.002$ Anxiety and pain $r = 0.33$ $p = 0.001$	20
Hertzog et al ⁵³ (2010)	Study design not reported 139 respondents recruited from patients receiving care at an outpatient heart failure clinic Mean age 70.6 years Gender – 24% women	Hierarchical cluster analysis,	Kansas City Cardiomyopathy Questionnaire α ranged from .78 to .89 except for self-efficacy which had an estimate of only .57 Heart Failure Symptom Survey (not reported)	Clusters 1) Shortness of breath, fatigue 2) Shortness of breath at rest, with activity, lying down, at night, fatigue, difficulty sleeping, forgetfulness 3) Fatigue, shortness of breath with activity, shortness of breath at rest, dizziness, bloating and swelling, depression	19
Jaing et al ³⁰ (2004)	Secondary analysis 291 heart failure patients admitted to the cardiology service of a university medical center Mean age 63 years 45.8% women	Pearson coefficient	Beck Depression Inventory (not reported) Spielberger State – Trait Anxiety Inventory (not reported)	Depression and anxiety State A – $r = 0.52$ $p < 0.01$ Trait A – $r = 0.59$, $p < 0.01$	19
Johansson et al ⁵⁵ (2010)	Study design not reported 331 subjects chosen	Principal component analysis with	Uppsala Sleep Inventory (not reported)	Clusters 1) Mean O ₂ saturation, dyspnea, NYHA class III, leg edema	19

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	from a previously investigated cohort of individuals living independently in a rural municipality Mean age 78 years 51% women	oblique rotation Structural equation modeling (SEM) Exploratory factor analysis Confirmatory factor analysis	Epworth Sleepiness Scale (not reported) Hospital Anxiety Depression Scale - Depression Scale(not reported) Dyspnea was assessed using a graded scale – 0 = no dyspnea, 10 = very severe dyspnea Polygraphic studies in the patient’s home	2) Excessive daytime sleepiness, fatigue, depression, non-restorative sleep	
Jurgens et al ⁵⁰ (2006)	Instrument Development Convenience sample of 201 community dwelling participants recruited from an urban tertiary care hospital, a suburban tertiary care hospital and a community hospital Mean age 70 years 44% female	Exploratory factor analysis using a principle component analysis with varimax rotation.	Heart Failure Somatic Awareness Scale (theta reliability = 0.71) Spielberger Trait Anxiety Inventory Form (not reported)	Factor 1 – orthropnea, dyspnea, paroxysmal nocturnal dyspnea Factor 2 – my feet were swollen, my shoes were tighter than usual Factor 3 – I could feel my heart beat faster, I felt pain in my chest, I had an upset stomach, I had a cough Factor 4 – fatigue, weight gain, dyspnea on exertion, unable to catch my breath	20
Jurgens et al ⁵¹ (2009)	Secondary Analysis 687 hospitalized patients with a confirmed diagnosis of heart failure drawn from a data registry of	Principal component analysis with oblique rotation Logistic	Minnesota Living with Heart Failure Questionnaire (Bartlett’s Test of Sphericity was reported as “acceptable”)	Inter-item correlation depression and worry .64 Inter-item correlation shortness of breath and fatigue low energy .65 Symptom Clusters Acute volume overload – shortness	21

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	the Heart Failure Quality of Life Trialist Collaborators. Mean age 71 years Gender – 51.7% female	Regression		of breath, tired, fatigue low energy, sleep problems Emotional cluster – depressed, worry, difficulty concentrating or remembering things Chronic volume overload – swelling, increased need to rest, dyspnea on exertion	
Lee et al ⁵⁴ (2010)	Secondary Analysis 331 patients. Data for this study were compiled from 3 prospective, longitudinal studies that had similar inclusion and exclusion criteria. Mean age 61 years Gender – “predominately male”	Cluster Analysis - hierarchical cluster agglomerative approach	Minnesota Living With Heart Failure Questionnaire (α .89)	Physical Symptom Cluster – dyspnea, fatigue increased need to rest, fatigue low energy, sleep disturbance Emotional/Cognitive Symptom Cluster – worrying, feel depressed, cognitive problems	17
Lesman-Leegte et al ³⁷ (2006)	Secondary analysis of data from a randomized control trial 572 patients hospitalized with heart failure Mean age 71 ± 12 years 38% women	Multivariable logistic regression	Center for Epidemiological Studies Depression Scale (not reported) Heart Failure symptoms were assessed by interview using 10 structured questions	Depression and sleep – odds ratio = 3.45, p 0.001 Depression and loss of appetite – odds ratio = 2.28 95%, p = 0.001.	18

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
Luyster et al ³¹ (2009)	Cross-sectional design 88 patients with a primary diagnosis of heart failure and treated with an ICD who were recruited from a large cardiology practice Mean age 70 years 23% female	Pearson correlation	Beck Depression Inventory (not reported) State-Trait Anxiety Inventory – Trait (not reported)	Depression and anxiety $r = 0.66$ $p < .01$ Depression and dietary adherence $r = -0.51$ $p < .01$	15
Paukert et al ³² (2009)	Study design not reported 104 Heart Failure patients identified using the outpatient and patient treatment databases of a large VA hospital Mean age 71.7 years Gender – 1% female	Spearman's rho correlations Pearson product-moment correlations	Geriatric Depression Scale (not reported) Geriatric Anxiety Inventory (not reported)	Depression and anxiety $r = 0.65$ $p < 0.01$	16
Principe-Rodriguez et al ⁴² (2005)	Design not reported 201 participants who were new patients referred to the heart failure program at a university hospital Mean age 57.7 years 48% women	Pearson coefficient	Cleveland Sleep Habits Survey (not reported)	Paroxysmal nocturnal dyspnea and sleepiness $r = 0.161$ $p = 0.02$ Paroxysmal nocturnal dyspnea and insomnia risk $r = -0.279$ $p = 0.004$	19
Ramasamy et al ⁴⁰ (2006)	Study design not reported 67 individuals with	Pearson's product moment	Dyspnea Scale (not reported) Hospital Anxiety and	Depression and dyspnea $r = 0.40$ $df = 46$ $p < 0.01$ Depression and sleep disturbance	14

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	<p>chronic, non-valvular heart failure who were well-compensated ambulatory patients or recently hospitalized with decompensated heart failure.</p> <p>Mean age 65.4 years Gender – 32% women</p>	correlations	<p>Depression Scale (not reported)</p> <p>Abbreviated visual-analog scale to assess severity of fatigue (0-10, not at all to a great deal) (not reported)</p> <p>A questionnaire about perception of health status adapted from the general health items on the SF-36</p> <p>Sleep questionnaire adapt from the Generalized Anxiety Disorder module of the Structured Clinical Interview for DSM-IV Axis 1 Disorders (not reported)</p>	<p>r = 0.26 df 41 p<0.11</p> <p>Anxiety and dyspnea r=-0.04 df 46 p<0.77</p> <p>Fatigue and dyspnea r= 0.62 df 36 p<0.01</p> <p>Dyspnea and sleep disturbance r = 0.26 df 41 p<0.11</p> <p>Dyspnea and fatigue r = 0.62 df 36 p <0.01</p> <p>Depression and fatigue – correlation was not statistically significant.</p>	
Redeker ²³ (2006)	<p>2 group cross-sectional design</p> <p>61 HF patients recruited from an outpatient heart failure program and 57 comparison group participants recruited from the surrounding community</p>	Bivariate correlations	<p>Centers for the Epidemiological Studies of Depression ($\alpha = .88$)</p> <p>Profile of Mood States – Short Form</p> <p>Dejection (Depression scale $\alpha = .92$, Tension Anxiety scale $\alpha = .88$)</p> <p>Hospital Anxiety and</p>	<p><u>Dyspnea</u></p> <p>Depression CESD r = 0.41, p<.001</p> <p>Depression POMS DD r=0.17 p 0.70</p> <p>Depression HADS r=0.03 p .790</p> <p><u>Fatigue</u></p> <p>Depression CESD r=.57 p<.001</p> <p>Depression POMS DD r=.34 p<.001</p> <p>Depression HADS r=.01 p .953</p> <p><u>Sleep Disturbance</u></p> <p>Depression CESD r = 0.53 p<.001</p>	18

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	<p>Mean age of the heart failure group 58.7 years</p> <p>Mean age of the comparison group 55.25 years</p> <p>Gender – Heart failure group 37.7% women, comparison group 52.6% women 61 HF patients and</p>		<p>Depression Scale (Anxiety scale $\alpha = .82$, Depression scale $\alpha = .72$)</p> <p>Pittsburgh Sleep Quality Index (not reported)</p> <p>Epworth Sleepiness Scale ($\alpha = .87$)</p> <p>Multidimensional Assessment of Fatigue Scale ($\alpha = .91$)</p> <p>Multidimensional Assessment of Dyspnea ($\alpha = 0.94$)</p>	<p>Depression POMS DD $r=0.39$ $p<.001$</p> <p>Depression HADS $r=0.19$ $p=.06$</p> <p><u>Excessive Daytime Sleepiness</u></p> <p>Depression CESD $r=.28$ $p=.003$</p> <p>Depression POMS DD $r = 0.30$ $p .002$</p> <p>Depression HADS $r =- 0.02$ $p = .855$</p> <p><u>Dyspnea</u></p> <p>Anxiety POMS – Tension Anxiety $r=0.22$ $p=0.20$</p> <p>Anxiety HADS $r=0.35$ $p=.001$</p> <p><u>Fatigue</u></p> <p>Anxiety POMS – TA $r=0.42$ $p<.001$</p> <p>Anxiety HADS $r=0.49$ $p<.001$</p> <p><u>Sleep Disturbance</u></p> <p>Anxiety POMS – TA $r = 0.41$ $p<.001$</p> <p>Anxiety HADS $r=0.42$ $p<.001$</p> <p><u>Excessive Daytime Sleepiness</u></p> <p>Anxiety POMS – TA $r=0.30$ $p = .002$</p> <p>Anxiety HADS $r=0.16$ $p = .110$</p>	
Redeker et al ³³ (2010)	<p>Cross-sectional observational</p> <p>173 stable chronic heart failure patients recruited from 5 heart failure disease management programs</p> <p>Mean age 60.3 years</p>	<p>Logistic Regression</p> <p>Confirmatory factor analysis</p>	<p>Multi-Dimensional Assessment of Fatigue ($\alpha = .91$)</p> <p>Center for Epidemiological Studies Depression Scale ($\alpha=.89$)</p> <p>Epworth Sleepiness</p>	<p>Patients with difficulty initiating and maintaining sleep were more likely to be depressed OR = 5.09 95% CI = 2.41 – 10.75.</p> <p>“The latent variable daytime symptoms (fatigue, depression, and excessive daytime sleepiness) had high factor loadings on fatigue(0.724</p>	20

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	Gender – 34.7% women		Scale ($\alpha=.78$ Sleep Habits Questionnaire ($\alpha = 0.83$) Items evaluating perception of quantitative sleep characteristics taken from the Pittsburgh Sleep Quality Index (not reported) Nocturnal polysomnography	SE = 0.056) and depression (0.716, SE = 0.057)	
Redeker et al ³⁶ (2010)	Cross-sectional observational design 170 patients with stable chronic HF recruited from 5 structured disease management programs Mean age 60.3 years Gender – 35.5% women	Linear and logistic regression	6 minute walk test Epworth Sleepiness Scale ($\alpha .77$) Multi-Dimensional Assessment of Fatigue ($\alpha = .93$) Center for Epidemiological Studies Depression Scale ($\alpha = .84$) Pittsburgh Sleep Quality Index ($\alpha = .79$) Wrist actigraph to measure daytime mobility Nocturnal polysomnography	Depression and severity of sleep disordered breathing – no statistically significant relationship Fatigue and severity of sleep disordered breathing – no statistically significant relationship Sleep quality and severity of sleep disordered breathing – no statistically significant relationship Excessive daytime sleepiness and severity of sleep disordered breathing – no statistically significant relationship	19

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
Sauve et al ⁴⁴ (2009)	Case-control design 50 patients with heart failure recruited from heart failure and cardiology clinics of a university hospital and 50 controls recruited from active adult communities Mean age: Heart failure patients 63 years, Controls 62 years. Gender: Heart failure patients 48% female, Controls 54% female	Pearson's r correlation, Kendall tau	Neurobehavioral Cognitive Status Examination Symbol-Digit Modalities Test Rey Auditory Verbal; Learning Test (not reported) Controlled Word Association Test (not reported) Finger Tapping Test (not reported) Memory Scanning Test (not reported) Visual Scanning Task(not reported) WAIS-R Vocabulary Subtest (not reported) The Mental Health Index (not reported) Short Form – 36 (not reported)	No statistically significant relationships between Depression and orientation Depression and attention Depression and memory Depression and motor performance Anxiety and orientation Anxiety and attention Anxiety and memory Anxiety and motor performance Pain and orientation Pain and attention Pain and memory Pain and motor performance	15
Skobel et al ³⁴ (2005)	Study design not reported 51 patients of an ambulatory clinic referred for evaluation of heart failure or consideration for a	Spearman rank correlation coefficients	Beck Depression Inventory (not reported) Pittsburg Sleep Quality Index (not reported) SF 36 – German version (not reported)	In heart failure patients with sleep disordered breathing Depression and sleep quality $r = 0.71$ $p = 0.001$ Depression and apnea hypopnea index $r = 0.88$ $p = 0.001$ Pain and apnea hypopnea index	15


Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	heart transplant <u>Mean age</u> of heart failure patients with a sleep-related breathing disorder - 61+ 11 years <u>Mean age</u> of heart failure patients without sleep-related breathing disorder - 52+10 29% female		Bicycle exercise protocol Overnight polygraphy	$r = -0.53$ $p = 0.006$	
Smith et al ⁵² (2009)	Prospective study design 381 CHF patients from an outpatient heart failure clinic Mean age 65.7 ± 10.4 years Gender 27.8% female	Principal component analysis with oblimin rotation, latent class cluster analysis	Maastricht questionnaire (not reported) Minnesota Living with Heart Failure Questionnaire (not reported)	Clusters 1) Absence of vital exhaustion 2) Manifest vital exhaustion, relative absence of cognitive affective, depressive symptoms, relative absence of sleep difficulties, decreased concentration, increased levels of fatigue 3) Manifest vital exhaustion, cognitive affective depressive symptoms, sleep difficulties	15
Smith et al ²⁶ (2007)	Prospective study design 136 CHF patients from an outpatient heart failure clinic Mean age 65.6 years Gender - 23.5%	Pearson's product moment correlation coefficients	Dutch Exertion Fatigue Scale (not reported) Fatigue Assessment Scale (not reported) Health Complaints Scale somatic complaints	Exertion Fatigue and: Sleep problem $r = .29$ $p = .001$ Dyspnea $r = .33$ $p < .001$ Cardiac pain $r = .42$ $p < .001$ Depressive symptoms $r = .33$ $p < .001$ General Fatigue and: Sleep problem $r = .42$ $p < .001$	16

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	women		subscale(not reported) Beck Depression Inventory (not reported) DS14 (not reported)	Dyspnea $r = .45$ $p < .001$ Cardiac pain $r = .44$ $p < .001$ Depressive symptoms $r = .42$ $p < .001$	
Song et al ⁷ (2010)	Prospective Study 421 heart failure patients recruited during an index hospitalization chronic heart failure exacerbation at tertiary medical centers Mean age 62 39.7% female	Agglomerative hierarchical clustering approach with Ward's method,	Memorial Symptom Assessment Scale – Heart Failure ($\alpha = .81$)	Dyspneic symptom cluster – shortness of breath, difficulty breathing while lying flat, waking up breathless at night Weary symptom cluster – lack of energy, lack of appetite, difficulty sleeping	17
Stephen ³⁹ (2008)	Descriptive, cross-sectional and correlational design Convenience sample of 53 patients with a diagnosis of HF and with documented systolic dysfunction recruited from outpatient clinics Mean age 77 years 32% women	Pearson product moment correlations	Profile of Mood States fatigue scale ($\alpha = .82$) Visual analogue scale for fatigue (not reported) Revised symptom checklist (not reported)	Fatigue Intensity and Breathless with activity $r = .25$ $p < .05$ Fatigue Intensity and Difficulty sleeping $r = .20$ $p > .05$ Fatigue Intensity and nocturia $r = .36$ $p < .01$	19
Sullivan et al ²⁴ (2004)	Prospective design 142 subjects and 88 spouses provided	Partial correlations	Hamilton Depression Rating Scale (not reported)	Baseline/ Follow-up Correlations Depression and bodily pain $-.37$ $p < .001$ / $-.47$ $p < .001$	21

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	<p>baseline data. A total of 113 subjects and 70 spouses also completed a follow-up assessment 6 to 12 months after baseline. Subjects were recruited from a heart failure/pretransplant clinic of a university medical center Mean age 53.6± 10 years Gender – 22.1% female</p>		<p>Hamilton Anxiety Rating Scale (not reported) System Checklist 20 Depression Scale (not reported) SF – 36 (not reported) Multidimensional Assessment of Fatigue Scale (not reported)</p>	<p>Depression and vitality -.44 p<.001/-.40 p<.001 Depression and breathlessness .56 p<.001/ .43 p<.001 Depression and chest pain .38 p<.001/.43 p<.001 Depression and fatigue .62 p<.001/ .57 p<.001</p>	
Tang et al. ²⁵ (2010)	<p>Cross-sectional correlation design Convenience sample of 107 CHF patients recruited from a cardiology ward of a medical center Mean age 69.3 years Gender – 46.7% women</p>	Pearson correlations	<p>Tang Fatigue Rating Scale (total scale – $\alpha = .96$, physiological scale – $\alpha = .83$, psychosocial scale – $\alpha = .92$, daily-living activities – $\alpha = .97$) Fatigue Visual Analog Scale (total scale $\alpha = .89$, energy subscale $\alpha = .97$, fatigue subscale $\alpha = .89$) Beck Depression Inventory ($\alpha 0.81$)</p>	Depression and fatigue $r=0.77$ $p<0.01$	22
Wang et al. ³⁵ (2010)	Predictive correlational design	Pearson product	Perceived Health Scale ($\alpha = .73$)	Depression and sleep quality $r = .402$ $p<0.001$	21

Citation	Design/Sample/Setting	Analytic Techniques	Instrumentation (Psychometrics in the present study)	Symptom Relationships	STROBE Score
	Convenience sample of 101 patients with heart failure recruited from cardiology clinics Mean age – 74 years Gender 32.8% female	moment correlations	Short Form Geriatric Depression Scale ($\alpha = .82$) Pittsburg Sleep Quality Index ($\alpha = .80$)		
Webel et al ⁴¹ (2007)	Descriptive correlational design 48 heart failure patients who were referred to the investigators by cardiologists and nurse practitioners- data collected for 30 days. Mean age 48 ± 15 years Gender – 46% female	Spearman correlation coefficients	Symptom Rating Scales – 10 point analog scale with 0 indicating absence of dyspnea or edema and 10 indicating the worst dyspnea or edema (not reported)	Dyspnea and edema – correlations calculated for each day of a 30 day period ranged from 0.45 to 0.63 $p \leq 0.001$ for all correlations	15
Yu et al ³⁸ (2004)	Cross-sectional study design. 227 patients with an index diagnosis of CHF were recruited from a medical unit of a university-affiliated hospital Mean age 77.1 years 52.4% women	Pearson product-moment correlation	Hospital Anxiety and Depression Scale – Chinese version (not reported) Chronic Heart Failure Questionnaire dyspnea and fatigue subscales - Chinese version (not reported)	Anxiety/Depression and dyspnea $r = -.17$ $p < .05$ Anxiety/Depression and fatigue $r = -.50$ $p < .001$	16

Figure 1. Search and Screening Results



Heart Failure Symptom Clusters and Functional Status

Janet K. Herr, RN, PhD

Jeanne Salyer, Ph.D., RN, Associate Professor
School of Nursing, Virginia Commonwealth University

Maureen Flattery, MS, RN, ANP, Heart Failure/Transplant Coordinator,
Pauley Heart Center
Virginia Commonwealth University Health System

Lauren Goodloe, Ph.D., RN, Clinical Associate Professor
Debra E. Lyon, Ph.D., RN, FNP-BC, FAAN, Professor School of Nursing
Virginia Commonwealth University

Christine Schubert Kabban, Ph.D, MBA, Assistant Professor
Department of Mathematics and Statistics/ENC
Air Force Institute of Technology
Wright-Patterson Air Force Base, Ohio

Dolores G. Clement, Dr.P.H. FACHE, Professor
Department of Health Administration, School of Allied Health Professions
Virginia Commonwealth University
Richmond, VA

Corresponding author: Janet K. Herr, Ph.D
School of Nursing, Virginia Commonwealth University
1100 East Leigh Street
Richmond, Virginia 23298-0567
Phone: 804-323-8281 Fax: 1-866-891-1105 E-mail: jkherr@vcu.edu

Acknowledgements: Support for this study was provided by The Center of Excellence for Biobehavioral Approaches to Symptom Management (P30 NRO11403), Mary Jo Grap, Ph.D., RN, ACNP, FAAN, Principle Investigator, Department of Adult Health and Nursing Systems, School of Nursing, Virginia Commonwealth University

There are no conflicts of interest.

Funding was not received for the conduct of this study.

6 Tables; 1 Figure

Abstract

Background: Clinical assessment of heart failure includes symptom identification and the evaluation of the relationship of symptoms to functional status. Symptom clusters are groups of at least 2 or 3 co-occurring symptoms that are related but are independent of other groups of symptoms. Since the publication of the NYHA functional classification symptom occurrence in cardiac conditions has been linked to functional status.

Objectives: The objectives of this study are to: (1) examine relationships among symptoms commonly experienced by individuals with heart failure, (2) identify symptoms that form clusters, and (3) evaluate the impact of heart failure symptom clusters on attributes of functional status: limitations and mobility

Methods: The Theory of Unpleasant Symptoms guided the conduct of this study. A convenience sample of individuals ($n = 117$) with a confirmed diagnosis of heart failure was recruited from an academic medical center. Prevalent symptoms (anxiety, depression, dyspnea, peripheral edema, pain, appetite, hunger, cognitive dysfunction and daytime function) were evaluated along with the outcome variables, functional limitations and mobility. Principle components analysis was used to extract symptom clusters and regression analysis was used to evaluate the relationship between the symptom clusters, their interaction terms, and the demographic variables, age and co-morbidity.

Results: Three symptom clusters, sickness behavior, discomforts of illness, and GI distress were extracted. Predictors of functional limitations ($F = 35.96$, $p = 0.0005$, $R^2 = 0.578$) included sickness behavior ($\beta = -.681$, $p \leq 0.0005$), discomforts of illness ($\beta = -.765$, $p = 0.0005$) and the interaction term between these two symptom clusters ($\beta = .649$, $p \leq 0.014$). This model predicted 59% of the variance in functional limitations. Predictors of limited mobility ($F =$

20.68, $p = 0.0005$, $R^2 = 0.275$) included sickness behavior ($\beta = -0.441$, $p \leq 0.0005$) and co-morbidity ($\beta = -0.200$, $p \leq 0.019$). This model predicted 28% of the variance in mobility.

Conclusions: Discomforts of illness and sickness behaviors were strong predictors of functional limitations. Functional limitations increase as discomforts of illness increase, but increases at a faster rate when sickness behaviors are increased. Sickness behavior and co-morbidity were predictors of limited mobility. Symptom management strategies focused on the symptom clusters may improve functional limitations and mobility for individuals experiencing heart failure symptoms.

Background

Heart failure is a progressive clinical syndrome that is estimated to affect approximately 6.6 million Americans over age 18 years and is projected to increase by 3 million people by 2030.¹ The ongoing clinical assessment of individuals with heart failure includes symptom identification and evaluation of the relationship of symptoms to functional status.² Symptoms have been referred to as threats to health³ and a stimulus for entry into the medical system.⁴

Symptoms may occur singly or in clusters of two or more. Dodd, Miaskowski and Paul⁵ and Kim, McGuire, Tulman and Barsevick⁶ characterize symptom clusters as a group of at least 2 or 3 symptoms that *co-occur, are related but are independent of other groups of symptoms*. The symptoms in a symptom cluster *may be related to a common source* such as a disease, treatment, life transition, or co-morbid conditions.⁶ Concurrence, stability, and relationship are important characteristics of a symptom cluster.

The influence of a symptom cluster on one or more patient outcomes may be useful in determining whether a symptom cluster is clinically significant.⁷ Since 1928, when the New York Heart Association (NYHA) published a classification of clinical cardiac disease severity based on the occurrence of symptoms with physical activity, functional status has been associated with symptoms of cardiac disease.⁸ Leidy⁹ and Wang¹⁰ concur that the term functional status refers to activities performed to meet basic needs, carry out usual roles, and maintain health and well-being. Additionally, Wang¹⁰ proposes that functional status is demonstrated by the performance of activities including physical, psychological, social, spiritual, intellectual, and roles associated with daily living.

In the heart failure patient population, relatively few studies have been conducted to identify symptom clusters. Fatigue and dyspnea is a pair of symptoms that have been included in

clusters reported by several investigators. For example, fatigue and dyspnea are included in symptom clusters with forgetfulness,¹¹ sleep disturbance,^{12,13} dizziness, and depression,¹¹ and manifestations of edema such as weight gain,¹⁴ and swelling.^{11,12} Dyspnea, dizziness, swelling and bloating formed a symptom cluster with fatigue and depression.¹¹ Associations between fatigue and pain^{15,16} and appetite loss¹⁵ are also reported by investigators conducting studies of single symptoms but they have not appeared in the heart failure symptom clusters reported at this point.

The relationship between fatigue and depression has been supported by several investigators.¹⁶⁻²¹ A cluster of symptoms which include fatigue, depression, and daytime sleepiness has been identified by Riegel and Ratcliff²² and is characterized as sickness behavior. As well, fatigue and depression have both been included in a symptom cluster with sleep disturbances such as non-restorative sleep,²³ excessive daytime sleepiness^{23,24} and sleep difficulties.²⁵ Investigators have also observed that depression and anxiety have been included in symptom clusters with cognitive problems.¹³

Evaluation of the clinical significance of a symptom cluster may include an assessment of the effect of the symptom cluster on important patient outcomes.⁷ For example, investigators have observed these relationships with outcomes such as death,^{13,26} re-hospitalization,^{13,25,26} health status,²⁵ and global perceived health.²³ In individuals with NYHA class IV heart failure, the sickness behavior symptom cluster was a predictor of the risk of death or hospitalization.²² The relationship between a cluster of heart failure symptoms and functional status is not reported in the literature.

The purpose of this study is to examine the presentation of symptom clusters in individuals who have heart failure and evaluate the relationship of the identified clusters to

functional status in these individuals. Specific aims are to: (1) examine symptoms commonly experienced by individuals with heart failure, (2) identify symptoms that form clusters, and (3) evaluate the impact of heart failure symptom clusters on two attributes of functional status: limitations and mobility.

Conceptual Framework

The Theory of Unpleasant Symptoms (TUS)³ illustrates the experience of concurrent symptoms. In a symptom cluster, concurrence refers to symptoms occurring together—for example, the occurrence of dyspnea, fatigue, and edema in an individual who has heart failure.⁶ The relationship among symptoms portrayed by the TUS is described as *multiplicative* instead of *additive* and symptoms may interact to augment the effects of each other. For example, several symptoms may individually have limited effect on an outcome, whereas, in a cluster, the impact may be highly significant. Model concepts included in the TUS are *influencing factors* that may trigger or shape the symptom experience (e.g., age and co-morbidities), *symptoms* experienced by an individual (e.g., dyspnea, fatigue, edema, sleep and cognitive disturbances, pain, anxiety, depression, hunger and appetite disturbances), and the *consequences of the symptom experience* (functional and mobility limitations). Consequences of the symptom experience are performance-related and are hypothesized to include effects on physical activity, activities of daily living, social activities, and role performance. The conceptual framework for this study, based on the TUS, is presented in Figure 1.

Methods

Design, sample and setting

A convenience sample of individuals (n=117) with a confirmed diagnosis of heart failure was recruited from the Heart Failure/Heart Transplant Program of Virginia Commonwealth

University. Potential participants in this cross-sectional study were 18 years of age or older, NYHA functional class II – IV, had been receiving standard therapy for heart failure for at least 3 months prior to enrollment, and were able to speak and read English well enough to complete the questionnaires. Standard therapy for heart failure includes beta blocker therapy, diuretics, and angiotensin converting enzyme inhibitor therapy.² Therapy for patients with symptomatic heart failure who have reduced left ventricular ejection fraction (<35%), a history of cardiac arrest, ventricular fibrillation or hemodynamically destabilizing ventricular tachycardia may also include an implantable cardioverter-defibrillator.²⁷

Data collection procedures

The study was approved by the Virginia Commonwealth University Institutional Review Board. After obtaining informed consent, study participants received a study packet containing a cover letter with instructions about completing the study materials, a copy of the consent form, and the study questionnaires. Study staff reviewed the contents of the packet with each participant and answered questions about completion of the questionnaires. Participants completed the questionnaires on the day of their clinic visit.

Measurement of Variables

Demographic variables (gender, age, education, employment and marital status) were self-reported on a demographic questionnaire. Clinical characteristics (NYHA functional classification of heart failure, ejection fraction [EF], etiology of heart failure, time since diagnosis of heart failure, prescribed medications, and co-morbidities) were collected from the medical record. The Charlson Co-morbidity Index,²⁸ which provides a weighted taxonomy of co-morbid conditions with prognostic implications, was used to summarize co-morbid conditions.

Dyspnea was measured using the shortness of breath subscale of the Cardiovascular Limitations and Symptoms Profile (CLASP).²⁹ The subscale consists of 5 questions assessing patient perception of shortness of breath. Scale items evaluating frequency, severity, and interference with activities are categorized as mild, moderate or severe (1=mild, or not at all; 5=severe, or a greater frequency). In individuals with coronary heart disease, internal consistency reliability, reported by Lopez, Chair, Thompson, Tsui, and Lewin,³⁰ is 0.70 for the CLASP and 0.82 for the shortness of breath subscale. In the present study, Cronbach's alpha for the shortness of breath subscale was 0.85.

Fatigue was evaluated using the Brief Fatigue Inventory³¹ which is composed of 9 items—three items assessing fatigue severity and 6 items assessing the amount of interference with aspects of life associated with fatigue. Items are rated on a 0 (no fatigue) to 10 (fatigue as bad as you can imagine) scale. Cronbach's alpha for the Brief Fatigue Inventory is reported to be 0.96. In this study, internal consistency reliability was 0.95.

Consistent with the strategy used by Webel and colleagues³² to evaluate daily variation of edema in heart failure patients, a visual analogue scale was selected to assess peripheral edema. The scale was anchored by the descriptors none (0) and worst I have ever had (10). Study participants were also asked to report the location of their edema (1=feet and ankles; 2=up to the knees; 3=above the knees).

The Hospital Anxiety and Depression Scale (HADS)³³ was used to assess anxiety and depression. The HADS is designed to provide reliable screening for clinically significant anxiety and depression. Each subscale is composed of 7 items on a 4-point Likert scale (0=absence; 3=occurrence most of the time). Chen et al.¹⁷ reported Cronbach's alpha for the HADS to be 0.78. Internal consistency reliability for the anxiety and depression subscales was 0.73 and 0.95,

respectively. Cronbach's alpha for the HADS in this study is 0.89 and for the anxiety and depression subscales, 0.84 and 0.81, respectively.

The Bodily Pain Scale of the Medical Outcomes Study (MOS) 36 – Item Short-Form Health Survey (SF-36)³⁴ was selected to evaluate pain. The 4 items assess the frequency of bodily pain or discomfort and the extent of interference with normal activities. Item responses range from 1 indicating none or very mild to 5 indicating frequent occurrence or very severe. McHorney, Ware and Raczek³⁵ reported a reliability of 0.77 for the SF-36 Bodily Pain Scale. In this study Cronbach's alpha for the Bodily Pain Scale was 0.83.

Appetite, the psychological desire for certain foods, and hunger, a physical sensation indicating the need to eat, were assessed using two items of the Food, Eating Experiences, and Diet (FEED) instrument.³⁶ The two items included visual analogue scales measuring hunger and appetite (1=no hunger sensations or no appetite/cravings; 10=hungriest I have ever been or my appetite was extremely good/strong cravings).

The General Sleep Disturbance Scale³⁷ is composed of 21 items evaluating initiation insomnia, maintenance insomnia, sleep quality, sleep quantity, daytime functioning, and medication/self help for sleep. Responses to each item are captured using an 8 point Likert scale (0 = no days; 7=every day). A Cronbach's alpha of 0.88 has been reported for the instrument.³⁷ In this study, only the daytime functioning subscale (5 items) was used. Excessive daytime sleepiness, a consequence of sleep disturbance, was chosen as a variable in this study because of its prevalence in individuals with heart failure³⁸ and association with worse functional status and lack of physical activity.³⁹ In the present study, Cronbach's alpha on this subscale was 0.74.

Cognitive impairment was assessed using the MOS Cognitive Function Scale.⁴⁰ which provides an assessment of the day-to-day difficulties with cognitive functioning of which an

individual has awareness. The 6-item Likert scale (1=all the time; 6=none of the time) was recoded so that higher scores reflected more difficulty with cognitive functioning. Hays, Sherbourne and Mazel⁴¹ reported Cronbach's alpha for the Cognitive Function Scale is 0.88. Cronbach's alpha was 0.91 in this study.

The outcome variable, functional status, conceptualized as functional limitations and mobility limitations, was evaluated using the MOS Physical Functioning Scale.⁴² Categories of activities in this evaluation of functional status include self-care limitations, mobility, and physical activities. There are 10 items that evaluate physical limitation (1=limited a lot, 2 = limited a little, 3 = not limited at all), and 3 items addressing mobility or independence in physical activity (1 = difficulty all of the time or every day, 5 = difficulty none of the time or never). Hays, Sherbourne and Mazel⁴¹ reported a Cronbach's alpha of 0.93 for the physical limitations scale and 0.74 for the mobility scale. In this study Cronbach's alpha for the physical limitations subscale was 0.91 and for the mobility subscale Cronbach's alpha was 0.65.

Statistical Analysis

Data were analyzed using descriptive statistics, Pearson's correlations, principle components analysis, and multiple regression analyses. The level of significance was set at $p < 0.05$. Histograms and scatter plots of variables were examined for skew, gaps in distribution of the data, multiple peaks, and outliers. We examined the frequency and percent of symptoms to confirm prevalence and co-occurrence. Correlation analysis using Pearson's r was used to confirm that the symptoms were related. The study sample was characterized using demographic and clinical variables. Means and standard deviations are reported for continuous variables and frequencies and percents are reported for categorical variables.

Principle components analysis (PCA) was used to identify symptoms forming clusters. To improve interpretability, we used oblimin rotation with Kaiser normalization. A 3-component solution was obtained. Loadings were evaluated to determine the symptoms retained on each component. After identification of the symptom clusters—characterized as sickness behaviors (5 symptoms), discomforts of illness (3 symptoms), and gastrointestinal (GI) distress (2 symptoms)—a composite score was calculated for each symptom cluster. This score was calculated by multiplying each symptom (scale/subscale) mean by the loading and summing the products of the calculation.

Assumptions of regression (normality of residuals and homogeneity of variance) were examined and two multiple regression analyses were performed with functional limitations and mobility as the dependent variables of each respective analysis.

Results

Sample characteristics

Participants (n=117) were 62% male and had a mean age of 56 years. A little less than half (42%) were receiving disability. Etiology of heart failure was primarily ischemic (48%) or idiopathic (43%). Mean EF was 25%. On average, time since the onset of heart failure was 5 years. The majority of the study participants were NYHA functional class 2 (44%) or class 3 (39%) and 13% were class 4. The most prevalent co-morbidities were diabetes mellitus (44%), renal disease (32%), chronic pulmonary disease (28%), and connective tissue disease/arthritis (21%). Twenty six participants (13.6%) had only heart failure. Over half of the study participants (62%) had an implanted cardiac defibrillator (ICD); 9% were receiving intravenous inotrope infusions; 21 were either listed or being evaluated for heart transplantation, and 10 were

being evaluated for mechanical circulatory support. Demographic and clinical characteristics are presented in Table 1.

All of the individuals in this study reported having shortness of breath at least once or twice during the previous 2 weeks. Some other symptoms with a high prevalence in this sample of individuals with heart failure are decreased hunger (95%), decreased appetite (85%), fatigue (73%), edema (73%), and pain (62%). Symptom prevalence is presented in Table 2.

Principal Components Analysis (PCA)

The PCA solution revealed 3 symptom clusters, characterized as sickness behaviors (5 symptoms), discomforts of illness (3 symptoms), and gastrointestinal (GI) distress (2 symptoms). The factor loadings are presented in Table 3. In Table 4 descriptive statistics for the model variables are presented for each cluster. Descriptive statistics for the symptom cluster composite scores may be found in Table 5.

Sickness Behavior. The sickness behavior cluster (mean=58.89; $SD=25.16$) was composed of anxiety, depression, daytime functioning, cognitive dysfunction, and fatigue. The most highly correlated among these symptoms are depression and fatigue ($r = 0.652, p \leq 0.0005$) and depression and anxiety ($r = 0.648, p \leq 0.0005$). Depression was correlated at the same level with daytime function ($r = 0.513, p = 0.0005$) and cognitive dysfunction ($r = 0.513, p \leq 0.0005$). Cognitive dysfunction was inversely correlated with anxiety ($r = -0.567, p \leq 0.005$), fatigue ($r = -0.469, p \leq 0.0005$) and daytime function (sleepiness) ($r = -0.432, p \leq 0.0005$).

Although the mean scores on the HADS of 7.67 ($SD=4.28$) on the depression subscale and 8.28 ($SD=4.64$) on the anxiety subscale fall into the doubtful category of anxiety and depression as defined by Zigmond and Snaith³³ it is notable that 32% of the participants had scores of 11 or higher on the anxiety subscale, indicating definite cases of anxiety, and 21% had

scores of 11 or higher on the depression subscale indicating definite cases of depression. Scores on the daytime functioning sub-scale of the General Sleep Disturbance Scale ranged from 0 to 19 out of a possible high score of 35. The mean daytime functioning score was 17.75 ($SD = 7.29$), which indicates some limitation in daytime function due to sleepiness every day of the week. Only 35% of the study participants reported feeling alert and energetic during the day on 4 or more days during the week. Feeling sleepy during the day on 4 or more days during the week was reported by 68% of study participants and 45% of study participants reported struggling to stay awake during the day for 4 or more days during the week.

The mean score on the cognitive functioning variable of 27.85 ($SD = 6.81$) out of a possible score of 36 (denoting good cognitive function) indicates difficulty some of the time with forgetfulness, concentrating, and problem solving. In fact 19% of the participants reported having difficulty with problem solving a good bit of the time to all of the time and 8.5% reported having difficulty with concentration or difficulty paying attention a good bit of the time to all of the time. Slow reactions or difficulty with forgetfulness were reported by 11.1% of participants a good bit or all of the time. Confusion a good bit of the time to all of the time was reported by 6% of participants.

Participants experienced a moderate level of interference with activities related to fatigue (mean 30, $SD = 17.59$). When asked to rate their fatigue “right now” on a visual analog scale, 14% of participants reported no fatigue at the time of the survey. However, over half (57%) rated their fatigue “right now” as 5 or higher on a 0-10 visual analogue scale. The usual level of fatigue during the past 24 hours was rated between 3 and 8 by 67% of study participants. This sample of heart failure patients rated fatigue interference with walking (64%), normal work (63%), and general activity (55%) as 5 or higher on a visual analogue scale.

Discomforts of illness. Shortness of breath, edema, and pain are the symptoms included in the discomforts of illness symptom cluster (mean=16.03; $SD=8.69$). The most highly correlated symptoms in this symptom cluster are shortness of breath and pain ($r = 0.484, p \leq 0.0005$) followed by shortness of breath and edema ($r = 0.426, p \leq 0.0005$) and edema with pain ($r = 0.414, p \leq 0.0005$).

On average, participants experienced moderate limitation associated with shortness of breath (mean=9.23; $SD=3.61$). While mostly good days or a mix of good and bad days was reported by 90% of the participants, 51% reported moderate to severe shortness of breath over the previous 2 weeks and 68% reported a moderate amount or a great deal of difficulty lying flat without becoming short of breath.

Participants reported a moderate level of peripheral edema (mean=3.91, $SD 3.63$). Peripheral edema was reported primarily in the feet and ankles (47.9%). Peripheral edema up to the knees was reported by 16.2% of the participants.

In this sample pain caused a moderate level of interference with activities (mean 8.1, $SD 6.98$). Pain was described as moderate to very severe by 78% of study participants who reported pain and was reported to last more than 2 days by 16% of the individuals reporting pain. Pain interfered at least a moderate amount with normal work for 63% of those who had pain. The sites of pain were lower extremities (41%), chest (37%), back (16%), upper extremities (12%) and abdomen (10%).

GI Distress. The third symptom cluster, GI distress, is composed of appetite, and hunger. On average participants experienced some loss of appetite (mean=6.46, $SD=2.74$) and hunger (mean=5.01, $SD=2.66$). Appetite and hunger were moderately correlated ($r = 0.416, p \leq 0.0005$). In this sample of individuals with heart failure 85% reported decreased appetite and 95%

reported decreased hunger. Among the study participants 32% felt full after eating a small amount of food and 26% said that being short of breath had an effect on the amount of food eaten. The limited variety of foods and the number of pills taken were reported by 20% to have an effect on the amount of food eaten.

There was a moderate correlation between the discomforts of illness symptom cluster and the sickness behavior symptom cluster ($r = 0.661, p \leq 0.0005$). The GI distress symptom cluster did not correlate with either the discomforts of illness symptom cluster ($r = -0.011, p = 0.455$) or the sickness behavior symptom cluster ($r = -0.059, p = 0.272$).

Regression analysis

Functional limitations and mobility were outcome variables in the regression analysis. Participants had a moderate amount of functional limitation (mean= 16.10, $SD = 5.09$) and a moderate amount of limitation in mobility (mean =9.96, $SD = 2.85$). 80% or more of the study participants reported limitation in activities such as walking several blocks, climbing stairs, kneeling or stooping, pushing a vacuum cleaner, and lifting or carrying groceries. Difficulty traveling around the community because of health issues was reported to occur at least some of the time by 27% of study participants and 38% reported spending most days or everyday in the bed or chair because of their health.

The initial model for *functional limitations* included the three symptom clusters, the 3 two-way interaction terms of the symptom clusters, and selected participant characteristics (age, co-morbidity) (See Table 6). Interaction terms were entered into the model to evaluate for cluster independence. Using a backward stepwise elimination model-building approach the model was trimmed of non-significant variables (age, co-morbidity, GI Distress, and the interaction of GI Distress with Discomforts of Illness and Sickness Behavior). This process

resulted in a model incorporating Discomforts of Illness ($\beta = -.765, p = 0.0005$), Sickness Behavior ($\beta = -.681, p = 0.0005$) and the interaction of Sickness Behavior and Discomforts of Illness ($\beta = .649, p = 0.014$). The interaction between these variables implies that not only do functional limitations increase as discomforts of illness increase, but increases at a faster rate when sickness behaviors are increased. This model explained approximately 58% of the variance in functional limitations ($F = 35.96, p = 0.0005, R^2 = 0.578$) (See Table 6).

These same variables were entered into a regression model of factors predicting *mobility*. Variables predictive of mobility include co-morbidity ($\beta = -.214, p = 0.019$) and sickness behavior ($\beta = -.328, p = 0.005$) (See Table 6). A model trimmed of non-significant variables (age, discomforts of illness, GI distress) using the backward stepwise elimination model-building approach, included sickness behavior ($\beta = -0.441, p = 0.0005$) and co-morbidity ($\beta = -0.200, p = 0.019$) (See Table 6). This suggests that as an individual with heart failure experiences increased sickness behavior symptoms, mobility decreases. This model ($F = 20.68, p = 0.0005, R^2 = 0.275$) explained almost 30% of the variance in mobility.

Discussion

This study examined heart failure symptoms, identified symptom clusters, and evaluated the effect of the identified heart failure symptom clusters on functional limitations and mobility. The incidence of heart failure is high in individuals over 65 years of age¹. The mean age of the study sample is 56 years. According to the Centers for Disease Control and Prevention (CDC), the percentage of adults age 45 – 64 years with two or more of nine selected chronic health conditions, which included heart disease and diabetes (a prevalent co-morbidity in our sample) increased during the periods 1999 – 2000 and 2009 – 2012.⁴³ Thus, the younger age of our sample is consistent with the younger age reported by the CDC. The individuals in this sample

had decreased cardiac function as indicated by their markedly reduced EF and over half of them required an ICD. While the majority of the study sample was NYHA class II or III, 13% of the sample was NYHA class IV. Patients who have NYHA class IV heart failure are those who have heart failure symptoms at rest and may be difficult to capture as study participants in the outpatient setting. In our participants, these patients were either listed for transplant or were being evaluated for transplant and/or mechanical circulatory support.

The cluster of symptoms (e.g., anxiety, depression, fatigue, daytime function, and cognitive changes) identified in this study, is similar to the constellation of symptoms known as sickness behavior identified by Dantzer⁴⁴ who describes the construct as a highly organized and purposeful response of an organism to activation of the immune system or to non-immune stimuli. Fatigue, depression and changes in daytime functioning due to sleepiness are features of sickness behavior. Kelly et al.⁴⁵ reported that hypersomnia, listlessness, and loss of interest in social activities are features of sickness behavior.

The sickness behavior symptom cluster identified in this study is consistent with but includes additional symptoms than identified by other investigators who studied heart failure symptom clusters.²²⁻²⁴ Riegel and Ratcliffe²² reported that adults with heart failure who are in NYHA class IV and who present with the group of symptoms known as sickness behaviors have an increased likelihood of being hospitalized or of dying than adults with NYHA class IV heart failure who do not exhibit symptoms associated with sickness behavior. The symptoms that were identified in the sickness behavior cluster included depression, fatigue, and sleepiness. The symptoms in the sickness behavior cluster that we identified empirically using PCA included these three symptoms as well as anxiety and cognitive dysfunction characterized, for example, as forgetfulness, confusion, decreased attention, and concentration.

Few investigators observed symptom clusters which included appetite or hunger. Song, et al.²⁶ isolated a symptom cluster that included lack of appetite with lack of energy and difficulty sleeping. In our study appetite and hunger composed a symptom cluster. Diet is a crucial component of effective heart failure management. The association of appetite and hunger with other symptoms in heart failure symptom clusters will need further investigation.

Our approach to the identification of symptom clusters was selection and measurement of individual symptoms prevalent in individuals with heart failure. From this analysis a discomforts of illness symptom cluster composed of dyspnea, pain, and edema emerged. Some of the symptoms in the discomforts of illness cluster (e.g., pain) may receive limited attention by clinicians during the assessment of individuals with heart failure. However, attention to this cluster may be important to helping an individual with heart failure preserve their lifestyle and engage in enjoyable activities.

Investigators have observed relationships between physical functioning and single symptoms--for example fatigue,^{17,19} sleep disordered breathing,^{46,47} pain,⁴⁸ and depression²⁰ in individuals with heart failure. However, among the studies examining the relationship of heart failure symptom clusters with outcomes, there are no reports about the relationship between heart failure symptom clusters and functional limitations or mobility. Perhaps the recognition of a need for energy conservation and redirection of activities toward recuperation from an exacerbation of heart failure needs to receive more attention in care planning for individuals experiencing a heart failure exacerbation.

A relationship was observed between the heart failure symptom clusters, sickness behaviors and discomforts of illness with functional limitations. In regression analysis, both increased sickness behaviors and increased discomforts of illness predicted limitations in

functional status. While functional limitations increased as sickness behaviors and discomforts of illness increased, functional limitations increased as a faster rate as sickness behaviors increased. This provides evidence to clinicians that when caring for an individual with heart failure it is important to consider a range of symptoms rather than only the classic heart failure symptoms as sources of functional limitations.

When helping an individual with heart failure improve mobility, addressing sickness behaviors, the strongest predictor of mobility in our study, has the potential to help patients improve functional status. A decrease in sickness behaviors and fewer comorbid conditions is associated with improved mobility which may result in the ability to get out into the community, to use public transportation, and to spend less time spent confined to the bed or chair. .

The TUS proved to be an adequate model for the study of heart failure symptom clusters. Three groups of heart failure symptoms that met the definition of a symptom cluster were identified. Although the independence of the symptom clusters sickness behaviors and discomforts of illness with respect to functional limitations may be questionable, we were able to demonstrate that 2 symptom clusters were related to the performance outcome functional limitations and one symptom cluster was related to performance outcome mobility.

Limitations

The study findings reported here provide insight into symptom clusters in patients with heart failure; however, we recognize some limitations. First, the findings may not be generalizable because the sample was a convenience sample of heart failure patients followed at one institution. In addition, the sample was composed of symptomatic outpatients followed in a tertiary care setting and the symptom profile of this sample may not be representative of the symptom profile observed in the acute care setting. Study participants could have magnified or

minimized their reporting of symptoms. However, by definition symptoms are an individual's perceptions which can only be recognized and confirmed by the individual.^{49,50} Instrument selection may have limited the range of symptom characteristics that were considered in developing the symptom clusters identified by this study.

Implications

The relationship of the three symptom clusters to functional limitations and mobility outcomes adds to the evidence validating the composition of heart failure symptom clusters. The findings of this study suggest that while it is important to include symptoms such as dyspnea, fatigue, and peripheral edema in heart failure patient assessment it is equally important to inquire about the presence of symptoms such as pain, daytime function, appetite, and hunger. As patients are instructed to manage their heart failure perhaps health care providers need to remember that participation in self-care activities may be limited because the individual with heart failure may suffer from the combination of anxiety, depression, and fatigue which is compounded by daytime dysfunction which includes sleepiness and decreased energy during the day and cognitive dysfunction which includes difficulty with problem solving, forgetfulness and difficulty concentrating.

Several areas for further research are suggested by the findings of this study. More study is needed to establish the stability of heart failure symptom clusters. The finding of an interaction between sickness behaviors and discomforts of illness associated with functional limitations indicates research to confirm the independence of these symptom clusters is warranted. Examination of the relationship of heart failure symptom clusters with biomarkers of heart failure such as the brain natriuretic peptide (BNP) may aid the identification of underlying mechanisms producing heart failure symptom clusters. Finally, additional exploration of

associations between symptom clusters and outcomes will advance the science of heart failure symptom cluster identification.

Summary and Conclusions

It is early in the evaluation of cluster formation among heart failure symptoms and stability of heart failure symptom clusters across samples and over time has not been established. However, this study lends support for the existence of a cluster of heart failure symptoms similar to those symptoms associated with sickness behaviors as described by Dantzler⁴⁴ and substantiated by Riegel and Ratcliffe²² in individuals with heart failure. The identification of a symptom cluster composed of symptoms associated with discomfort highlights the significance of illness discomforts such as pain, edema and shortness of breath, in living with heart failure. This finding represents a signal that measures to alleviate symptoms such as pain and shortness of breath need to be prominent in nursing care planning for individuals with heart failure. The relationship of appetite and hunger with other heart failure symptoms has not received much attention by investigators and provides an opportunity for additional research.

Assessment and care for individuals with heart failure should focus on managing a cluster of symptoms which may includes some less typical heart failure symptoms rather than single symptoms. Interventions targeting a cluster of symptoms as opposed to a single symptom can provide a more focused approach to decreasing functional limitations and improving mobility so individuals with heart failure can engage in activities such as moving about the community to do grocery shopping, filling prescriptions, walking, or even sports such as golf

References

1. Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics – 2011 update: A report from the American Heart Association. *Circulation*. 2011;123(4): e18 -e209. doi:10.1161/CIR.0b013e3182009701. Accessed August 11, 2011.
2. Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: A report of the American College of Cardiology/American Heart Association task force on practice guidelines. *Circulation*. 2005; 112(12):e154 – e235. doi: 10.1161/CIRCULATIONAHA.105.167586
3. Lenz ER, Pugh LC, Milligan RA, Gift A, Suppe F. The Middle-Range Theory of Unpleasant symptoms: an update. *ANS Adv Nurs Sci*. 1997; 19(3):14 – 27.
<http://ovidsp.tx.ovid.com.proxy.library.vcu.edu/sp-3.5.1a/ovidweb.cgi> Accessed April 11, 2011
4. Wilson IB, Cleary PD. Linking clinical variables with health-related quality of life, a conceptual model of patient outcomes. *JAMA*. 1995; 273(1):59 – 65.
<http://jamanetwork.com.proxy.library.vcu.edu/article.aspx?volume=273&page=59> Accessed April 11,2011.
5. Dodd MJ, Miaskowski C, Paul SM. Symptom clusters and their effect on the functional status of patients with cancer. *Onc Nurs Forum*. 2001;28(3):465 – 470.
<http://web.ebscohost.com.proxy.library.vcu.edu/ehost/pdfviewer/pdfviewer?sid=b271ab0b-faae-430a-9568-99976fc9ab95%40sessionmgr12&vid=2&hid=25>
6. Kim H-J, McGuire DB, Tulman L, Barsevick AM. Symptom clusters, concept analysis and clinical implications for cancer nursing. *Cancer Nurs*. 2005;28(4):270 – 282.

<http://ovidsp.tx.ovid.com.proxy.library.vcu.edu/sp-3.5.1a/ovidweb.cgi?QS2>. Accessed April 21, 2011.

7. Miaskowski C, Dodd M, Lee, K. Symptom clusters: The new frontier in symptom management research. *J Natl Cancer Inst Monogr*. 2004; 2004(32): 17 – 21. doi: 10.1093/jncimonographs/lgh023

8. American Heart Association. 1994 Revisions to classification of functional capacity and objective assessment of patients with diseases of the heart. 2007.

<http://www.americanheart.org/presenter.jhtml?identifier=1712>. Accessed July 16, 2007.

9. Leidy NK. Functional status and the forward progress of merry-go-rounds: toward a coherent analytical framework. *Nurs Res*. 1994; 43(4):196 – 202.

<http://ovidsp.tx.ovid.com.proxy.library.vcu.edu>

10. Wang T-J. Concept analysis of functional status. *Int J Nurs Stud*. 2004; 41(4):457 – 462. doi: 10.1016/j.ijnurstu.2003.09.004

11. Hertzog MA, Pozehl B, Duncan K. Cluster analysis of symptom occurrence to identify subgroups of heart failure patients, a pilot study. *J Cardiovasc Nurs*. 2010; 25(4):273 – 283. doi: 10.1097/JCN.0b013e318cfbb6c

12. Jurgens, C.Y., Moser, D.K., Armola, R., et al. Symptom clusters of heart failure. *Res Nurs Health*. 2009; 32(5): 551 – 560. doi:10.1002/nur.20343

13. Lee KS, Song EK, Lennie TA, et al. Symptom clusters in men and women with heart failure and their impact on cardiac event-free survival. *J Cardiovasc Nurs*. 2010; 25(4): 263 – 272.

14. Jurgens CY, Fain JA, Riegel B. Psychometric testing of the heart failure somatic awareness scale. *J Cardiovasc Nurs*. 2006; 21(2): 95-102.
<http://ovidsp.tx.ovid.com.proxy.library.vcu.edu/sp-3.5.1a/ovidweb.cgi?QS2>. Accessed August 8, 2011.
15. Falk K, Patel H, Swedberg K, Ekman I. Fatigue in patients with chronic heart failure – a burden associated with emotional and symptom distress. *Euro J Cardiovasc Nurs*. 2009; 8(2):91 – 96. doi: 10.1016/j.ejcnurse.2008.07.002
16. Smith ORF, Michielsen HJ, Pelle AJ, Schiffer AA, Winter JB, Denollet J. Symptoms of fatigue in chronic heart failure patients: clinical and psychological predictors. *Eur J Heart Fail*. 2007; 9(9): 922-927. doi:10.1016/j.ejheart.2007.05.016
17. Chen L-H, Li C-Y, Shieh S-M., YinW-H, Chiou A-F. Predictors of fatigue in patients with heart failure. *J Clin Nurs*. 2010;19(11-12):1588-1596. doi:10.1111/j.1365-2010.03218.x
18. Evangelista LS, Moser DK, Westlake C, Pike N, Ter-Galstanyan A, Dracup, K. Correlates of fatigue in patients with heart failure. *Prog Cardiovasc Nurs*. 2008; 23(1):12 – 17.
<http://web.ebscohost.com.proxy.library.vcu.edu/ehost/pdfviewer/pdfviewer?sid=45a802e2-e9c3-4dae-85cd-0be4389c6159%40sessionmgr4&vid=2&hid=25>. Published May 2, 2008. Updated
19. Fink AM, Sullivan SL, Zerwic JJ, Piano MR. Fatigue with systolic heart failure. *J Cardiovasc Nurs*. 2009; 24(5):410 – 417. doi:10.1097/JCN.0b013e3181ae1e84
20. Sullivan M, Levy WC, Russo JE, Spertus JA. Depression and health status in patients with advanced heart failure: A prospective study in tertiary care. *J Card Fail*. 2004; 10(5):390-396. doi: 10.1016/j.cardfail.2004.01.011

21. Tang WR, Yu CY, Yeh SJ. Fatigue and its related factors in patients with chronic heart failure. *J Clin Nurs*. 2010; 19(1-2):69-78. doi:10.1111/j.1365-2702.2009.02959.x
22. Riegel B, Ratcliffe SJ. Abstract 8268: sickness behavior predicts event free survival in adults with heart failure. *Circulation*. 2011;124 (21)(suppl).
http://circ.ahajournals.org/cgi/content/meeting_abstract/124/21_MeetingAbstracts/A8268.
23. Johansson P, Alehagen U, Svensson E, Svanborg E, Dahlström U, Broström A. Determinants of global perceived health in Community-dwelling elderly screened for heart failure and sleep-disordered breathing. *J Cardiovasc Nurs*. 2010; 25(5), E16 – E26.
doi:10.1097/JCN.0b013e3181d6de6f
24. Redeker NS, Jeon S, Muench U, Campbell D, Walseleben J, Papoport DM. (2010). Insomnia symptoms and daytime function in stable heart failure. *Sleep*. 2012; 33(9):1210-1216.
<http://www.ncbi.nlm.nih.gov.proxy.library.vcu.edu/pmc/articles/PMC2938862/pdf/aasm.33.9.1210.pdf>. Accessed April 19, 2011.
25. Smith ORF, Gidron Y, Kupper N, Winter JB, Denollet J. Vital exhaustion in chronic heart failure: symptom profiles and clinical outcome. *J Psychosom Res*. 2009; 66(3):195-201.
doi:10.1016/j.jpsychores.2009.10.021
26. Song EK, Moser DK, Rayens MK, Lennie TA. Symptom clusters predict event-free survival in patients with heart failure. *J Cardiovasc Nurs*. 2010; 25(4): 284-291.
doi:10.1097/JCN.0b013e318cfbcb.

27. Jessup M, Abraham WT, Casey DE, et al. 2009 Focused update: ACCF/AHA guidelines for the diagnosis and management of heart failure in adults. *Circulation*, 2009;119(14): 1977 – 2016. doi: 10.1161/CIRCULATIONAHA.109.192064
28. Charlson ME, Pompei P, Ales, KL, MacKenzie CR. A New Method of Classifying Prognostic Comorbidity in Longitudinal Studies: Development and Validation. *J Chronic Dis*. 1987; 40(5): 373 – 383.
29. Lewin, R.J.P., Thompson, D.R., Martin, C.R., et al. Validation of the cardiovascular limitations and symptoms profile (CLASP) in chronic stable angina. *J Cardiopulm Rehabil*. 2002; 22(3):184 – 191. Retrieved from <http://ovidsp.tx.ovid.com.proxy.library.vcu.edu>
30. Lopez V, Chair SY, Thompson DR, Tsui YY, Lewin RJP. (2008). A psychometric evaluation of the Chinese version of the cardiovascular limitations and symptoms profile in patients with coronary heart disease. *J Clin Nurs*. 2008; 17(17):2327 – 2334. doi: 10.1111/j.1365-2702.2008.02294.x
31. Mendoza TR, Wang X S, Cleeland CS, et al. The rapid assessment of fatigue severity in cancer patients, use of the brief fatigue inventory. *Cancer*. 1999; 85(5): 1186 – 1196. doi: 10.1002/(SICI)1097-0142(19990301)
32. Webel AR, Frazier SK, Moser, DK, Lennie TA. Daily variability in dyspnea, edema and body weight in heart failure patients. *Eur J Cardiovasc Nurs*. 2007; 6(1):60-65. doi: 10.1016/j.ejcnurse.2006.04.003
33. Zigmond AS, Snaith RP. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*. 1983; 67(6): 361 – 370.

34. Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). *Medical Care*. 1992; 30(6):473 – 438. <http://ovidsp.tx.ovid.com.proxy.library.vcu.edu> Accessed June 24, 2012.
35. McHorney CA, Ware JE, Raczek AE. The MOS 36-item short form health survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Medical Care*. 1993; 31(3): 247 – 263. <http://ovidsp.tx.ovid.com.proxy.library.vcu.edu>
36. Lennie TA, Moser DK, Heo S, Chung ML, Zambroski CH. Factors influencing food intake in patients with heart failure, a comparison with healthy elders. *J Cardiovasc Nurs*. 2006; 21(2):123 – 129. <http://ovidsp.tx.ovid.com.proxy.library.vcu.edu>.
37. Lee KA. Self-reported sleep disturbances in employed women. *Sleep*. 1992; 15(6):493 – 498.
38. Reddeker NS, Stein S. Characteristics of sleep in patients with stable heart failure versus a comparison group. *Heart Lung*. 2006; 35(4):252-261. doi: 10.1016/j.hrtlng.2005.10.007
Accessed November 25, 2012.
39. Riegel B, Ratcliffe SJ, Sayers, SL, et al. Determinants of excessive daytime sleepiness and fatigue in adults with heart failure. *Clin Nurs Res*. 2012;21(3):271-293. doi:
10.1177/1054773811419842
40. Stewart, A.L., Ware, J.E., Sherbourne, C.D. & Wells, K.B. Psychological distress/well-being and cognitive functioning measures. In: Stewart AL, Ware JE eds. *Measuring functioning and well-being*. Durham, NC: Duke University Press; 1992: 102 – 142.

41. Hayes RD, Sherbourne CD, Mazel RM. *Users manual for the medical outcomes study (MOS) core measures of health-related quality of life*. Santa Monica, CA: Rand; 1995.
http://www.rand.org/content/dam/rand/pubs/monograph_reports/2008/MR162.pdf
42. Stewart AL, Kamberg CJ. Physical functioning measures. In: Stewart AL, Ware JE eds. *Measuring functioning and well-being*. Durham, NC: Duke University Press; 1992: 86 – 101.
43. Freid VM, Bernstein AB, Bush MA. *Multiple chronic conditions among adults aged 45 and over: trends over the past 10 years*, NCHS Data Brief No. 100. Atlanta, GA: Centers for Disease Control and Prevention; 2012. <http://www.cdc.gov/nchs/data/databrief/db100.pdf>
44. Dantzer, R. Cytokine-induced sickness behavior: mechanisms and implications. *Ann N Y Acad Sci*. 2001; 993(1):222 – 234. doi:10.1111/j.1749-6632.2001.1b05827.x
December 31, 2010. Accessed 8/14/2011.
45. Kelly KW, Bluthé R.-M, Dantzer R, et al. Cytokine-induced sickness behavior. *Brain, Behavior and Immunity*. 2003;17(1)(suppl 15): S112 – S118.
<http://www.sciencedirect.com.proxy.library.vcu.edu/science/article/pii/S0889159102000776>
46. Redeker NS, Muench U, Zucker,MJ, et al. (2010). Sleep disordered breathing, daytime symptoms, and functional performance in stable heart failure. *Sleep*. 2010; 33(4):551 – 560.
<http://www.ncbi.nlm.nih.gov.proxy.library.vcu.edu/pmc/articles/PMC2849795/pdf/aasm.33.4.551.pdf>. Accessed April 13, 2011.
47. Skobel E, Norra C, Sinha A, Breuer C, Hanrath P, Stellbrink C. Impact of sleep-related breathing disorders on health-related quality of life in patient with chronic heart failure. *Eur J Heart Fail*. 2005; 7(4): 505-511. doi:10.1016/j.eheart.2004.07.020.

48. Goebel JR, Doering LV, Shugarman LR, et al. Heart failure: The hidden problem of pain. *J Pain Symptom Manage*. 2009; 38(5):698 – 707. doi: 10.1016/j.jpainsymman. 2009.04.022
49. Fu MR, LeMone P, McDaniel RW. An integrated approach to an analysis of symptom management in patients with cancer. *Oncol Nurs Forum*. 2004; 31(1):65 – 70.
doi: 10.1188/04.ONF.65-70
50. Gapstur RL. Symptom burden: a concept analysis and implications for oncology nurses. *Oncol Nurs Forum*. 2007; 34(3):673 – 680. doi: 10.1188/07.ONF.673-680

What's New?

- Three symptom clusters, sickness behavior, discomforts of illness, and GI distress were identified.
- To reduce functional limitations and improve mobility in individuals with heart failure it is necessary to consider symptoms such as anxiety, depression, pain, daytime dysfunction, and cognitive dysfunction.
- Reducing the level of effect by the symptoms in the Sickness Behavior symptom cluster could improve a heart failure patient's mobility from being bed or chair bound all or most days to only being bed or chair bound some days.

Table 1 Sample Characteristics (N=117)

Variable :	f (%)	Mean (SD)
Age		56 (13)
20 – 39 years	15 (13%)	
40 – 59 years	49(42%)	
60 – 79 years	50 (43%)	
≥ 80 years	1 (1%)	
Gender		
Female	44(38%)	
Male	73 (62%)	
Race		
White	55 (47%)	
African American	59(50%)	
American Indian/Alaska Native	2 (2%)	
Education		
Did not complete high school	20(17%)	
High school graduate	60 (51%)	
College graduate	37 (32%)	
Marital Status		
Never married	25 (21%)	
Previously married	28 (24%)	
Currently married	64 (55%)	
Employment Status		
Unemployed (but looking for work)	8 (7%)	
Employed part time	2 (2%)	
Employed full time	18 (15%)	
Homemaker	1 (1%)	
Receiving disability	49 (42%)	
Retired	36 (31%)	
Student	1 (1%)	
Volunteer	2 (2%)	
Length of time with heart failure		5.2 (5.3)
<1 year	15 (14%)	
1– 5 years	51 (48%)	
6 – 10 years	27 (25%)	
11– 15 years	10 (9%)	
>15 years	3 (3%)	
NYHA Classification		2.7 (0.70)
Class 2	51 (44%)	
Class 3	46 (39%)	
Class 4	15 (13%)	
Ejection Fraction (EF)		25% (10%)

F =frequency *SD* = standard deviation

Table 2: Symptom Prevalence

Symptom	<i>f</i> (%)	Mean (<i>SD</i>)
Shortness of breath at least once or twice in the past 2 weeks	117 (100%)	3 days (1.8)
Decreased appetite (9 or less on a visual analogue scale with 0 = no appetite)	99(85%)	6.5 (2.7)
Decreased hunger (9 or less on a visual analogue scale with 0 = no hunger sensation)	110 (95%)	5 (2.6)
Depression	25 (21%)	7.7 (4.3)
Anxiety	38 (32.5%)	8.3 (4.6)
Daytime functioning: Felt sleepy during the day 4 or more days during the week	78(67.8%)	4.5 days (2.4)
Fatigue: Felt unusually tired or fatigued in the past week	85 (73%)	0.84 (.89)
Fatigue: Usual level of fatigue during the past week		4.8 (2.8)
Edema (1 or greater on a visual analogue scale with 0 = no edema and 10 = worst edema I have ever had)	84 (73%)	3.9(3.6)
Pain (1=yes; 0=no)	73 (62%)	3.2 (1.6)
Cognitive function: Difficulty reasoning and solving problems	49 (42%)	4.7(1.4)
Cognitive function: Difficulty with concentration and thinking at least some of the time	51(43%)	4.5(1.4)

F =frequency *SD* = standard deviation

Table 3 Symptom Factor Loadings

Symptom	Component 1 (Sickness Behavior)	Component 2 (GI Disturbance)	Component 3 (Discomforts of Illness)
Shortness of Breath			.758
Edema			.824
Pain			.737
Anxiety	.832		
Depression	.822		
Daytime Function	.741		
Cognitive Function	.761		
Fatigue	.757		
Appetite		.829	
Hunger		.819	

Table 4: Model Variables

Variable	Mean	Standard Deviation	Minimum Score	Maximum Score
Sickness Behaviors				
• Anxiety	8.28	4.64	0	19
• Depression	7.67	4.28	0	21
• Fatigue	30.04	17.59	0	60
• Daytime Function	17.75	7.29	3	32
• Cognitive Dysfunction	27.85	6.81	6	36
Discomforts of Illness				
• Shortness of Breath	9.23	3.61	1	15
• Edema	3.91	3.63	0	10
• Pain	8.08	6.98	0	20
GI Distress				
• Appetite	6.46	2.74	1	10
• Hunger	5.01	2.66	0	10
Participant Characteristics				
• Age	56	12.95	20	84
• Co-morbidity	3.21	1.93	1	11
Outcomes				
• Functional Limitations	16.10	5.09	10	29
• Mobility	9.96	2.85	3	13

Table 5: Descriptive Statistics of Symptom Clusters

Symptom Cluster	Mean	Standard Deviation	Minimum	Maximum
Sickness Behaviors	58.89	25.16	11.27	110.50
Discomforts of Illness	16.03	8.69	0.76	33.59
GI Disturbance	9.45	3.75	0.83	16.48


Table 6: Regression Analyses of Functional Limitations and Mobility

Model:	β	Standard Error	Beta	Significance
Functional Limitations ($F = 17.68, p = 0.0005, R^2 = 0.591$)				
• Age	0.014	0.029	0.034	0.631
• Co-morbidity	-0.359	0.188	-0.135	0.059
• Discomforts of Illness	-0.410	0.164	-0.700	0.014
• GI Distress	0.192	0.229	0.142	0.404
• Sickness Behavior	-0.139	0.053	-0.687	0.010
• Sickness Behavior* Illness Discomforts	0.004	0.002	0.645	0.018
• Sickness Behavior * GI Distress	0.001	0.005	0.071	0.820
• Illness Discomforts* GI Distress	-0.006	0.013	-0.120	0.662
Trimmed Model of Functional Limitations ($F = 35.96, p = 0.0005, R^2 = 0.578$)				
• Discomforts of Illness	-0.456	0.106	-0.765	0.0005
• Sickness Behavior	-0.139	0.031	-0.681	0.0005
• Sickness Behavior*Illness Discomforts	0.004	0.002	0.649	0.014
Mobility ($F = 8.96, p = 0.0005, R^2 = 0.305$)				
• Age	0.027	0.027	0.089	0.325
• Co-morbidity	-0.412	0.173	-0.214	0.019
• Sickness Behavior	-0.048	0.017	-0.328	0.005
• Discomforts of Illness	-0.064	0.048	-0.151	0.188
• GI Distress	0.013	0.083	0.013	0.880
Trimmed Model of Mobility ($F = 20.68, p = 0.0005, R^2 = 0.275$)				
• Co-morbidity	-0.388	0.163	-0.200	0.019
• Sickness Behavior	-0.065	0.012	-0.441	0.0005

 B = unstandardized beta coefficient

Beta – standardized beta coefficient

Figure 1: Conceptual Framework*



*Adapted from the Theory of Unpleasant Symptoms (Lenz, Pugh, Milligan & Gift, 1997)

Vita

Janet Kay Herr was born June 13, 1951, in Alton, Illinois and is an American citizen. She graduated from Huguenot High School, Richmond, Virginia in 1969. She received a Bachelor of Science in Nursing from the University of Virginia in 1973. She has been employed as a Registered Nurse at Chippenham Hospital in Richmond, Virginia since graduation from nursing school. Her current position is as a clinical data specialist in quality management. She received a Master of Science in Nursing from Virginia Commonwealth University in 1993 and a Post-master's Certificate in Aging Studies from Virginia Commonwealth University in 2007.