



VCU

Virginia Commonwealth University
VCU Scholars Compass

Theses and Dissertations

Graduate School

2022

Black Hearts Matter: A Comparative Study of Sociodemographic, Clinical, and Psychosocial Characteristics in Black and White Persons With Heart Failure

Hannah M Anderson Hughes
Virginia Commonwealth University

Follow this and additional works at: <https://scholarscompass.vcu.edu/etd>



Part of the [Biological Psychology Commons](#), [Health Psychology Commons](#), [Multicultural Psychology Commons](#), [Other Nursing Commons](#), [Psychiatric and Mental Health Nursing Commons](#), [Public Health and Community Nursing Commons](#), [Quality Improvement Commons](#), [Social Justice Commons](#), and the [Somatic Psychology Commons](#)

© The Author

Downloaded from

<https://scholarscompass.vcu.edu/etd/7190>

This Dissertation is brought to you for free and open access by the Graduate School at VCU Scholars Compass. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Black Hearts Matter: A Comparative Study of Sociodemographic, Clinical, and Psychosocial
Characteristics in Black and White Persons With Heart Failure

A dissertation submitted in partial fulfillment of the requirements for the degree of Doctor of
Philosophy at Virginia Commonwealth University

by

Hannah Marie Anderson Hughes

Associate degree in Nursing, Norfolk State University, 2004

Bachelor of Science in Nursing, Norfolk State University, 2005

Master of Science in Nursing Education, Walden University, 2009

Post-Master's Certificate in Nursing Informatics, Walden University, 2012

Master of Business Administration, Jack Welch Management Institute, 2020

Chair: Jo Lynne W. Robins, PhD, RN, ANP-BC, AHN-C, FAANP

Associate Professor, School of Nursing

November 10, 2022

ABSTRACT

Heart failure (HF) is a chronic, progressive, debilitating illness that disproportionately affects Black individuals, often leading to poor quality of life (QOL). Various sociodemographic, clinical, and psychological characteristics are related to QOL in patients with HF. Because most studies do not include representative Black samples or report findings from racial subgroup analyses, these relationships and any existing racial differences are not well understood. The purpose of this descriptive, comparative study was to explore the relationships among sociodemographic, clinical, and psychological characteristics and QOL in patients with HF and to determine if there were racial differences between Black and White individuals. Mean participant age was 56.29 ($SD = 12.42$). SPSS 27.0 was the program used to conduct secondary data analysis. Among the sociodemographic characteristics, racial differences were apparent in age, gender, and marital status. Racial differences in ejection fraction, New York Heart Association Classification, comorbidities, and functional status emerged as clinical characteristics. Race was not significantly associated with QOL. Physical functioning and anxiety were significant predictors of QOL in the initial analysis. However, in the full model, older age and greater depression remained significant, suggesting they were the strongest predictors of QOL. Patients with HF should undergo regular screening for depression and anxiety. Future researchers should explore sociodemographic, clinical, and psychological characteristics in larger samples; consider the timing and measurement approaches for depression, anxiety, and QOL; and further explore racial differences.

ACKNOWLEDGMENTS

I would first like to thank God, my Heavenly Father, who gave me the ability to pursue a terminal degree within my beloved profession of nursing. He has never left me while I traveled this winding, uphill, journey to completion. Thank you, God!

I would like to thank Alminis Hughes for being my most consistent support throughout this journey. I could not complete the PhD program without his support and contributions. Alminis, thank you for supporting my goals, even at your inconvenience. Thank you for your many sacrifices. Thank you for encouraging me to persist, despite so many obstacles. Thank you for the meals prepared, the work done, and for your prayers that covered me. Thank you for making sure that I never had to worry about Myles, and for all of the support that you offered in many ways. I will never be able to repay you, but please accept my heartfelt gratitude and love.

I would like to thank my son, Myles, who is 9 years old. He was born on Day 6 of my doctoral journey. He has sacrificed much time with me and has been a significant motivator in me completing this journey. Son, thank you for believing in your “Doctor Mom!” Thank you for encouraging me! Thank you for asking when I would *finally* be finished and be able to wear that huge robe. I hope you are inspired to pursue excellence and I hope my displays of resilience, tenacity, and commitment to my goals and higher education inspire you along your life’s journey. I love you! You are forever my favorite son!

As a first-generation college graduate, I must recognize my father, the late James Anderson. Although my father’s formal education concluded in elementary school, he was a strong proponent of education and paid for my 13 years of private school education. Unfortunately, he passed away when I was 18 years old and a freshman in college, so he did not

get to witness any of my higher education or professional accomplishments. He would be so proud! I honor him!

I also thank my mother who has always been a strong proponent of education. My mother made so many sacrifices for her children over the years. She came to help care for my son as I began my doctoral journey, which was essential for my being able to focus on doctoral coursework with a newborn. My mother developed heart failure secondary to chemotherapy treatment for breast cancer while I was studying heart failure. Although painful, she became a part of my “why” for this research in one sense. Thank you, Mommy, for your many sacrifices over the years, investing into me, and always encouraging us to pursue our highest potential. This PhD is also for you!

I thank Ms. Floris Williams, who supported me early in my journey. She relocated to help me take care of my son, which gave me the time to focus on my doctoral coursework and milestones. She cooked and helped take care of the household, and I did not have to worry about much. I am grateful for her investments in making this goal my reality. Thank you!

I want to recognize my siblings: Conrad and Deborah Maynes, Lincoln and Jenelle Maynes, Jermaine and Abigail Young, and George and Elizabeth Slusher. I am forever grateful for your love, prayers, and support in thought, word, and deed. Thank you to Dr. Lincoln Maynes, the first doctorally prepared of my mother’s offspring. With your focus, determination, and hard work, you inspired us regarding what’s possible! You set your goals and pursued them relentlessly. You set a strong example for us to follow! Thank you!

I also recognize the Hughes family for all that you have been to me and done for me and us over the years. Thank you for being supportive of us and for your many prayers.

I would like to thank the Virginia Commonwealth School of Nursing for its unwavering belief in and support of me and for giving me the opportunity to complete the PhD in Nursing, despite numerous challenges and setbacks. Your care extends beyond academic rigor, and I will always be grateful that I chose VCU SON and that VCU SON repeatedly chose me.

I would like to thank my dissertation committee members, Dr. Jo Lynne Robins (Chair), Dr. Rebecca Etz, Dr. Jeanne Salyer, and Dr. Debra Barksdale, for their support and for guiding me toward the completion of this degree and milestone in my life. Your unwavering support, commitment, and many contributions are so appreciated. I am forever grateful to you for investing into me and leading me to completion. I would also like to thank Ms. Fonda Neal, whose support of the doctoral programs and students at VCU has been invaluable. I also thank Dr. R. K. Elswick for his statistical consultation support.

I offer heartfelt gratitude to my mentors, Dr. Janice Brewington, Dr. Nicole Morris, Dr. Michael Plater, Dr. Jacqueline Palmer, Dr. Carline Jean-Gilles, Dr. Deborah Jones, and Dr. Kimberly Adams-Tufts. Your guidance, unwavering support, faith in my abilities, “truth-telling,” and consistent presence have made a world of difference. Thank you!

I extend sincere appreciation to the faculty and staff of Norfolk State University Department of Nursing for laying the foundation of my nursing career. You prepared me well to grow and develop my career while remaining true to the ideals of the profession of nursing. My nursing professors included Dr. W. F. White-Parson, Dr. Sheila Cannon, Dr. Rosalyn Gardiner, Dr. Candace Rogers, Ms. Karen Yancey, and the late Berthenia Cromwell.

I would like to thank Ms. Dina Eagle, my friend and confidante. You have walked with me through so many seasons of life in these recent years. You have been my sounding board, my voice of reason, my harsh critic, my “other lens” from which to look at a situation, and even my

“antagonizer.” I appreciate you so much! Thank you, Dina! To my best friend of over 25 years, Ms. Melissa Pinney. Thank you for always being there. Your quiet, yet supportive nature is always felt. You are the epitome of persistence, determination, tenacity, and focus on your goals. Thank you for being the example. Thank you for being my true and consistent friend. Thank you to Tiffany Eversley for her friendship and most of all her prayers. I really appreciate your genuine love and friendship!

I appreciate the Black Docs Group so much! You have been “community” for me. I appreciate the friendships, encouragement, and support cultivated among Black doctoral pursuers and completers. I will forever be indebted to this community of scholars.

I am grateful to so many of my family members and friends who have supported me in various ways along the way, from being a listening ear, offering a wonderful meal, or providing a word of encouragement. I also thank so many other nursing and personal friends and colleagues who have been so supportive to me along my journey. I appreciate each of you so much!

Finally, to my community where it all started: Thank you for pouring into me and grooming me to pursue my highest potential. I am forever indebted to my home, St. Croix, U.S. Virgin Islands, and the many people who helped to build, develop, and groom “Hannah.” This accomplishment is for America’s Paradise, too!

DEDICATION

First, I dedicate this dissertation to my mother, Charmaine Anderson, a retired licensed practical nurse who aspired to be a registered nurse. My mother sacrificed greatly for my four siblings and me, and I honor her investments and sacrifice as I become Dr. Hannah.

Second, I dedicate this dissertation to my father, the late James Hogan Anderson, who passed away when I was 18 years old and a freshman in college. Although he invested so much into me, passionately believed in my abilities, and was so proud of my accomplishments, he did not live long enough to see me graduate from college, far less earn a PhD. Da, they will call me Dr. Hannah now! You have a doctor! Although you only had an elementary school education yourself, you firmly believed in education for your children and invested into it. Thank you for all that you did to lay the educational foundation for the day I would defend my dissertation. You would have been so proud!

I also dedicate this dissertation to my dear sister friend, Rosalyn Ann Lee (June 15, 1985 to January 14, 2021). Rosalyn embodied the spirit of nursing! She was one of the most caring, compassionate, strong, determined, resilient, and kind people I have ever met. A graduate of practical nursing education and an aspiring registered nurse herself, she found joy in meticulously caring for the sick and assisting the less fortunate in her home care business and her volunteer work in the community. Rosalyn extended this love and care to her family and friends and everyone she encountered. Although her life's journey was short due to COVID-19-related illness and an underlying diagnosis of multiple sclerosis, she made a great impact on so many. I was looking forward to so many more accomplishments from her—so, Ros, we'll share this one. Ros, you are forever in my heart and I'll love you forever. Thoughts of your measure of strength, tenacity, and resilience helped me to finish this degree, even after you left us so soon and while I

was writing this dissertation. Amid my sorrow and heartbreak over losing you, I found the will to finish anyway, just as you would have done. I pushed past the pain, hurt, grief, shock, and disappointment, just as you did every time life gave you enough reasons to cry. Instead, you always chose to smile! We will always miss your smile! Thank you, Sister! Your legacy continues to inspire and push me. Even in death, you helped me finish!

I also dedicate this terminal degree to the place where I received my foundation in nursing: my HBCU, Norfolk State University! I am Spartan proud and will forever be grateful for the many investments made into me by my nursing professors, such as Dr. W. F. White-Parson, Dr. Rosalyn Gardiner, Dr. Candace Rogers, Ms. Karen Yancey, Ms. Berthenia Cromwell, and Dr. Sheila Cannon. Thank you for introducing me to the profession I love, for investing into me, and for leading by example.

As I reflect on my life, I also dedicate this dissertation to my home, St. Croix, U.S. Virgin Islands. I was born and raised on an island—an American territory of just 84 square miles—yet, I always knew I had limitless potential. Thank you for investing in me. Thank you for believing in me. Thank you for your support, nurture, and love. Thank you for the friendships and the partnerships that extended to St. Thomas, St. John, and the British Virgin Islands. You laid the foundation upon which everything else was built! I would also like to thank two local organizations from my hometown: the Virgin Islands Department of Education, for awarding me a scholarship to pursue undergraduate nursing education, and HOVENSA, LLC, which provided a 4-year scholarship that helped to fund my undergraduate education. Your investment would go on to yield for years to come!

TABLE OF CONTENTS

ABSTRACT.....	ii
ACKNOWLEDGMENTS	iii
DEDICATION.....	vii
TABLE OF CONTENTS.....	ix
CHAPTER 1: INTRODUCTION AND STATEMENT OF PROBLEM	1
Chronic Illness Challenges, Chronic Illness With Blacks, Disparities, Depression and Anxiety, Sociodemographic and Clinical Characteristics, and Quality of Life.....	1
Heart Failure Overview and Statistics	3
Heart Failure and Racial Disparities Among Blacks	4
Summary, Research Focus, and Research Implications	6
Organization of the Dissertation	8
References.....	10
CHAPTER 2: LITERATURE REVIEW AND THEORETICAL FRAMEWORK.....	31
Conceptual Model.....	31
Conceptual Model for the Current Study.....	34
Literature Review.....	42
Sociodemographic and Clinical Characteristics	42
Clinical Characteristics	57
Psychological Characteristics	67
Summary/Conclusion.....	85
References.....	87
CHAPTER 3: METHODOLOGY	136

Introduction.....	136
Rationale for Conducting a Secondary Data Analysis.....	137
Setting, Sample, Data Collection, and Human Subjects Protection	137
Study Variables and Measures.....	138
Sociodemographic Characteristics.....	138
Clinical Characteristics	139
Psychological Characteristics	141
Quality of Life.....	142
Statistical Analysis.....	144
Aim 1	145
Aim 2	145
Limitations and Delimitations.....	146
References.....	148
CHAPTER 4: RESULTS.....	161
Introduction.....	161
Data Analysis	161
Approach and Phases of Data Analysis	161
Data Preparation and Management.....	162
Creation of New Variables.....	162
Reliability.....	163
Statistical Power Analysis.....	163
Descriptive Analysis	163
Sociodemographic Characteristics.....	164

Clinical Characteristics	165
Quality of Life.....	166
Analysis for Aim 3.....	172
Bivariate Analysis.....	Error! Bookmark not defined.
Sociodemographic Characteristics.....	172
Clinical Characteristics	172
Psychological Characteristics	172
Multivariate Analysis.....	174
References.....	178
CHAPTER 5: DISCUSSION.....	179
Introduction.....	179
Discussion.....	180
Aim 1: Racial Differences and Associations Between Sociodemographic, Clinical, and Psychological Characteristics	180
Aim 2: Sociodemographic, Clinical, and Psychological Predictors of Quality of Life.....	190
Limitations	196
Conclusion	197
Recommendations and Future Research.....	198
Alignment With National Institute for Nursing Research	219
Advancing Health Disparities in Heart Failure Research	219
References.....	221

CHAPTER 1: INTRODUCTION AND STATEMENT OF PROBLEM

Chronic Illness Challenges, Chronic Illness With Blacks, Disparities, Depression and Anxiety, Sociodemographic and Clinical Characteristics, and Quality of Life

Chronic illnesses are prevalent (Boersma et al., 2020; NCCDPHP, 2021a; Raghupathi & Raghupathi, 2018) and costly (Buttorff et al., 2017; Martin et al., 2021; NCCDPHP, 2021b), lead to functional status impairment and disability (van Campen & Cardol, 2009), and require patients to make many life adjustments to effectively manage and cope with the associated physiological and psychosocial changes (Bury, 1991; Charmaz, 1995; Chin et al., 2000; Loeb, 2006; Schulman-Green et al., 2016; Sharpe & Curran, 2006; Sidell, 1997). Many individuals live with multiple chronic illnesses (Boersma et al., 2020; Denton & Spencer, 2010; Vogeli et al., 2007), contributing to even more pronounced physical limitations, psychosocial challenges, and disability (Iezzoni, 2010). Black individuals are disproportionately affected by chronic illnesses (Calvin et al., 2003; Ezeokpo & Ugwueze, 2022; Kanchi et al., 2018; Piccolo et al., 2013; Quinones et al., 2019; Smalls et al., 2020; Smedley et al., 2002) with higher morbidity, comorbidity (Marrie et al., 2008; McGee et al., 1996), disability (Thorpe et al., 2016; Whitson et al., 2011), and mortality rates (Howard et al., 2016; Smalls et al., 2020; Smedley et al., 2002) than Whites.

Living with chronic illness is challenging for patients because of the physical symptoms and psychosocial issues (Ingram & South, 2020; Schulman-Green et al., 2016), the complex treatment plan requirements, and the functional status limitations they often experience (Charmaz, 1995; Schulman-Green et al., 2016; Sidell, 1997). Among the psychosocial challenges faced by chronically ill patients are independently occurring depression and anxiety (Brown et al., 2010; Di Benedetto et al., 2014; Eisner et al., 2010; Fattouh et al., 2019; Li et al., 2008;

Thomas et al., 2003), and cooccurring depression and anxiety, with more pronounced adverse effects (DeJean et al., 2013; Liew, 2012; von Leupoldt et al., 2011; Yohannes et al., 2010).

Various sociodemographic and clinical characteristics influence depression and anxiety in chronically ill patients (Ani et al., 2009; Bell et al., 2005; El-Gabalawy et al., 2011; Rutledge et al., 2006; Thomas et al., 2003). Among these sociodemographic and clinical characteristics is race, which is a key variable that might influence depression and anxiety in chronically ill Blacks; however, its potential influence is not well understood (Assari, 2014; Assari et al., 2015; Hill-Joseph, 2019; Johnson-Lawrence et al., 2013; Li et al., 2008; Mouzon, 2017). Some investigators have found that Blacks with chronic illnesses have more depression (Assari, 2014; Assari et al., 2015; Mouzon, 2017), whereas others suggest that Blacks might have less depression than Whites (Ani et al., 2009; Hollingshaus & Utz, 2013) or found no racial differences in depression (Ghods et al., 2008) and anxiety (Ellis et al., 2015). However, it is generally known that Blacks do not receive the same rates of screening, diagnosis, and treatment for depression as White individuals (Barnes & Bates, 2017; Lagomasino et al., 2011; Stockdale et al., 2008). Depression underdiagnosis in Blacks might be due to its presentation differing from that of White individuals (Bailey et al., 2019). Black individuals experience more fluctuations in depressive symptoms, which suggests that the measurement timing might confound screening attempts and study findings in this racial minority group (Hill-Joseph, 2019).

Anxiety is also associated with chronic illnesses in Black people (Assari, 2016; Watkins et al., 2015). Although some researchers have found Black individuals are less likely to experience anxiety than Whites, anxiety in Blacks is associated with greater negative effects, including more significant functional impairment (Himle et al., 2009). Yet, it remains unclear

whether Black individuals living with chronic illnesses experience anxiety at similar rates as their White counterparts or whether race influences rates of anxiety in chronically ill Blacks.

Depression and anxiety in chronically ill patients, particularly when undiagnosed and untreated, contribute to decreased life satisfaction, poor health-related quality of life (HRQOL) (Balcells et al., 2010; Brown et al., 2010; Chuquilín-Arista et al., 2021; Lebel et al., 2020; Liu et al., 2020; Sadlonova et al., 2021; Strine et al., 2008; von Leupoldt et al., 2011), and negative clinical outcomes (Assari, 2014; McLaughlin et al., 2005; Ryu et al., 2016; Sidell, 1997; Whittemore & Dixon, 2008), including greater morbidity, healthcare utilization (Ani et al., 2009), and mortality (McLaughlin et al., 2005; Sadlonova et al., 2021). Chronically ill Blacks have the worst functional status and quality of life (QOL; Jackson-Triche et al., 2000). More effort is necessary to reduce and eliminate persistent racial disparities among Black, chronically ill patients (Smedley et al., 2002). Research focused on understanding potential racial differences could provide a better understanding of whether depression and anxiety contribute to the persistent disparities.

Heart Failure Overview and Statistics

Heart failure (HF) is one of the most prevalent, costly, and debilitating chronic illnesses (AHA, 2017; Centers for Disease Control and Prevention, 2020) that contributes to morbidity, healthcare utilization, and mortality (Sullivan et al., 2002; Virani, Alonso, Aparicio, et al., 2020; Virani, Alonso, Benjamin, et al., 2021). HF is challenging for patients because of its chronicity, multibody system involvement, progressive nature, association with disability, and negative impact on psychosocial wellness (Riegel et al., 2009; Wong et al., 2011). HF affects 6.2 million American adults 20 years and older, with approximately 960,000 new diagnoses annually. As of 2017, HF accounted for about 80,480 deaths yearly (Virani, Alonso, Aparicio, et al., 2020;

Virani, Alonso, Benjamin, et al., 2021). Patients with HF have significant healthcare utilization resulting from frequent exacerbations that require medical intervention (Urbich et al., 2020). Annually, patients with HF have 1,932,000 primary care visits, 414,000 emergency department visits, and 809,000 HF-related hospital readmissions (Virani et al., 2020). HF is the leading cause of rehospitalization for patients ages 65 and older, accounting for 24% of all-cause 30-day rehospitalizations (Desai & Stevenson, 2012; Heidenreich et al., 2011). HF management places a significant burden on the U.S. healthcare system, with annual costs of approximately \$30.7 billion (Desai & Stevenson, 2012; Sullivan et al., 2002; Virani et al., 2020). Projections show that by 2030, HF-related costs will increase to \$160 billion, which reflects a 46% increase; Heidenreich et al., 2013; Tsao et al., 2022).

Heart Failure and Racial Disparities Among Blacks

Similar to other chronic illnesses, HF disproportionately affects Black individuals (Benjamin et al., 2017; Go et al., 2013; Roger et al., 2012; Tsao et al., 2022; Virani, Alonso, Aparicio, et al., 2020; Virani, Alonso, Benjamin, et al., 2021), and several disparities exist among Black individuals with HF. Black people have the highest risk of developing HF (Virani et al., 2020), are a majority of the HF patient population (Heidenreich et al., 2011), tend to be younger (Vaccarino et al., 2002), and have a higher incidence before age 50 (Benjamin et al., 2017; Bibbins-Domingo et al., 2009; Go et al., 2013). Blacks are also more likely to have comorbidities and greater mortality than their White counterparts (Evangelista, Dracup, & Doering, 2002; Mensah et al., 2005; Mentz et al., 2013; Roger et al., 2012; Vaccarino et al., 2002). Other disparities include: Blacks are more likely to delay seeking HF-related treatment (Cuyjet & Akinboboye, 2014; Virani et al., 2020), to receive emergency department care and require readmissions (Alhurani et al., 2015; Cuyjet & Akinboboye, 2014; Evangelista, Dracup, &

Doering, 2002), have the highest rates of HF-related hospitalizations (Virani et al., 2020), have greater functional decline following hospitalization (Vaccarino et al., 2002), and have higher mortality rates than Whites (Mensah et al., 2005; Mentz et al., 2013; Roger et al., 2012; Virani et al., 2020). According to projections, HF will substantially negatively impact Blacks more than Whites over the next several years. Interventions to decrease racial disparities are needed (Van Nuys et al., 2018).

Because depression and anxiety are common in patients with HF (Easton et al., 2016; Ishak et al., 2020; Johansson et al., 2006; Joynt et al., 2004; Konstam et al., 2005; MacMahon & Lip, 2002; Norra et al., 2008; Rustad et al., 2013; Rutledge et al., 2006b; Sokoreli et al., 2016; Yohannes et al., 2010), undiagnosed depression and anxiety might be a contributing factor to the disparate clinical outcomes observed among Blacks, such as higher rates of mortality and rehospitalization (Virani, Alonso, Aparicio, et al., 2020; Virani, Alonso, Benjamin, et al., 2021). Similar to the effects of depression and anxiety on QOL in other chronic illness populations (Lim et al., 2011), depression and anxiety negatively impact QOL in patients with HF (Ahmed et al., 2006; Bekelman et al., 2007; Garin et al., 2009; Rumsfeld et al., 2003; Rustad et al., 2013; Samartzis et al., 2013; Trivedi et al., 2009). Improving depression and anxiety might improve QOL in Black patients with HF (Blumenthal, 2012); however, the influence of depression and anxiety on Black individuals with HF is not well understood because Blacks are underrepresented in the published literature and race has not been investigated for its potential influence on QOL.

To date, research on the prevalence and influence of depression and anxiety in Black patients with HF and contributing factors, such as sociodemographic and clinical characteristics, and outcomes related to depression and anxiety, such as QOL, are rarely stratified by race.

Among the limited studies conducted with Black individuals with HF, findings vary. Factors that might contribute to disparate outcomes among Black individuals with HF, such as depression and anxiety, require further exploration to determine whether racial differences exist.

Summary, Research Focus, and Research Implications

Despite advancements in understanding psychological and psychosocial functioning in patients with HF, research focused on depression, anxiety, and their comorbid existence remains limited, and any differences that might exist between Blacks and Whites are not well understood (Dickson et al., 2013; Evangelista, Ter-Galstanyan, et al., 2009). Even when systematic reviews and meta-analyses are conducted using studies of patients with HF, it is difficult to compare and draw conclusions from findings related to depression and anxiety because included studies are heterogenous and consider different covariates (Celano et al., 2018; Sokoreli et al., 2016).

Although Black individuals comprise the majority racial group within the HF patient population (Virani, Alonso, Aparicio, et al., 2020; Virani, Alonso, Benjamin, et al., 2021), they are underrepresented in most studies of depression and anxiety (Gaffey et al., 2022). Generally, investigators did not consider race as an important construct in their study design and only reported racial groups descriptively, indicating there were no baseline differences; grouped all minorities in their analysis; did not include representative Black samples; and did not conduct racial subgroup analyses, even when recruiting representative samples of Black patients. For this reason, the body of research focused on the factors that influence depression and anxiety in Blacks is even more limited and lacks a cultural lens (Dickson et al., 2013). It is unknown whether existing findings can be generalized to Blacks (Bean et al., 2009). Yet, how race influences health, chronic illness, and healthcare should be a variable of interest when conducting research (Bhopal et al., 2021; Culley, 2006). This knowledge gap could contribute to

the persistent negative and disparate clinical outcomes among Black patients with HF. A better understanding of depression and anxiety in Black individuals with HF, the sociodemographic and clinical factors that could influence any relationships, and how depression and anxiety might impact QOL differently in Blacks compared to Whites is needed.

Additional research is needed to explore the incidence and impact of comorbid depression and anxiety in Black patients with HF compared to their White counterparts. Understanding the relationships among and influences on sociodemographic and clinical variables on depression and anxiety in Black individuals with HF is essential to inform evidence that guides patient care. Such evidence could ultimately improve morbidity, mortality, care quality, reduces rehospitalizations and care costs (Dickson et al., 2013) and enhance overall clinical outcomes (MacMahon & Lip, 2002) for all patients with HF, but especially for those who self-identify as Black and who continue to experience disparate clinical outcomes (Evangelista, Ter-Galstanyan, et al., 2009).

The literature suggests that sociodemographic characteristics and clinical characteristics influence depression and anxiety in patients with HF and that depression and anxiety negatively influence QOL in patients with HF. Researchers have not stratified their findings by race or examined race as a sociodemographic variable of potential influence. As a result, there is a poor understanding of the potential impact of race on depression and anxiety in patients with HF. Also unclear is whether racial differences are sociodemographic and clinical characteristics and the impact of depression and anxiety on QOL. There is also relatively no literature focused on anxiety in Black individuals with HF or comorbid depression and anxiety in Black individuals with HF. Additional research is needed to better understand any racial differences between Black and White people, which might help decrease disparities and improve clinical outcomes among

Black individuals with HF. The purpose of this descriptive, comparative study was to explore the relationships among sociodemographic, clinical, and psychological characteristics and QOL in patients with HF and to determine if there were racial differences between Blacks and Whites. The research question was: *What are the differences between Blacks and Whites with HF with respect to sociodemographic and clinical characteristics and quality of life?* The specific aims were to (a) to describe the differences and examine the relationships between sociodemographic, clinical, and psychological characteristics and QOL in Black and White persons with HF; and (b) examine the influence of sociodemographic, clinical, and psychological characteristics on QOL in Black and White persons with HF.

Findings from this study will provide foundational new knowledge to help better understand how psychosocial factors might impact Black individuals with HF and inform interventions to improve HF care and decrease disparate clinical outcomes for Black patients with HF. This study aligns with the National Institute for Nursing Research's (NINR; 2016) future strategic direction related to examining social determinants of health and their impact on health outcomes and QOL, health equity (reducing and eliminating systemic and structural inequities among disadvantaged groups), and tertiary prevention approaches for reducing disease severity, symptoms, and progression, with an emphasis on eliminating health disparities (NINR, 2022).

Organization of the Dissertation

Chapter 2 will include a discussion of the conceptual framework that guided the study design. There will be a review of the existing literature regarding depression and anxiety within the context of chronic illness, specifically HF; sociodemographic and clinical influencers of depression and anxiety; and the impact of depression and anxiety on QOL in patients with HF.

Chapter 3 presents the methodology used to conduct the study, and Chapter 4 includes the findings. Chapter 5 is a discussion of the findings, study weaknesses/limitations, and future directions to further advance this body of knowledge.

References

- Ahmed, A., Ali, M., Lefante, C. M., Mullick, M. S. I., & Kinney, F. C. (2006). Geriatric heart failure, depression, and nursing home admission: An observational study using propensity score analysis. *The American Journal of Geriatric Psychiatry, 14*(10), 867–875. <https://doi.org/10.1097/01.JGP.0000209639.30899.72>
- Alhurani, A. S., Dekker, R. L., Abed, M. A., Khalil, A., Al Zaghal, M. H., Lee, K. S., Mudd-Martin, G., Biddle, M. J., Lennie, T. A., & Moser, D. K. (2015). The association of comorbid symptoms of depression and anxiety with all-cause mortality and cardiac rehospitalization in patients with heart failure. *Psychosomatics, 56*(4), 371–380. <https://doi.org/10.1016/j.psych.2014.05.022>
- American Heart Association. (2017). *What is heart failure?* Retrieved February 26, 2021, from <https://www.heart.org/en/health-topics/heart-failure/what-is-heart-failure>
- Ani, C., Bazargan, M., Hindman, D., Bell, D., Rodriguez, M., & Baker, R. S. (2009). Comorbid chronic illness and the diagnosis and treatment of depression in safety net primary care settings. *Journal of the American Board of Family Medicine, 22*(2), 123–135. <https://doi.org/10.3122/jabfm.2009.02.080035>
- Assari, S. (2014). Chronic medical conditions and major depressive disorder: Differential role of positive religious coping among African Americans, Caribbean Blacks and Non-Hispanic Whites. *International Journal of Preventive Medicine, 5*(4), 405–413.
- Assari, S. (2016). Race and ethnic differences in additive and multiplicative effects of depression and anxiety on cardiovascular risk. *International Journal of Preventive Medicine, 2016*(7), 22–22. <https://doi.org/10.4103/2008-7802.173931>

- Assari, S., Burgard, S., & Zivin, K. (2015). Long-term reciprocal associations between depressive symptoms and number of chronic medical conditions: Longitudinal support for Black–White health paradox. *Journal of Racial and Ethnic Health Disparities*, 2(4), 589–597. <https://doi.org/10.1007/s40615-015-0116-9>
- Bailey, R. K., Mokonogho, J., & Kumar, A. (2019). Racial and ethnic differences in depression: Current perspectives. *Neuropsychiatric Disease and Treatment*, 15, 603–609. <https://doi.org/10.2147/NDT.S128584>
- Balcells, E., Gea, J., Ferrer, J., Serra, I., Orozco-Levi, M., de Batlle, J., Rodriguez, E., Benet, M., DonaireGonzalez, D., Anto, J. M., & Garcia-Aymerich, J. (2010). Factors affecting the relationship between psychological status and quality of life in COPD patients. *Health and Quality of Life Outcomes*, 8(1), 108–108. <https://doi.org/10.1186/1477-7525-8-108>
- Barnes, D. M., & Bates, L. M. (2017). Do racial patterns in psychological distress shed light on the Black–White depression paradox? A systematic review. *Social Psychiatry and Psychiatric Epidemiology*, 52(8), 913–928. <https://doi.org/10.1007/s00127-017-1394-9>
- Bean, M. K., Gibson, D., Flattery, M., Duncan, A., & Hess, M. (2009). Psychosocial factors, quality of life, and psychological distress: Ethnic differences in patients with heart failure. *Progress in Cardiovascular Nursing*, 24(4), 131–140. <https://doi.org/10.1111/j.1751-7117.2009.00051.x>
- Bekelman, D. B., Havranek, E. P., Becker, D. M., Kutner, J. S., Peterson, P. N., Wittstein, I. S., Gottlieb, S. H., Yamashita, T. E., Fairclough, D. L., & Dy, S. M. (2007). Symptoms, depression, and quality of life in patients with heart failure. *Journal of Cardiac Failure*, 13(8), 643–648. <https://doi.org/10.1016/j.cardfail.2007.05.005>

Bell, R. A., Smith, S. L., Arcury, T. A., Snively, B. M., Stafford, J. M., & Quandt, S. A. (2005).

Prevalence and correlates of depressive symptoms among rural older African Americans, Native Americans, and Whites with diabetes. *Diabetes Care*, 28(4), 823–829.

<https://doi.org/10.2337/diacare.28.4.823>

Benjamin, J. E., Blaha, J. M., Chiuve, E. S., Cushman, R. M., Das, D. S., Deo, R. R., De

Ferranti, C. S., Floyd, C. J., Fornage, E. M., Gillespie, H. C., Isasi, T. C., Jiménez, H. M., Jordan, E. L., Judd, W. S., Lackland, K. D., Lichtman, R. J., Lisabeth, J. L., Liu, J. S.,

Longenecker, A. C., ... Muntner, S. P. (2017). Heart disease and stroke statistics—2017 update: A report from the American Heart Association. *Circulation*, 135(10), e146–e603.

<https://doi.org/10.1161/CIR.0000000000000485>

Bhopal, R., Gruer, L., Agyemang, C., Davidovitch, N., de-Graft Aikins, A., Krasnik, A.,

Martinez-Donate, A. P., Miranda, J. J., Pottie, K., Segal, U., Zwi, A., & Kumar, B.

(2021). The Global Society on Migration, Ethnicity, Race and Health: Why race can't be ignored even if it causes discomfort. *European Journal of Public Health*, 31(1), 3–4.

<https://doi.org/10.1093/eurpub/ckaa191>

Bibbins-Domingo, K., Pletcher, M. J., Lin, F., Vittinghoff, E., Gardin, J. M., Arynchyn, A.,

Lewis, C. E., Williams, O. D., & Hulley, S. B. (2009). Racial differences in incident heart failure among young adults. *New England Journal of Medicine*, 360(12), 1179–1190.

<https://doi.org/10.1056/NEJMoa0807265>

Blumenthal, J. A. (2012). Effects of exercise training on depressive symptoms in patients with

chronic heart failure: The HF-ACTION randomized trial. *JAMA*, 308(17), 1742–1742.

<https://doi.org/10.1001/jama.2012.13837>

- Boersma, P., Black, L. I., & Ward, B. W. (2020). Prevalence of multiple chronic conditions among US adults, 2018. *Preventing Chronic Disease*, 2020(17), Article 200130. <https://doi.org/10.5888/pcd17.200130>
- Brown, L. F., Kroenke, K., Theobald, D. E., Wu, J., & Tu, W. (2010). The association of depression and anxiety with health-related quality of life in cancer patients with depression and/or pain. *Psycho-Oncology*, 19(7), 734–741. <https://doi.org/10.1002/pon.1627>
- Bury, M. (1991). The sociology of chronic illness: A review of research and prospects. *Sociology of Health & Illness*, 13(4), 451–468. <https://doi.org/10.1111/j.1467-9566.1991.tb00522.x>
- Buttorff, C., Ruder, T., & Bauman, M. (2017). *Multiple chronic conditions in the United States*. RAND Corporation. https://www.rand.org/content/dam/rand/pubs/tools/TL200/TL221/RAND_TL221.pdf
- Calvin, R., Winters, K., Wyatt, S. B., Williams, D. R., Henderson, F. C., & Walker, E. R. (2003). Racism and cardiovascular disease in African Americans. *The American Journal of the Medical Sciences*, 325(6), 315–331. <https://doi.org/10.1097/00000441-200306000-00003>
- Celano, C. M., Villegas, A. C., Albanese, A. M., Gaggin, H. K., & Huffman, J. C. (2018). Depression and anxiety in heart failure: A review. *Harvard Review of Psychiatry*, 26(4), 175–184. <https://doi.org/10.1097/HRP.0000000000000162>
- Centers for Disease Control and Prevention. (2020). *Heart failure*. Retrieved February 26, 2022, from https://www.cdc.gov/heartdisease/heart_failure.htm
- Charmaz, K. (1995). The body, identity, and self: Adapting to impairment. *Sociological Quarterly*, 36(4), 657–680. <https://doi.org/10.1111/j.1533-8525.1995.tb00459.x>

- Chin, M. H., Polonsky, T. S., Thomas, V. D., & Nerney, M. P. (2000). Developing a conceptual framework for understanding illness and attitudes in older, urban African Americans with diabetes. *The Diabetes Educator*, 26(3), 439–449.
<https://doi.org/10.1177/014572170002600311>
- Chuquilín-Arista, F., Álvarez-Avellón, T., & Menéndez-González, M. (2021). Impact of depression and anxiety on dimensions of health-related quality of life in subjects with Parkinson's disease enrolled in an association of patients. *Brain Sciences*, 11(6), Article 771. <https://doi.org/10.3390/brainsci11060771>
- Culley, L. (2006). Transcending transculturalism? Race, ethnicity and health-care. *Nursing Inquiry*, 13(2), 144–153. <https://doi.org/10.1111/j.1440-1800.2006.00311.x>
- Cuyjet, A. B., & Akinboboye, O. (2014). Acute heart failure in the African American patient. *Journal of Cardiac Failure*, 20(7), 533–540.
<https://doi.org/http://dx.doi.org/10.1016/j.cardfail.2014.04.018>
- DeJean, D., Giacomini, M., Vanstone, M., & Brundisini, F. (2013). Patient experiences of depression and anxiety with chronic disease: A systematic review and qualitative meta-synthesis. *Ontario Health Technology Assessment Series*, 13(16), 1–33.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3817854/>
- Denton, F. T., & Spencer, B. G. (2010). Chronic health conditions: Changing prevalence in an aging population and some implications for the delivery of health care services. *Canadian Journal on Aging*, 29(1), 11–12. <https://doi.org/10.1017/S0714980809990390>
- Desai, S. A., & Stevenson, W. L. (2012). Rehospitalization for heart failure: Predict or prevent? *Circulation*, 126(4), 501–506. <https://doi.org/10.1161/CIRCULATIONAHA.112.125435>

Di Benedetto, M., Lindner, H., Aucote, H., Churcher, J., McKenzie, S., Croning, N., & Jenkins, E. (2014). Co-morbid depression and chronic illness related to coping and physical and mental health status. *Psychology, Health & Medicine, 19*(3), 253–262.

<https://doi.org/10.1080/13548506.2013.803135>

Dickson, V. V., McCarthy, M. M., & Katz, S. M. (2013). How do depressive symptoms influence self-care among an ethnic minority population with heart failure? *Ethnicity & Disease, 23*(1), 22–28.

Easton, K., Coventry, P., Lovell, K., Carter, L.-A., & Deaton, C. (2016). Prevalence and measurement of anxiety in samples of patients with heart failure: Meta-analysis. *Journal of Cardiovascular Nursing, 31*(4), 367–379.

<https://doi.org/10.1097/JCN.0000000000000265>

Eisner, M. D., Blanc, P. D., Yelin, E. H., Katz, P. P., Sanchez, G., Iribarren, C., & Omachi, T. A. (2010). Influence of anxiety on health outcomes in COPD. *Thorax, 65*(3), 229–234.

<https://doi.org/10.1136/thx.2009.126201>

El-Gabalawy, R. M. A., Mackenzie, C. S., Shoostari, S., & Sareen, J. (2011). Comorbid physical health conditions and anxiety disorders: A population-based exploration of prevalence and health outcomes among older adults. *General Hospital Psychiatry, 33*(6), 556–564.

<https://doi.org/10.1016/j.genhosppsy.2011.07.005>

Ellis, E. M., Orom, H., Giovino, G. A., & Kiviniemi, M. T. (2015). Relations between negative affect and health behaviors by race/ethnicity: Differential effects for symptoms of depression and anxiety. *Health Psychology, 34*(9), 966–969.

<https://doi.org/10.1037/hea0000197>

- Evangelista, L. S., Dracup, K., & Doering, L. V. (2002). Racial differences in treatment-seeking delays among heart failure patients. *Journal of Cardiac Failure*, 8(6), 381–386.
<https://doi.org/10.1054/jcaf.2002.129234>
- Evangelista, L. S., Ter-Galstanyan, A. M., Moughrabi, S. R., & Moser, D. K. (2009). Anxiety and depression in ethnic minorities with chronic heart failure. *Journal of Cardiac Failure*, 15(7), 572–579. <https://doi.org/10.1016/j.cardfail.2009.03.005>
- Ezeokpo, B. C., & Ugwueze, C. V. (2022). The racial and ethnic differences in type 2 diabetes mellitus: A Black race narrative review. *Journal of Diabetology*, 13(1), 25–32.
https://doi.org/10.4103/jod.jod_127_21
- Fattouh, N., Hallit, S., Salameh, P., Choueiry, G., Kazour, F., & Hallit, R. (2019). Prevalence and factors affecting the level of depression, anxiety, and stress in hospitalized patients with a chronic disease. *Perspectives in Psychiatric Care*, 55(4), 592–599.
<https://doi.org/10.1111/ppc.12369>
- Gaffey, A. E., Cavanagh, C. E., Rosman, L., Wang, K., Deng, Y., Sims, M., O'Brien, E. C., Chamberlain, A. M., Mentz, R. J., Glover, L. M., & Burg, M. M. (2022). Depressive symptoms and incident heart failure in the Jackson Heart Study: Differential risk among Black men and women. *Journal of the American Heart Association*, 11(5), Article e022514. <https://doi.org/10.1161/JAHA.121.022514>
- Garin, O., Ferrer, M., Pont, À., Rué, M., Kotzeva, A., Wiklund, I., Van Ganse, E., & Alonso, J. (2009). Disease-specific health-related quality of life questionnaires for heart failure: A systematic review with meta-analyses. *Quality of Life Research*, 18(1), 71–85.
<https://doi.org/10.1007/s11136-008-9416-4>

Ghods, B. K., Roter, D. L., Ford, D. E., Larson, S., Arbelaez, J. J., & Cooper, L. A. (2008).

Patient–physician communication in the primary care visits of African Americans and Whites with depression. *Journal of General Internal Medicine*, 23(5), 600–606.

<https://doi.org/10.1007/s11606-008-0539-7>

Go, A. S., Mozaffarian, D., Roger, V. L., Benjamin, E. J., Berry, J. D., Blaha, M. J., Dai, S.,

Ford, E. S., Fox, C. S., Franco, S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Huffman, M. D., Judd, S. E., Kissela, B. M., Kittner, S. J., ... Turner, M. B. (2013). Heart disease and stroke statistics—2014 update: A report from the American Heart Association. *Circulation*, 129(3), e28–e292.

<https://doi.org/10.1161/01.cir.0000441139.02102.80>

Heidenreich, P. A., Albert, N. M., Allen, L. A., Bluemke, D. A., Butler, J., Fonarow, G. C.,

Ikonomidis, J. S., Khavjou, O., Konstam, M. A., Maddox, T. M., Nichol, G., Pham, M., Piña, I. L., & Trogon, J. G. (2013). Forecasting the impact of heart failure in the United States: A policy statement from the American Heart Association. *Circulation: Heart Failure*, 6(3), 606–629. <https://doi.org/10.1161/HHF.0b013e318291329a>

Heidenreich, P. A., Trogon, J. G., Khavjou, O. A., Butler, J., Dracup, K., Ezekowitz, M. D.,

Finkelstein, E. A., Hong, Y., Johnston, S. C., Khera, A., Lloyd-Jones, D. M., Nelson, S. A., Nichol, G., Orenstein, D., Wilson, P. W., & Woo, Y. J. (2011). Forecasting the future of cardiovascular disease in the United States: A policy statement from the American Heart Association. *Circulation*, 123(8), 933–944.

<https://doi.org/10.1161/CIR.0b013e31820a55f5>

- Hill-Joseph, E. A. (2019). Coping while Black: Chronic illness, mastery, and the Black-White health paradox. *Journal of Racial and Ethnic Health Disparities*, 6(5), 935–943.
<https://doi.org/10.1007/s40615-019-00594-9>
- Himle, J. A., Baser, R. E., Taylor, R. J., Campbell, R. D., & Jackson, J. S. (2009). Anxiety disorders among African Americans, Blacks of Caribbean descent, and non-Hispanic Whites in the United States. *Journal of Anxiety Disorders*, 23(5), 578–590.
<https://doi.org/10.1016/j.janxdis.2009.01.002>
- Hollingshaus, M. S., & Utz, R. L. (2013). Depressive symptoms following the diagnosis of major chronic illness. *Society and Mental Health*, 3(1), 22–39.
<https://doi.org/10.1177/2156869312464788>
- Howard, G., Moy, C. S., Howard, V. J., McClure, L. A., Kleindorfer, D. O., Kissela, B. M., Judd, S. E., Unverzagt, F. W., Soliman, E. Z., Safford, M. M., Cushman, M., Flaherty, M. L., & Wadley, V. G. (2016). Where to focus efforts to reduce the Black–White Disparity in stroke mortality: Incidence versus case fatality? *Stroke*, 47(7), 1893–1898.
<https://doi.org/10.1161/STROKEAHA.115.012631>
- Iezzoni, L. I. (2010). Multiple chronic conditions and disabilities: Implications for health services research and data demands. *Health Services Research*, 45(5p2), 1523–1540.
<https://doi.org/10.1111/j.1475-6773.2010.01145.x>
- Ingram, S. H., & South, S. C. (2020). Dimensional latent factor structure of psychopathology in adults with chronic illness. *Psychological Assessment*, 32(12), 1145–1157.
<https://doi.org/10.1037/pas0000954>

- Ishak, W. W., Edwards, G., Herrera, N., Lin, T., Hren, K., Peterson, M., Ngor, A., Liu, A., Kimchi, A., & Spiegel, B. (2020). Depression in heart failure: A systematic review. *Innovations in Clinical Neuroscience, 17*(4–6), 27–38.
- Jackson-Triche, M. E., Greer Sullivan, J., Wells, K. B., Rogers, W., Camp, P., & Mazel, R. (2000). Depression and health-related quality of life in ethnic minorities seeking care in general medical settings. *Journal of Affective Disorders, 58*(2), 89–97.
[https://doi.org/10.1016/S0165-0327\(99\)00069-5](https://doi.org/10.1016/S0165-0327(99)00069-5)
- Johansson, P., Dahlstrom, U., & Brostrom, A. (2006). Consequences and predictors of depression in patients with chronic heart failure: Implications for nursing care and future research. *Progress in Cardiovasc Nursing, 21*(4), 202–211.
<https://doi.org/10.1111/j.0889-7204.2006.05415.x>
- Johnson-Lawrence, V., Griffith, D. M., & Watkins, D. C. (2013). The effects of race, ethnicity, and mood/anxiety disorders on the chronic physical health conditions of men from a national sample. *American Journal of Men's Health, 7*(4_suppl), 58S–67S.
<https://doi.org/10.1177/1557988313484960>
- Joynt, K. E., Whellan, D. J., & O'Connor, C. M. (2004). Why is depression bad for the failing heart? A review of the mechanistic relationship between depression and heart failure. *Journal of Cardiac Failure, 10*(3), 258–271.
<https://doi.org/10.1016/j.cardfail.2003.09.008>

- Kanchi, R., Perlman, S. E., Chernov, C., Wu, W., Tabaei, B. P., Trinh-Shevrin, C., Islam, N., Seixas, A., Rodriguez-Lopez, J., & Thorpe, L. E. (2018). Gender and race disparities in cardiovascular disease risk factors among New York City adults: New York City Health and Nutrition Examination Survey (NYC HANES) 2013–2014. *Journal of Urban Health*, 95(6), 801–812. <https://doi.org/10.1007/s11524-018-0287-x>
- Konstam, V., Moser, D. K., & De Jong, M. J. (2005). Depression and anxiety in heart failure. *Journal of Cardiac Failure*, 11(6), 455–463. <https://doi.org/10.1016/j.cardfail.2005.03.006>
- Lagomasino, I. T., Stockdale, S. E., & Miranda, J. (2011). Racial-ethnic composition of provider practices and disparities in treatment of depression and anxiety, 2003–2007. *Psychiatric Services*, 62(9), 1019–1025. https://doi.org/10.1176/ps.62.9.pss6209_1019
- Lebel, S., Mutsaers, B., Tomei, C., Leclair, C. S., Jones, G., Petricone-Westwood, D., Rutkowski, N., Ta, V., Trudel, G., Laflamme, S. Z., Lavigne, A.-A., & Dinkel, A. (2020). Health anxiety and illness-related fears across diverse chronic illnesses: A systematic review on conceptualization, measurement, prevalence, course, and correlates. *PLoS ONE*, 15(7), Article e0234124. <https://doi.org/10.1371/journal.pone.0234124>
- Li, C., Barker, L., Ford, E. S., Zhang, X., Strine, T. W., & Mokdad, A. H. (2008). Diabetes and anxiety in US adults: Findings from the 2006 Behavioral Risk Factor Surveillance System. *Diabetic Medicine*, 25(7), 878–881. <https://doi.org/10.1111/j.1464-5491.2008.02477.x>
- Liew, H.-P. (2012). Depression and chronic illness: A test of competing hypotheses. *Journal of Health Psychology*, 17(1), 100–109. <https://doi.org/10.1177/1359105311409788>

- Lim, L., Jin, A.-Z., & Ng, T.-P. (2011). Anxiety and depression, chronic physical conditions, and quality of life in an urban population sample study. *Social Psychiatry and Psychiatric Epidemiology*, 47(7), 1047–1053. <https://doi.org/10.1007/s00127-011-0420-6>
- Liu, X., Haagsma, J., Sijbrands, E., Buijks, H., Boogaard, L., Mackenbach, J. P., Erasmus, V., & Polinder, S. (2020). Anxiety and depression in diabetes care: Longitudinal associations with health-related quality of life. *Scientific Reports*, 10(1), Article 8307. <https://doi.org/10.1038/s41598-020-57647-x>
- Loeb, S. J. (2006). African American older adults coping with chronic health conditions. *Journal of Transcultural Nursing*, 17(2), 139–147. <https://doi.org/10.1177/1043659605285415>
- MacMahon, K. M., & Lip, G. Y. (2002). Psychological factors in heart failure: A review of the literature. *Archives of Internal Medicine*, 162(5), 509–516. <https://doi.org/10.1001/archinte.162.5.509>
- Marrie, R. A., Horwitz, R., Cutter, G., Tyry, T., Campagnolo, D., & Vollmer, T. (2008). Comorbidity, socioeconomic status and multiple sclerosis. *Multiple Sclerosis*, 14(8), 1091–1098. <https://doi.org/10.1177/1352458508092263>
- Martin, A. B., Hartman, M., Lassman, D., & Catlin, A. (2021). National health care spending in 2019: Steady growth for the fourth consecutive year. *Health Affairs*, 40(1), 14–24. <https://doi.org/10.1377/hlthaff.2020.02022>
- McGee, D., Cooper, R., Liao, Y., & Durazo-Arvizu, R. (1996). Patterns of comorbidity and mortality risk in Blacks and Whites. *Annals of Epidemiology*, 6(5), 381–385. [https://doi.org/10.1016/S1047-2797\(96\)00058-0](https://doi.org/10.1016/S1047-2797(96)00058-0)

- McLaughlin, T. J., Aupont, O., Bambauer, K. Z., Stone, P., Mullan, M. G., Colagiovanni, J., Polishuk, E., Johnstone, M., & Locke, S. E. (2005). Improving psychologic adjustment to chronic illness in cardiac patients: The role of depression and anxiety. *Journal of General Internal Medicine*, *20*(12), 1084–1090. <https://doi.org/10.1111/j.1525-1497.2005.00256.x>
- Mensah, G. A., Mokdad, A. H., Ford, E. S., Greenlund, K. J., & Croft, J. B. (2005). State of disparities in cardiovascular health in the United States. *Circulation*, *111*(10), 1233–1241. <https://doi.org/10.1161/01.CIR.0000158136.76824.04>
- Mentz, R. J., Bittner, V., Schulte, P. J., Fleg, J. L., Pina, I. L., Keteyian, S. J., Moe, G., Nigam, A., Swank, A. M., Onwuanyi, A. E., Fitz-Gerald, M., Kao, A., Ellis, S. J., Kraus, W. E., Whellan, D. J., & O'Connor, C. M. (2013). Race, exercise training, and outcomes in chronic heart failure: Findings from Heart Failure – A Controlled Trial Investigating Outcomes in Exercise Training (HF-ACTION). *American Heart Journal*, *166*(3), 488–495. <https://doi.org/10.1016/j.ahj.2013.06.002>
- Mouzon, D. M. (2017). Religious involvement and the Black–White paradox in mental health. *Race and Social Problems*, *9*(1), 63–78. <https://doi.org/10.1007/s12552-017-9198-9>
- National Center for Chronic Disease Prevention and Health Promotion. (2021a). *About chronic diseases*. Retrieved October 7, 2021, from <https://www.cdc.gov/chronicdisease/about/index.htm>
- National Center for Chronic Disease Prevention and Health Promotion. (2021b). *Health and economic costs of chronic diseases*. Retrieved October 7, 2021, from <https://www.cdc.gov/chronicdisease/about/costs/index.htm>

National Institute of Nursing Research. (2016, September). *The NINR strategic plan: Advancing science, improving lives*.

https://www.ninr.nih.gov/sites/files/docs/NINR_StratPlan2016_reduced.pdf

National Institute of Nursing Research. (2022). *NINR Strategic Plan (2022–2026)* – Under development. Retrieved April 9, 2022, from <https://www.ninr.nih.gov/aboutninr/ninr-mission-and-strategic-plan>

Norra, C., Skobel, E. C., Arndt, M., & Schauerte, P. (2008). High impact of depression in heart failure: Early diagnosis and treatment options. *International Journal of Cardiology*, *125*(2), 220–231. <https://doi.org/10.1016/j.ijcard.2007.05.020>

Piccolo, R. S., Yang, M., Bliwise, D. L., Yaggi, H. K., & Araujo, A. B. (2013). Racial and socioeconomic disparities in sleep and chronic disease: Results of a longitudinal investigation. *Ethnicity & Disease*, *23*(4), 499–507.

Quinones, A. R., Botosaneanu, A., Markwardt, S., Nagel, C. L., Newsom, J. T., Dorr, D. A., & Allore, H. G. (2019). Racial/ethnic differences in multimorbidity development and chronic disease accumulation for middle-aged adults. *PLoS ONE*, *14*(6), Article e0218462. <https://doi.org/10.1371/journal.pone.0218462>

Raghupathi, W., & Raghupathi, V. (2018). An empirical study of chronic diseases in the United States: A visual analytics approach to public health. *International Journal of Environmental Research and Public Health*, *15*(3), Article 431.

<https://doi.org/10.3390/ijerph15030431>

Riegel, B., Moser, D. K., Anker, S. D., Appel, L. J., Dunbar, S. B., Grady, K. L., Gurvitz, M. Z., Havranek, E. P., Lee, C. S., & Lindenfeld, J. (2009). State of the science promoting self-care in persons with heart failure: A scientific statement from the American Heart Association. *Circulation*, *120*(12), 1141–1163.

<https://doi.org/10.1161/CIRCULATIONAHA.109.192628>

Roger, V. L., Go, A. S., Lloyd-Jones, D. M., Benjamin, E. J., Berry, J. D., Borden, W. B., Bravata, D. M., Dai, S., Ford, E. S., Fox, C. S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Kissela, B. M., Kittner, S. J., Lackland, D. T., Lichtman, J. H., ... Turner, M. B. (2012). Heart disease and stroke statistics—2012 update: A report from the American Heart Association. *Circulation*, *125*(1), e2–e220.

<https://doi.org/10.1161/CIR.0b013e31823ac046>

Rumsfeld, J. S., Havranek, E., Masoudi, F. A., Peterson, E. D., Jones, P., Tooley, J. F., Krumholz, H. M., Spertus, J. A., & Consortium, C. O. R. (2003). Depressive symptoms are the strongest predictors of short-term declines in health status in patients with heart failure. *Journal of the American College of Cardiology*, *42*(10), 1811–1817.

<https://doi.org/10.1016/j.jacc.2003.07.013>

Rustad, J. K., Stern, T. A., Hebert, K. A., & Musselman, D. L. (2013). Diagnosis and treatment of depression in patients with congestive heart failure: A review of the literature. *The Primary Care Companion for CNS Disorders*, *15*(4).

<https://doi.org/10.4088/PCC.13r01511>

- Rutledge, T., Reis, V. A., Linke, S. E., Greenberg, B. H., & Mills, P. J. (2006a). Depression in heart failure: A meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. *Journal of the American College of Cardiology*, 48(8), 1527–1537. <https://doi.org/10.1016/j.jacc.2006.06.055>
- Ryu, E., Chamberlain, A. M., Pendegraft, R. S., Petterson, T. M., Bobo, W. V., & Pathak, J. (2016). Quantifying the impact of chronic conditions on a diagnosis of major depressive disorder in adults: A cohort study using linked electronic medical records. *BMC Psychiatry*, 16, Article 114. <https://doi.org/10.1186/s12888-016-0821-x>
- Sadlonova, M., Wasser, K., Nagel, J., Weber-Krüger, M., Gröschel, S., Uphaus, T., Liman, J., Hamann, G. F., Kermer, P., Gröschel, K., Herrmann-Lingen, C., & Wachter, R. (2021). Health-related quality of life, anxiety and depression up to 12 months post-stroke: Influence of sex, age, stroke severity and atrial fibrillation – A longitudinal subanalysis of the Find-AF_{RANDOMISED} trial. *Journal of Psychosomatic Research*, 142, Article 110353. <https://doi.org/10.1016/j.jpsychores.2020.110353>
- Samartzis, L., Dimopoulos, S., Tziogourou, M., & Nanas, S. (2013). Effect of psychosocial interventions on quality of life in patients with chronic heart failure: A meta-analysis of randomized controlled trials. *Journal of Cardiac Failure*, 19(2), 125–134. <https://doi.org/10.1016/j.cardfail.2012.12.004>
- Schulman-Green, D., Jaser, S. S., Park, C., & Whittemore, R. (2016). A metasynthesis of factors affecting self-management of chronic illness. *Journal of Advanced Nursing*, 72(7), 1469–1489. <https://doi.org/10.1111/jan.12902>
- Sharpe, L., & Curran, L. (2006). Understanding the process of adjustment to illness. *Social Science & Medicine*, 62(5), 1153–1166. <https://doi.org/10.1016/j.socscimed.2005.07.010>

- Sidell, N. L. (1997). Adult adjustment to chronic illness: A review of the literature. *Health & Social Work, 22*(1), 5–11. <https://doi.org/10.1093/hsw/22.1.5>
- Smalls, B. L., Ritchwood, T. D., Bishu, K. G., & Egede, L. E. (2020). Racial/ethnic differences in glycemic control in older adults with type 2 diabetes: United States 2003–2014. *International Journal of Environmental Research and Public Health, 17*(3), Article 950. <https://doi.org/10.3390/ijerph17030950>
- Smedley, B. D., Stith, A. Y., & Nelson, A. R. (Eds.). (2002). *Unequal treatment: Confronting racial and ethnic disparities in health care*. National Academies Press. <https://doi.org/10.17226/12875>
- Sokoreli, I., Vries, J., Pauws, S., & Steyerberg, E. (2016). Depression and anxiety as predictors of mortality among heart failure patients: Systematic review and meta-analysis. *Heart Failure Review, 21*(1), 49–63. <https://doi.org/10.1007/s10741-015-9517-4>
- Stockdale, S. E., Lagomasino, I. T., Siddique, J., McGuire, T., & Miranda, J. (2008). Racial and ethnic disparities in detection and treatment of depression and anxiety among psychiatric and primary health care visits, 1995–2005. *Medical Care, 46*(7), 668–677. <https://doi.org/10.1097/MLR.0b013e3181789496>
- Strine, T. W., Chapman, D. P., Balluz, L. S., Moriarty, D. G., & Mokdad, A. H. (2008). The associations between life satisfaction and health-related quality of life, chronic illness, and health behaviors among U.S. community-dwelling adults. *Journal of Community Health, 33*(1), 40–50. <https://doi.org/10.1007/s10900-007-9066-4>
- Sullivan, M., Simon, G., Spertus, J., & Russo, J. (2002). Depression-related costs in heart failure care. *Archives of Internal Medicine, 162*(16), 1860–1866. <https://doi.org/10.1001/archinte.162.16.1860>

- Thomas, J., Jones, G., Scarinci, I., & Brantley, P. (2003). A descriptive and comparative study of the prevalence of depressive and anxiety disorders in low-income adults with type 2 diabetes and other chronic illnesses. *Diabetes Care*, 26(8), 2311–2317.
<https://doi.org/10.2337/diacare.26.8.2311>
- Thorpe, R. J., Jr., Wynn, A. J., Walker, J. L., Smolen, J. R., Cary, M. P., Szanton, S. L., & Whitfield, K. E. (2016). Relationship between chronic conditions and disability in African American men and women. *Journal of the National Medical Association*, 108(1), 90–98. <https://doi.org/10.1016/j.jnma.2015.12.012>
- Trivedi, R. B., Blumenthal, J. A., O'Connor, C., Adams, K., Hinderliter, A., Dupree, C., Johnson, K., & Sherwood, A. (2009). Coping styles in heart failure patients with depressive symptoms. *Journal of Psychosomatic Research*, 67(4), 339–346.
<https://doi.org/10.1016/j.jpsychores.2009.05.014>
- Tsao, C. W., Aday, A. W., Almarzooq, Z. I., Alonso, A., Beaton, A. Z., Bittencourt, M. S., Boehme, A. K., Buxton, A. E., Carson, A. P., CommodoreMensah, Y., Elkind, M. S. V., Evenson, K. R., Eze-Nliam, C., Ferguson, J. F., Generoso, G., Ho, J. E., Kalani, R., Khan, S. S., Kissela, B. M., ... Martin, S. S. (2022). Heart disease and stroke statistics – 2022 update: A report from the American Heart Association. *Circulation*, 145(8), e153–e639. <https://doi.org/10.1161/CIR.0000000000001052>
- Urbich, M., Globe, G., Pantiri, K., Heisen, M., Bennison, C., Wirtz, H. S., & Di Tanna, G. L. (2020). A systematic review of medical costs associated with heart failure in the USA (2014–2020). *Pharmacoeconomics*, 38(11), 1219–1236. <https://doi.org/10.1007/s40273-020-00952-0>

- Vaccarino, V., Gahbauer, E., Kasl, S. V., Charpentier, P. A., Acampora, D., & Krumholz, H. M. (2002). Differences between African Americans and Whites in the outcome of heart failure: Evidence for a greater functional decline in African Americans. *American Heart Journal*, 143(6), 1058–1067. <https://doi.org/10.1067/mhj.2002.122123>
- van Campen, C., & Cardol, M. (2009). When work and satisfaction with life do not go hand in hand: Health barriers and personal resources in the participation of people with chronic physical disabilities. *Social Science & Medicine*, 69(1), 56–60. <https://doi.org/10.1016/j.socscimed.2009.04.014>
- Van Nuys, K. E., Xie, Z., Tysinger, B., Hlatky, M. A., & Goldman, D. P. (2018). Innovation in heart failure treatment: Life expectancy, disability, and health disparities. *JACC: Heart Failure*, 6(5), 401–409. <https://doi.org/10.1016/j.jchf.2017.12.006>
- Virani, S. S., Alonso, A., Aparicio, H. J., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Cheng, S., Delling, F. N., Elkind, M. S. V., Evenson, K. R., Ferguson, J. F., Gupta, D. K., Khan, S. S., Kissela, B. M., Knutson, K. L., Lee, C. D., Lewis, T. T., ... Tsao, C. W. (2021). Heart disease and stroke statistics – 2021 update: A report from the American Heart Association. *Circulation*, 143(8), e254–e743. <https://doi.org/10.1161/CIR.0000000000000950>
- Virani, S. S., Alonso, A., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Chang, A. R., Cheng, S., Delling, F. N., Djousse, L., Elkind, M. S. V., Ferguson, J. F., Fornage, M., Khan, S. S., Kissela, B. M., Knutson, K. L., Kwan, T. W., Lackland, D. T., ... Tsao, C. W. (2020). Heart disease and stroke statistics – 2020 update: A report from the American Heart Association. *Circulation*, 141(9), e139–e151. <https://doi.org/10.1161/CIR.0000000000000757>

- Vogeli, C., Shields, A. E., Lee, T. A., Gibson, T. B., Marder, W. D., Weiss, K. B., & Blumenthal, D. (2007). Multiple chronic conditions: Prevalence, health consequences, and implications for quality, care management, and costs. *Journal of General Internal Medicine*, 22(S3), 391–395. <https://doi.org/10.1007/s11606-007-0322-1>
- von Leupoldt, A. P., Taube, K. M. D., Lehmann, K., Fritzsche, A. P., & Magnussen, H. M. D. (2011). The impact of anxiety and depression on outcomes of pulmonary rehabilitation in patients with COPD. *Chest*, 140(3), 730–736. <https://doi.org/10.1378/chest.10-2917>
- Watkins, D. C., Assari, S., & Johnson-Lawrence, V. (2015). Race and ethnic group differences in comorbid major depressive disorder, generalized anxiety disorder, and chronic medical conditions. *Journal of Racial and Ethnic Health Disparities*, 2(3), 385–394. <https://doi.org/10.1007/s40615-015-0085-z>
- Whitson, H. E., Hastings, S. N., Landerman, L. R., Fillenbaum, G. G., Cohen, H. J., & Johnson, K. S. (2011). Black–White disparity in disability: The role of medical conditions. *Journal of the American Geriatrics Society*, 59(5), 844–850. <https://doi.org/10.1111/j.1532-5415.2011.03401.x>
- Whittemore, R., & Dixon, J. (2008). Chronic illness: The process of integration. *Journal of Clinical Nursing*, 17(7b), 177–187. <https://doi.org/10.1111/j.1365-2702.2007.02244.x>
- Wong, C. Y., Chaudhry, S. I., Desai, M. M., & Krumholz, H. M. (2011). Trends in comorbidity, disability, and polypharmacy in heart failure. *The American Journal of Medicine*, 124(2), 136–143. <https://doi.org/10.1016/j.amjmed.2010.08.017>

Yohannes, A. M., Willgoss, T. G., Baldwin, R. C., & Connolly, M. J. (2010). Depression and anxiety in chronic heart failure and chronic obstructive pulmonary disease: Prevalence, relevance, clinical implications and management principles. *International Journal of Geriatric Psychiatry*, 25(12), 1209–1221. <https://doi.org/10.1002/gps.2463>

CHAPTER 2: LITERATURE REVIEW AND THEORETICAL FRAMEWORK

This chapter includes two sections. The first is the theoretical framework and conceptual model that guided this study. Next, there will be a review of what is known regarding the relationships among sociodemographic, clinical, and psychological characteristics and quality of life (QOL) in Black and White patients with heart failure (HF).

Conceptual Model

The guiding conceptual model comprised the heart failure racial disparities reduction framework, a new framework I proposed based on a synthesis of the HF literature; the National Institute on Minority Health and Health Disparities (NIMHD) research framework; and Doll's (2018) epidemiological framework. These sources collectively informed the relevance of this exploratory comparative study, highlighted the importance of investigating the differences between Black and White individuals with HF, and contributed to the scholarly work of reducing disparities among Black individuals with HF.

The literature shows differences between Black and White individuals with HF (Afzal et al., 1999; Cuyjet & Akinboboye, 2014; Dickson et al., 2015; Durstenfeld et al., 2016; Glynn et al., 2019; Goyal et al., 2022; Mwansa et al., 2021; Nayak et al., 2020; Sharma et al., 2014; Smith et al., 2005; Su et al., 2019; Vaccarino et al., 2002), suggesting that race influences various phenomena in patients with HF. Because race-based health disparities persist and HF affects Black people more negatively (Tsao et al., 2022; Virani et al., 2020, 2021; Ziaeeian et al., 2017), studies designed with a disparities lens to examine racial differences between Black and White patients with HF are needed (Akwo et al., 2018; Echouffo-Tcheugui et al., 2022; Gordon et al., 2010; Lewsey & Breathett, 2021; Mentz et al., 2015; Nayak et al., 2020; Rathore et al., 2003; Sharma et al., 2014; Spahillari et al., 2017; Yancy, 2001).


The NIMHD provides a lens from which to conduct health disparities research. This framework suggests that individual patient domains affect health outcomes (NIMHD Minority Health and Health Disparities, 2017). This study aligned because it was an exploration of the relationships among individual patient factors, such as sociodemographic, clinical, and psychological characteristics, and their influence on QOL. The primary focus was comparing the observed relationships between Black and White patients with HF.

The third component of the heart failure racial disparities reduction framework (see Figure 3) is an epidemiological approach to investigating Black-White disparities in gynecologic cancer (Doll, 2018). Using Doll's (2018) framework in this study enabled comparing relationships observed in Black versus White patients with HF and helped interpret inequalities. Although inequity and inequality are different concepts, in U.S. society, systemic inequity contributes to specific health-related inequalities, which then play a role in perpetuating persistent racial disparities among patients with HF. Doll's framework was appropriate in this study to illuminate inequalities in this patient population that might contribute to disparities. It includes elements of three other frameworks for studying racial inequity—eco-social theory of disease distribution, fundamental cause theory, and public health critical race praxis—indicating four disease phases (see Figure 1):

1. Phase I: Natural Inequalities
2. Phase II: Increasing Inequalities
3. Phase III: Reducing Inequalities
4. Phase IV: Reduced Mortality

Figure 1

Doll's Theoretical Framework



Note. From “Investigating Black-White Disparities in Gynecologic Oncology: Theories, Conceptual Models, and Applications,” by K. M. Doll, 2018, *Gynecologic Oncology*, 149(1), pp. 78-83 (<https://doi.org/10.1016/j.ygyno.2017.10.002>). Copyright 2018 by Elsevier.

Each phase represents a stage on a continuum of evolution to reduce racial disparities. According to the heart failure racial disparities reduction framework (see Figure 2), racial disparities persist, increase, improve, or resolve based on inequalities. Addressing these inequalities involves designing targeted, efficacious interventions to generate knowledge and decrease racial disparities.

During Phase I: Natural Inequalities, there is limited knowledge about risk factors and few treatments available to reduce disparities within a patient population . As a result, the mortality rate remains stable (Doll, 2018). Characterizing Phase II: Increasing Inequalities are unequal diffusion of innovations, reduction of risk factors, and treatment strategies within the patient population. During this phase, mortality due to the illness decreases. However, there are increasing disparities because of the inequality of effective interventions across all subpopulations of individuals affected. Phase III: Reducing Inequalities involves increased

access to new knowledge and innovation applied to the patient population, including those with disparate outcomes, decreasing mortality rates, and disparities. Finally, Phase IV, Reduced Mortality, is characterized by widely available prevention strategies and treatment applied equitably, contributing to minimized or absent mortality rates and little to no remaining disparities.

Doll's (2018) theory indicates that health disparities can persist in a specific racial group despite increasing innovation and resources to decrease the illness's overall burden and effects. There is a need for research to address the racial differences that might contribute to persistent inequalities and health disparities. Targeted studies could inform tailored interventions to reduce disparities and promote more equitable patient outcomes.

Conceptual Model for the Current Study

This study incorporated the four phases of the heart failure racial disparities reduction framework (see Figure 2) into Doll's (2018) original framework (see Figure 3). However, rather than disparities in gynecological cancer, the four phases of inequality applied to inequalities, specifically, racial disparities in HF related to depression and anxiety in Black and White patients. Although Doll investigated Black–White disparities in gynecologic oncology patient populations, the framework is applicable to disparities research focused on potential differences between Black and White patients with HF. This study and future studies to identify racial differences could contribute to a growing body of knowledge on disparities in HF research.

Racial disparities persist within the HF patient population. Black individuals have worse outcomes than White persons (Tsao et al., 2022; Virani et al., 2020, 2021; Ziaieian et al., 2017), which is consistent with Phase II: Increasing Inequalities. Despite resources and innovations

directed at primary and secondary HF prevention, there has been no reduction in disparities among Black patients. Black patients with HF are understudied

The Heart Failure Racial Disparities Reduction Framework. The heart failure racial disparities reduction framework (see Figure 2) is an application of Doll's (2018) epidemiological framework with application to the HF patient population (see Figure 3). Consistent with Doll, Phase I includes the inequalities in HF influenced by sociodemographic characteristics (race, age, gender, marital status, education, and employment status), clinical characteristics (disease severity, comorbidities, and functional status), and psychological characteristics (depression and anxiety), characteristics that collectively influence QOL. The framework suggests that these characteristics are inherently different in Black and White patients with HF. In addition, the characteristics influence depression and anxiety in ways that perpetuate disparities among Blacks, including underdiagnosis and undertreatment (Dickson et al., 2013; Evangelista et al., 2009). The current study did not address Phase I on the continuum because morbidity and mortality rates among patients with HF are no longer stable. Instead, there is stability and continued growth in the morbidity, mortality, and disparity between Black and White patients with HF (Tsao et al., 2022; Virani et al., 2020, 2021).

The heart failure racial disparities reduction framework was appropriate for this study to explore phenomena in Black individuals' and determine whether racial differences exist by examining the potential influences of sociodemographic and clinical characteristics on psychological characteristics and their collective influence on QOL. This study focused on the interval between Phase II (Increasing Inequalities) and Phase III (Reducing Inequalities; see Figure 3). The persistent health disparities observed among Black individuals with HF align with the unequal diffusion of innovations, reduction of risk factors, and treatment strategies (Doll,

2018), perpetuated by the underrepresentation of Black patients in HF research (Sharma et al., 2014). Medical professionals have attempted to address racial disparities in patients with HF, albeit with limited success. There have been investments and scientific advances to better treat HF, improve clinical outcomes, and decrease morbidity and mortality within the general HF patient population (Braunwald, 2013; Lewsey & Breathett, 2021). However, these knowledge increases and innovative treatment approaches have received unequal application among Black individuals (Azam & Colvin, 2021; Heiat et al., 2002; Khayyat et al., 2021; Lekavich & Barksdale, 2016; Sullivan et al., 2018). Thus, the increases and advancements have not significantly contributed to improving clinical outcomes or decreasing health disparities among Black individuals with HF.

Researchers have investigated depression and anxiety in patients with HF, albeit with Black participants underrepresented in their samples (Sharma et al., 2014). Most investigators have only reported race descriptively, grouped all races and minorities, and inconsistently tested for racial differences. Rather than considering race as a significant predictor or contributor to the persistent disparities observed in Black individuals with HF, scholars have rarely unexplored race as a variable of potential influence. Depression and anxiety remain underdiagnosed and undertreated in Black individuals with HF (Dickson et al., 2013; Evangelista et al., 2009). Clinical outcome disparities among Black patients persist, including greater morbidity, greater health care utilization, more pronounced disability, and higher mortality (Tsao et al., 2022; Virani et al., 2020, 2021).

By underrepresenting Black patients in studies of depression and anxiety in patients with HF and not stratifying findings by race or comparing them to explore potential racial differences, researchers have contributed to persistent disparities among Black patients with HF. The

knowledge generated cannot be generalized to Blacks. Scientific innovation has improved clinical outcomes for the overall HF patient population (Braunwald, 2013; Lewsey & Breathett, 2021), yet Black individuals have not benefited as much from improvements (Tsao et al., 2022; Virani et al., 2020, 2021). Addressing this pattern of disparity requires a better understanding of depression and anxiety in Black patients and the sociodemographic and clinical characteristics that influence these psychological characteristics. Insight into the associations between and among sociodemographic, clinical, and psychological characteristics and QOL and whether racial differences exist is also warranted. The heart failure racial disparities reduction framework (see Figure 3) shows race as a central construct in HF health disparities research. The goal of applying this framework was to identify and account for any observed racial differences and ultimately decrease racial disparities by better understanding any differences between Black and White patients with HF so scholars and practitioners can address them.

The HF racial disparities reduction framework (see Figure 3) was appropriate due to the limited research on depression and anxiety in Black individuals with HF. The minimal literature on this subpopulation is insufficient to determine whether there are racial differences in the rates of depression and anxiety in Black and White individuals. The limited published literature provides insufficient insight into the sociodemographic and clinical characteristics that influence depression and anxiety (psychological characteristics) or how these factors might collectively influence QOL. Because most researchers have not included representative Black samples or conducted racial subgroup analyses when there are representative samples, more research is needed. Knowledge regarding racial differences is limited (Dickson et al., 2013; Evangelista et al., 2009), the findings are not generalizable to Black individuals (Bean et al., 2009), and disparate clinical outcomes, including mortality, persist (Tsao et al., 2022; Virani et al., 2020,

2021). Because race influences health, illness, and health care (Bhopal et al., 2021; Culley, 2006), it is a necessary variable in investigating disparities. Black individuals might perceive depression and anxiety differently than people from other races (Dickson et al., 2013). A cultural lens is necessary to investigate depression and anxiety within this subgroup of the HF patient population (Dickson et al., 2013; Nayak et al., 2020). In the era of persistent race-based disparities among Black individuals with HF, a better understanding of the prevalence of depression and anxiety in this patient population is needed to reduce disparities, illness burden, and mortality; improve care quality; decrease costs; and improve QOL (Celano et al., 2018; Cully et al., 2009; Dekker et al., 2014; Konstam et al., 2005; MacMahon & Lip, 2002; Rutledge et al., 2006; Yohannes et al., 2010).

Similar to its influence on health, race specifically influences HF, as patients of diverse racial groups do not experience HF the same as White patients (Dickson et al., 2013; Jaarsma et al., 2021; Riegel et al., 2008, 2009). Overall, Black individuals have a greater mortality risk than Whites, influenced by sociodemographic factors, including age, gender, and socioeconomic status (Beydoun et al., 2016). Black individuals with HF have higher mortality rates (Tsao et al., 2022; Virani et al., 2020, 2021), and although depression and anxiety are common in patients with HF, little research on depression and anxiety in patients with HF addresses potential racial differences. Researchers who consider sociodemographic characteristics and racial differences to understand patients' experiences without bias are more likely to better understand racially influenced phenomena (Doll, 2018; Nayak et al., 2020; Sharma et al., 2014).

Doll (2018) proposed stratifying research by race using univariate and multivariate modeling to consider the socioeconomic and sociocultural influences on the outcome of interest. Further, Doll encouraged exploring race as a variable to assess unique features within data sets

and to test mechanisms of known disparities, which this study aimed to accomplish. Although most research on patients with HF does not have findings stratified by race, scholars who choose to explore race as a variable of potential influence have identified racial disparities (Doshi et al., 2020; Minhas et al., 2022). Thus, stratification by race among patients with HF could help identify critical racial disparities to address. The current state of HF and accompanying disparities aligned with Phase II of the theoretical framework (see Figure 3). Research is needed to progress to Phase III, characterized by a reduction in inequalities secondary to new knowledge that informs the development of new interventions.

Figure 2

The Heart Failure Racial Disparities Reduction Framework





Figure 3

Application of Doll's Theoretical Framework to Disparities in HF



This and subsequent studies generate new knowledge to foster progression to Phase III: Reducing Inequalities. The current study was an exploration and comparison of the sociodemographic (race, age, gender, marital status, education, and employment status), clinical (disease severity, comorbidities, and functional status), and psychological (depression and anxiety) characteristics; how they influence each other; and how they might influence QOL in Black individuals with HF differently from White individuals (see Figure 3). Comparing these sociodemographic, clinical, and psychological characteristics was a means to identify racial differences between Black and White patients with HF that might contribute to the increasing inequalities phase (Phase II) among Black patients with HF (see Figure 3). The current study entailed testing whether differences exist, measuring the magnitude of any differences, and measuring the influence of depression and anxiety in Black individuals with HF. Findings from this study could inform interventions that address and ultimately decrease racial disparities in clinical outcomes among Black individuals with HF. This research generated foundational new knowledge to inform the diagnosis, care, and treatment of depression and anxiety in Black individuals with HF.

Phase III of the adapted version of Doll's (2018) heart failure racial disparities reduction framework suggests that developing increased knowledge and innovative approaches could improve the diagnosis and treatment of depression and anxiety in patients with HF (see Figure 3), with better QOL and overall HF clinical outcomes. Additional contributions were novel treatment strategies, culturally competent care, and culturally responsive interventions that account for racial differences. Success in Phase III will lead to Phase IV: Reduced Mortality, which occurs when prevention and treatment are widely available to all patients with HF, with racial disparities significantly reduced or eliminated (see Figure 3).

Leveraging Doll's adapted framework (see Figure 3) to inform the heart failure racial disparities reduction framework (see Figure 2) might help to advance knowledge regarding disparities within the HF patient population and advance health equity. HF patients, their care, and the healthcare providers caring for them are not isolated components. The greater context of society, which includes race as a societal construct, could influence phenomena in the HF patient population and thus merited research consideration (Doll, 2018).

Literature Review

This literature review is an exploration of what is known about the influence of sociodemographic and clinical characteristics on psychological characteristics when comparing Black and White persons with HF. The published research regarding the influence of sociodemographic, clinical, and psychological characteristics on QOL receives discussion, with a focus on comparing potential differences between Black and White individuals with HF.

Sociodemographic and Clinical Characteristics

Sociodemographic characteristics, including age (Cleland et al., 2007; Li et al., 2008), gender (Boersma et al., 2020; Cleland et al., 2007; Kalsoom, 2020; Liew, 2012), marital status (Pindikura et al., 2019; Wang et al., 2021; A.-Z. Zhang et al., 2016), education (Liew, 2012; Pindikura et al., 2019; Wang et al., 2021), and employment status (Almawi et al., 2014; Balcells et al., 2010), influence depression and anxiety in patients with chronic illnesses. Likewise, clinical characteristics, such as disease severity (Di Marco et al., 2006; Kim et al., 2014), comorbidities (Ahn et al., 2017; Bell et al., 2005; Lim et al., 2011), and functional status (Gerogianni et al., 2014; McCann & Boore, 2000; Pumar et al., 2014) also influence depression and anxiety in chronically ill individuals. Investigators have found associations between having depression and anxiety and having one or multiple physical chronic illnesses (Assari, 2016;

Balcells et al., 2010; Johnson-Lawrence et al., 2013; Katon et al., 2007; Liew, 2012; Read et al., 2017; Watkins et al., 2015). In addition, some Black individuals are more likely than persons of other races to have multiple chronic illnesses (Johnson-Lawrence et al., 2013; Warner et al., 2019). Blacks could be at the greatest risk for depression and anxiety. Depression and anxiety in chronically ill persons negatively influence functional status (Lim et al., 2011), often already diminished in patients with HF.

Sociodemographic Characteristics. Based on existing research, sociodemographic (race, gender, marital status, education, and employment status) and clinical (disease severity, comorbidities, and functional status) characteristics might also influence psychological characteristics (depression and anxiety) in patients with HF (Ahmed et al., 2006; Chung, Moser, et al., 2009; Dekker, Lennie, et al., 2014; Freedland, Rich, et al., 2003; Gottlieb et al., 2004; Jiang et al., 2001; Johansson et al., 2006ab; Vaccarino, Kasl, et al., 2001; Rohyans & Pressler, 2009; Rutledge et al., 2006; Sharma et al., 2009; Trivedi et al., 2009). These characteristics could affect depression and anxiety in Black individuals in diverse ways compared to White individuals with HF, which further investigation might indicate.

Because sociodemographic factors play a role in depression and anxiety, there is a strong likelihood of racial, ethnic, and sociocultural factors having diverse impacts on the experiences of Black patients with HF and depression and anxiety (Dickson, McCarthy, et al., 2013ab; Riegel et al., 2011). However, race has not been a variable of potential influence in most studies of depression and anxiety in patients with HF. Investigators studying depression and, to a lesser extent, anxiety in patients with HF have controlled for several sociodemographic characteristics, including age (Dekker et al., 2014; Freedland, Rich, Skala, et al., 2003), gender (Dekker et al., 2014; Gottlieb et al., 2004; Jackson & Emery, 2011), marital status (Luttik et al., 2006; Rohyans

& Pressler, 2009), education (Dekker et al., 2014; Gottlieb et al., 2004; Vaccarino et al., 2001), and employment status (Freedland, Rich, Skala, et al., 2003), and clinical variables, including disease severity (ejection fraction [EF] and New York Heart Association Classification [NYHA Class]; Benatar et al., 2003; Gottlieb et al., 2004), and comorbidities (Freedland, Rich, Skala, et al., 2003; Jiang et al., 2001; Sharma et al., 2009). However, researchers investigating depression and anxiety in patients with HF have rarely stratified depression and anxiety by race. There are likely racial differences when comparing minority patients, specifically Black and White (Rutledge et al., 2006).

Race. Race is a biological and social construct that influences health (Corbie-Smith et al., 2008) and a fluid concept used to group people based on various criteria, including ancestral background, social identity, and visible physical characteristics (Institute, n.d.). Race is an important consideration when investigating health disparities (Hill et al., 2015). In the present study, the conceptualization of race was the racial group (Black or White) with which a participant self-identified. Race received discussion within the context of knowledge regarding the sociodemographic, clinical, and psychological characteristics and QOL explored in this study.

Age. Age is the length of time from the beginning of a person's existence to a specific point in time (Merriam-Webster, n.d.-a). Age influences health in diverse ways (Charles & Piazza, 2009; Schöllgen et al., 2016; World Health Organization, n.d.) and should be a consideration when investigating various aspects of disorders (Geifman et al., 2013; Schöllgen et al., 2016). In this study, the concept of age was a self-reported numeric value that reflects an adult participant's chronological age.

Age is an important consideration when studying HF because HF symptoms vary based on age (Sethares & Chin, 2021). Research findings suggest that age influences depression and anxiety in patients with HF, psychological characteristics that might vary based on patient age (Celik et al., 2016; Dekker et al., 2014; Dickson et al., 2013; Easton et al., 2016; Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Rohyans & Pressler, 2009; Trivedi et al., 2009; Vaccarino et al., 2001). However, these age-related findings differ are unstratified by race. Few researchers have investigated whether age might influence depression and anxiety differently in Black individuals than White individuals with HF.

Some investigators have found that younger age is commonly associated with greater depression (Athilingam et al., 2017; Chung, Lennie, et al., 2009; Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Koenig, 1998; Mentz et al., 2015; Rohyans & Pressler, 2009; Stamp et al., 2014; Xiong et al., 2015) and anxiety (Dekker et al., 2014) in the general population of patients with HF. For example, Chung et al. (2009) found that younger age was associated with more depression. However, the sample was predominantly White (86%), and the researchers did not conduct a racial subgroup analysis on the Black sample. In another study, Celik et al. (2016) suggested that older age was associated with more depression but did not report race. However, in an all-Black sample of patients with HF, older age (> 62) was also significantly associated with more depression ($t = 2.147$, $p = 0.168$; Dickson et al., 2013).

Black individuals with HF tend to be younger than patients of other racial groups (Tandon et al., 2020; Virani et al., 2020, 2021) and might be at greater risk for depression and anxiety. Age could influence depression and anxiety differently among Black individuals. However, racial differences related to its influence are not well understood. In one of the few studies assessing relationships between age and anxiety, Dekker et al. (2014) found that younger

age was associated with more anxiety. However, the researchers did not stratify their findings by race and grouped all minorities, which prevented isolating the findings related to Black individuals with HF. Because researchers have not sufficiently studied the intersection of age and race as influencers on depression and anxiety in the HF patient population, the potential relationship and whether there are Black-White racial differences in the influence of age on depression and anxiety in patients with HF is unclear. Further research on this topic is needed to account for covariates (Sokoreli et al., 2016). This study was a test for age-related racial differences in depression and anxiety among patients with HF.

Gender. Gender is a social construct determined by biological factors and societal norms, roles, and values. It is a social and cultural determinant of health that influences health-related outcomes in various ways (ORWH, n.d.). Exploring relationships between gender and illness could help identify within-group and between-group differences (Phillips, 2005). In addition, including gender in research informs prevention strategies and interventions for both men and women (ORWH, n.d.; Phillips, 2005; World Health Organization, n.d.). In this study, gender was characterized by the group (male or female) with which participants self-identified.

Although gender influences symptoms in chronically ill individuals (O'Neill & Morrow, 2001), women are underrepresented in HF research (Reza et al., 2022). However, HF symptoms, including depression (Heo et al., 2019) and various phenomena in patients with HF vary by gender (Hopper et al., 2016). Gender might also influence depression and anxiety in Black and White individuals with HF, yet these potential relationships have not received sufficient study. Some researchers have found that women are more depressed than men (Ahmed et al., 2006; Gottlieb et al., 2004; Jiang et al., 2001; Vaccarino et al., 2001; Xiong et al., 2015). In contrast,

others identified no statistically significant differences in depression based on gender (Celik et al., 2016; Rohyans & Pressler, 2009; Sharma et al., 2009).

Ahmed et al. (2006) found that women in their prematched sample ($N = 2,040$) were more likely to be depressed than men ($p < 0.0001$), yet they did not stratify the findings by race. In another study, investigators found that women (64%) were more significantly depressed than men (44%), and Black women were more depressed than White women (Gottlieb et al., 2004), yet Black men were less depressed than White men. These varied findings suggest there might be racial differences in depression and anxiety based on gender, although more research is needed to determine potential relationships.

Although depression and anxiety are highly correlated (Celano et al., 2018; Herr et al., 2015), investigators have found differing relationships between anxiety and gender. For example, in a study conducted in Turkey, Celik et al. (2016) found that women were more anxious than men ($p < 0.05$). They observed no statistically significant relationship when comparing depression rates in males and females ($p > 0.05$). Race was not a variable reported in the study.

Because most researchers did not address the intersection of race and gender, it is unknown whether there are racial differences in the influence of gender on depression and anxiety in patients with HF. Some investigators (Dekker et al., 2014; Freedland et al., 2003; Gottlieb et al., 2004; Griffin et al., 2007; Jackson et al., 2011; Jiang, 2011; Rohyans & Pressler, 2009; Sharma et al., 2009; Vaccarino et al., 2001) have considered the influence of gender on depression and anxiety in patients with HF, but few have explored evidence suggesting there might be gender-related differences in depression and anxiety between Black and White patients with HF. As a result, the research is limited, and findings differ across studies. Although some investigators who have tested for gender differences have found that women with HF are more

depressed than men, women are not well represented in the existing research (Reza et al., 2022). It is unclear whether depression and anxiety prevalence differ in men and women or whether other racial subgroup differences exist. An exploration of the impact of gender on depression and anxiety and its intersection with race is needed to enhance understanding. This study was a means to test for the potential influence of gender on depression and anxiety and differences in this influence based on race.

Marital Status. Marital status is defined as the state of being married or unmarried according to the civil laws or customs of a country and can include subcategories, such as separated, single, divorced, or widowed (Merriam-Webster n.d.-b; OECD, n.d.). In this study, marital status was an individual's self-identified membership in one of the categories delineating currently, previously, or never married. Examining marital status as a variable that could influence depression and anxiety aligns with the premises that marital status can influence mental health (Waldron et al., 1997), and married individuals have better health and well-being (Grundström et al., 2021) and lower mortality (Drefahl, 2012; Manzoli et al., 2007). Married persons with chronic illnesses experience less depression (Lam & Perales, 2017; Wang et al., 2021) than unmarried individuals, and there have been observable differences by race (Williams et al., 1992). Married individuals with cardiovascular illnesses have better clinical outcomes, including fewer adverse events and lower mortality rates than unmarried persons (Dhindsa et al., 2020; Manfredini et al., 2017; Potluri et al., 2019; Schultz et al., 2017).

Limited research regarding the influence of marital status on depression in patients with HF has shown varied results. Marriage can serve as a protective factor in patients with HF (Chung, Lennie, et al., 2009; Rohrbaugh et al., 2006), perhaps by providing companionship and decreasing the social isolation that patients with HF often experience (Dickson, McCarthy, &

Katz, 2013). Living alone is linked to depression in patients with HF (Johansson et al., 2006). Conversely, a live-in partner can provide social support that buffers some of the adverse effects of depression in patients with HF (Lee et al., 2017). Some investigators found that married patients have more depression (Celik et al., 2016). In a study conducted in Greece with a hospitalized sample of 190 patients, Polikandrioti et al. (2019) found that unmarried (single, divorced, and widowed) participants were significantly more likely to have major depression (35.6% vs. 19.2%, $p = 0.046$). However, they did not report race. In comparison, Rohyans and Pressler (2009) found no statistically significant differences in depression between married and unmarried patients ($t = 1.41$, $p = 0.16$) in a sample of patients with HF ($N = 150$; 31% Black). Their findings were not stratified by race, so the presence of racial differences is unclear. Although Chung et al. (2009) found no significant differences in depression among married and nonmarried patients with HF (10.9 vs. 12.2, $p = 0.39$), the sample was predominantly White (86%). The researchers reported no stratified findings and no data regarding the racial composition of the non-White sample.

Despite minimal research on the relationship between marital status and anxiety, the limited evidence suggests that marital status and anxiety are not associated (Celik et al., 2016). Although Celik et al. (2016) found a 33% anxiety prevalence in their sample, they observed no relationships between marital status and anxiety ($p = 0.98$) and did not report race. Despite infrequent explorations of the relationship between marital status and depression, investigators frequently examine relationships between marital status and clinical outcomes (i.e., rehospitalization and mortality). Chung et al. (2009) conducted a longitudinal study of a sample of patients with HF ($N = 166$) recruited from outpatient clinics at an academic medical center. The results showed that even when depressed, married patients with HF have lower morbidity

and mortality rates than unmarried patients ($p = 0.012$), suggesting the potentially protective influence of having a marital partner. However, the researchers did not measure anxiety or report race among the sociodemographic variables.

In a study of patients with HF conducted in the Netherlands, investigators found that being married positively influenced clinical outcomes, such as rehospitalization rates, mortality, and QOL (Luttik et al., 2006). These relationships might be related to less depression and anxiety due to marital status and unmarried patients' lack of social support from a spouse. However, Luttik et al. (2006) did not report race, and more evidence is needed to support these potential relationships. Similarly, a recent meta-analysis of 10 studies to explore the influence of marital status on HF outcomes showed that unmarried status is associated with negative clinical outcomes, including increased risk of mortality (pooled OR = 1.52, 95% CI [1.30–1.78], $p < .001$), increased risk of rehospitalization (pooled OR = 1.80, 95% CI [1.18–2.74], $p = .007$), and increased risk of combined endpoint of mortality and rehospitalization (pooled OR = 1.72, 95% CI = 1.36–2.17, $p < .001$; Kewcharoen et al., 2021). These negative clinical outcomes could be related to undiagnosed depression and anxiety among unmarried patients with HF. Marital status might influence outcomes in patients with HF, but more evidence is needed to support this potential relationship.

Although some researchers found evidence suggesting that marital status might influence clinical outcomes in patients with HF, other researchers concluded that marital status did not influence clinical outcomes in patients with HF. In a study of 357 patients with HF, Watkins et al. (2013) found no association between marital status and negative clinical outcomes (HR = 0.71; 95% CI [0.35–1.49]). There were missing data, and although the sample included 183

Black patients, findings were not stratified by race, preventing comparisons between Black and White patients.

Marital status might influence depression and anxiety in patients with HF, and racial differences could exist. Still, there is very little specific evidence to support this potential relationship, and existing findings have not been stratified by race. Further investigation is needed to understand these potential relationships and determine if there are racial differences in the influence of marital status on depression and anxiety in patients with HF. This study included an assessment of the potential influence of marital status on depression and anxiety, with results stratified by race to better understand any potential relationships and if there are racial differences between Black and White individuals.

Education. Educational attainment level is “the highest level of schooling completed” or “the status of learning that has been achieved by a student” (American Psychological Association, 2022) attained by meeting a set of standards for acquired knowledge that individuals must master before completing an educational level. In this study, educational attainment level was characterized as education and referred to the highest level of education completed by each participant. Relationships between education and health are well-established (Ross & Wu, 1995), with greater educational attainment contributing to more positive health outcomes (Brunello et al., 2016; Ross & Mirowsky, 1999). Higher educational status is associated with greater health literacy (Andrus & Roth, 2002; Kickbusch, 2001), leading to better health outcomes (Berkman et al., 2011; Ferguson & Pawlak, 2011). Researchers have established relationships between higher levels of education and less depression and anxiety (Bjelland et al., 2008), with other studies showing that more-educated patients with cardiovascular illnesses have less depression and anxiety (Gorini et al., 2020).

Among patients with HF, depression and anxiety might differ based on education. However, few investigators have explored the relationships between education, depression, and anxiety (Celik et al., 2016; Gottlieb et al., 2004; Vaccarino et al., 2002), and there are few, if any, studies exploring the influence of education on depression and anxiety at the intersection of race. In a study with a majority-White patient sample (86% of $N = 166$), Chung, Lennie, et al. (2009) found that patients with HF who were less educated had more depression (44.5% vs. 63.6%, $p = 0.032$). Because the researchers did not report the races of the remaining 14% of the sample, whether there were racial differences in this observed relationship is unknown. Similarly, in a study with 231 participants in Greece, Aggelopoulou et al. (2017) found that lower educational status was associated with high levels of depression and anxiety. The researchers did not report race.

In the United States, Dekker et al. (2014) found an association between fewer years of education and more depression (10.8 vs. 13.5, $p < 0.001$) and more anxiety (33% vs. 17%, $p < 0.001$) in patients with HF. The sample included Black individuals, but the researchers grouped all minorities ($n = 206$); as a result, findings related to Blacks cannot be determined. Vaccarino et al. (2001) reported similar findings, indicating that less education was significantly associated with more depression ($p = 0.004$) in patients with HF. However, the limitations included a small sample size ($N = 79$), only 20% Black participants, missing data, and findings not stratified by race. In another study conducted in the United States, Gottlieb et al. (2004) found no differences in depression based on educational status ($p = 0.232$) in a sample of 155 patients with HF (57% Blacks, $n = 89$), yet without stratifying the results by race.

In a study conducted in Turkey, Celik et al. (2016) reported that lower educational status was associated with more severe depression ($p = 0.02$). There was no association between

anxiety and education ($p = 0.28$). As with other studies, the researchers did not report race.

Likewise, in a study in the Netherlands with a sample of 268 patients with HF, greater depression was associated with higher educational status ($\beta = 1.41$, $SE = 0.48$, $p = < 0.005$). However, anxiety was unmeasured and race unreported.

Few researchers have explored the relationship between education and anxiety in patients with HF. Dekker et al. (2014) found higher levels of anxiety among participants in their American sample ($N = 556$) who were less educated (less than a high school diploma; 33% vs. 17%, $p < 0.001$). Although the researchers included approximately 200 minorities in the sample, they did not isolate race among the minority group or stratify the findings by race. Yet, this finding on the overall sample is consistent with other studies that showed associations between lower levels of education and greater anxiety (Bjelland et al., 2008; Thurston et al., 2006).

Although education is not a variable frequently studied in relation to depression and anxiety in patients with HF, these potential relationships require exploration because low health literacy is related to lower educational status in patients with HF (Cajita et al., 2016; Peterson et al., 2011) and health literacy is often lower among Black individuals (Chaudhry et al., 2011; Morrow et al., 2006; Westlake et al., 2013). Additionally, lower health literacy is related to poor adherence to the prescribed HF regimen (Cajita et al., 2016; Deek et al., 2020) and negative clinical outcomes (Fabbri et al., 2020; Peterson et al., 2011) and might be associated with greater depression and anxiety. Poor adherence often leads to increased symptom burden and other adverse clinical outcomes, indirectly increasing depression and anxiety in patients with HF (De Jong et al., 2011; Easton et al., 2016; Rechenberg et al., 2020). Low health literacy might be indirectly related to educational status, but this potential relationship requires further exploration with a focus on testing for potential racial differences between Black and White individuals.

Education has implications for socioeconomic status (SES) and is an important variable in health disparities research (Shavers, 2007; Winkleby et al., 1992). Due to the limited explorations of relationships between education, depression, and anxiety, there is very little known about the presence of racial differences in the influence of education on depression and anxiety in Black and White individuals with HF. Because this study focused on potential racial differences between Black and White individuals with HF, education was a variable of interest to determine its association with depression and anxiety in patients with HF and whether there were racial differences between Black and White individuals.

Employment Status. Employment status is a person's situation in relation to the labor force. This status includes those who are engaged in work for pay or profit (employed), those who are jobless but available to work and have actively sought work within the past 4 weeks (unemployed), those out of the labor force (retired), and those who are eligible for the labor force but are neither employed nor unemployed (HealthyPeople.gov, n.d.). In this study, employment status was characterized as paid or volunteer work completed by the participant. Employment status is an important variable to consider when investigating health disparities, chronic illnesses, and health outcomes (Jones & Crews, 2013; Landsbergis et al., 2018) in Black individuals because employment has implications for SES (Stronks et al., 1997). In addition, financially distressed Black individuals are at risk for depression (Assari, 2019), highlighting the need to investigate the relationship between depression, anxiety, and employment status.

Although employment status is a critical sociodemographic variable to study within the HF patient population, it receives little consideration for its potential relationship to depression and anxiety in this patient population. However, because many patients with HF are disabled and unable to work, and unemployment can influence depression and anxiety (Ohlsson et al., 2021),

there could be an unexplored relationship. In a study of inpatients with HF, Freedland et al. (2003) found that unemployment due to disability was significantly related to depression ($n = 289, p = .0002$, and $n = 259, p = .0001$), respectively, when assessed using two different measures of depression. However, the investigators did not stratify the findings by race. Ohlsson et al. (2021) identified a relationship between unemployment and increased mortality (adjusted HR = 1.76; 1.47, 2.11) among patients with HF, and adjusting for age and gender did not significantly change this estimate. However, the researchers conducted their study in Sweden and did not report race.

Unemployed and disabled patients with HF have greater mortality and worse clinical outcomes (García-Olmos et al., 2019; Ohlsson et al., 2021; Rørth et al., 2018). When considering disparities in depression and anxiety among Black and White individuals with HF, employment status merits investigation because economic burden contributes to the development and progression of depression in patients with HF (Havranek et al., 2004) and might contribute to negative clinical outcomes. Compared to White individuals, Black persons traditionally have lower SES (including income; Anderson & Armstead, 1995; Averbuch et al., 2021) and significantly more disabilities (Van Nuys et al., 2018). These racial differences might contribute to greater depression and anxiety in Black patients. However, there is insufficient evidence to support a relationship between employment status, depression, and anxiety in Blacks and whether differences are apparent when comparing Black and White patients.

Patients with HF need income from employment to meet their financial obligations, including the costly requirements of HF care and treatment. However, most are unemployed and receive disability benefits, as HF's chronic and debilitating nature limits their ability to maintain employment (Seo et al., 2021). Because disability benefits are often lower than patients' previous

earnings, their financial needs might exceed available resources. Patients with HF who are less comfortable with their financial position are more likely to be depressed (Chung et al., 2009). For example, Cirelli et al. (2018) reported a relationship between home ownership and depression in a sample of patients with HF ($N = 309$), suggesting that patients with HF who have financial concerns might be at risk for depression. However, although the sample included 60 Black participants, the findings were not stratified by race.

In addition, multiple HF symptoms experienced by patients often significantly limit activities of daily living (ADLs), and many patients with HF have pronounced functional status limitations (Blinderman et al., 2008; Falk et al., 2008; Herr, Salyer, Flattery, et al., 2015; Herr, Salyer, Lyon, et al., 2014; Moser et al., 2014), which might limit their participation in physically demanding occupations they held prior to their HF diagnosis (Dickson et al., 2008). Patients must also engage in complex HF management that takes enough time to be considered work (Granger et al., 2009), which limits their capacity for employment. Individuals living with HF experience exacerbations that result in worsened symptoms, the need for healthcare provider intervention and hospitalization, and a restricted physical inability to work productively. This cyclical pattern of exacerbations and rehospitalizations make it challenging for patients with HF to maintain employment (Dickson et al., 2008).

Black individuals are diagnosed with HF at younger ages than their White counterparts (Virani et al., 2020, 2021). Younger patients might be at greater risk for depression because their illness prevents them from functioning in their expected roles, including employment (Johansson et al., 2006). The age and clinical outcome disparities among Black individuals with HF suggest that employment status could be an important variable to consider in this subpopulation of patients with HF. Specifically, these younger Black individuals might have financial obligations

unmet by disability benefits, contributing to depression and anxiety. Financial strain can contribute to nonadherence with the prescribed treatment regimen (Tansey, 2010), leading to increased rehospitalizations, poor QOL, and a cyclical pattern of depression and anxiety. Conversely, employment can negatively impact patients' ability to effectively manage HF and engage in the prescribed regimen. However, employment can be a mood enhancer for some patients with HF (Dickson et al., 2008). Thus, employment status needs further exploration.

Despite the extremely limited study of employment status, especially in Black individuals with HF, employment is an essential measurement when considering racial disparities among chronically ill individuals (Beatty & Joffe, 2006; Vassilev et al., 2014; von Houtum et al., 2015; Walker, 2010; Wilson, 2001) and thereby, those with HF. Although investigations of relationships between depression, anxiety, and employment status are scarce, there could be relationships because of the economic burden of unemployment. Further study is needed to measure the influence of employment status on depression and anxiety in Black individuals with HF and compare this outcome to White individuals to determine whether there are racial differences. This study was an investigation of the potential relationships between employment status, depression, and anxiety and differences between Black and White individuals with HF.

Clinical Characteristics

Investigators have found associations between clinical characteristics, including disease severity (Dekker et al., 2014; Evangelista et al., 2009; Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Jiang et al., 2001; McCarthy et al., 2015; Mentz et al., 2015; Rohyans & Pressler, 2009; Sharma et al., 2009), comorbidities (Freedland et al., 2003; Sharma et al., 2009), and functional status (Dekker et al., 2014; Herr, Salyer, Flattery, et al., 2015; Shen et al., 2010) with depression and anxiety. However, the findings differ and are not stratified by race. Clinical

characteristics might influence anxiety, but these potential relationships in patients with HF is minimal and race-based findings are seldom, if ever, explored. Furthermore, instead of examinations for their potential influence on depression and anxiety, clinical variables are most often only reported descriptively in research studies.

Disease Severity. In patients with HF, disease severity refers to the accepted clinical criteria that denote the magnitude of symptoms and how HF limits physical activity and usual functioning (American Heart Association, 2017). In this study, disease severity conceptualization was with two variables: the left ventricular EF and NYHA Class. Left ventricular EF is the quantity of blood pumped out by the heart's left ventricle with each heartbeat. Expressed as a percentage, EF has a normal range of 50% to 70%. The categories of EF are preserved (diastolic HF; $< 40\%$) and reduced (systolic HF; $< 35\%$). The heart muscle contracts normally with a preserved EF, but the ventricles do not relax as they should when refilling with blood. A reduced EF is evident when the heart does not contract effectively, thus pumping out less oxygen-rich blood to the body (Katz, 2013). In this study, EF was characterized as a percentage and was one of two values used to measure disease severity.

Relationships between depression, anxiety, and disease severity receive infrequent investigation; however, racial differences might exist when comparing Black and White individuals with HF. The NYHA Class provides a simple means of measuring the extent of HF, classifying patients into one of four categories based on their limitations during physical activity regarding normal breathing and varying degrees of shortness of breath and angina pain. NYHA Class includes four categories based on intensity, from Class I (mild) to Class IV (most severe). Patients in Class I have no activity limitations and no symptoms with ordinary activity. Patients with HF classified as NYHA Class II have slight limitations and slight to moderate symptoms

with normal activity. Patients categorized as Class III have marked activity limitations and symptoms with less-than-normal activity. Finally, Class IV patients might have symptoms at rest and be unable to carry out physical activity without discomfort (American Hospital Association, 2017; Fleg et al., 2000). In this study, NYHA Class was conceptualized as a second measure of disease severity because it might have influenced depression, anxiety, and QOL differently in Black and White individuals with HF.

Ejection Fraction. Although this is a relationship not well explored by race, disease severity, as measured by the EF, is associated with depression in patients with HF (Adelborg et al., 2016). Freedland et al. (2003) found that depression was more prevalent among participants with lower EF in their sample ($N = 682$), which included 278 Black individuals (41%), but findings were not stratified by race. In a study of 172 patients with HF, Steca et al. (2013) reported an inverse relationship between disease severity and depression, suggesting that lower EF (worse disease severity) was associated with greater depression ($p < 0.001$). However, they conducted the study in Italy and did not report race. Adelborg et al. (2016) conducted another large study of patients with HF ($N = 204,523$) sampled from the Danish medical registries. The results showed that a history of depression was associated with all-cause mortality in patients with HF who had an $EF \leq 35\%$ (MRR = 1.14, 95% CI [1.12–1.17]), which represented an approximately 20% higher mortality rate than patients without a history of depression.

If more investigators explored racial differences in outcomes based on EF, perhaps there would be more knowledge regarding the influence of EF on depression and anxiety by race. However, there is little knowledge because only a few scholars have explored relationships between EF, depression, and anxiety. Depression and anxiety might be associated with greater disease severity, as measured by the EF, and there could be differences between Black and White

individuals. More evidence is needed to support potential relationships, especially related to anxiety. This study was an exploration of relationships between disease severity as measured by the EF, depression, and anxiety, comparing findings in Black and White patients with HF.

NYHA Class. Relationships between NYHA Class and depression and anxiety have received minimal study in patients with HF. Although some researchers have found associations between disease severity (NYHA Class) and greater depression (Bean et al., 2009; Chung et al., 2009; De Jong et al., 2011; de Leon et al., 2009; Dekker et al., 2014; Dickson, McCarthy, & Katz, 2013; Freedland, Rich, et al., 2003; Gottlieb et al., 2004; Rohyans & Pressler, 2009; Sharma et al., 2009; Trivedi et al., 2009), few considered race. In a study of 268 patients with HF in the Netherlands, Brouwers et al. (2014) found that depression was significantly associated with NYHA Class ($\beta = 2.43$, $SE = 0.75$, $p = 0.001$) at baseline (2.44 , $SE = 0.74$, $p = 0.001$) and the 12-month follow-up ($\beta = 2.37$, $SE = 0.85$, $p = 0.005$; $OR = 4.54$, $p = 0.02$), even after adjusting for sociodemographic factors ($OR = 4.85$, $p = 0.02$). These results suggest a strong correlation between NYHA Class as a measure of disease severity and depression. Brouwers et al.'s study had limitations, as they did not examine anxiety or report race. In research conducted in Turkey, Celik et al. (2016) found no relationship between anxiety and disease severity as measured by the NYHA Class ($p = 0.31$), although a relationship emerged between depression and the NYHA Class ($p \leq 0.001$). Race was not reported. In a U.S. study, Chung, Moser, et al. (2009) identified an association between depression and higher NYHA Class (greater disease severity). Patients with depression had worse functional status (NYHA Class III–IV 83.3% vs. 50.5%, $p < 0.001$). However, White patients accounted for the majority of this sample (86%) and no findings were reported by race.

Dekker, Lennie, et al. (2014) found that patients with more severe HF (higher NYHA Class III or IV) were significantly more likely to be depressed (75% vs. 43%, $p < 0.001$) and anxious (64% vs. 41%, $p < 0.001$). Although the sample included 200 participants reported as a group of “minorities” who were “mostly Black,” the findings were not stratified by race and cannot be generalized to Black patients. In a mixed methods study of an all-Black sample of patients with HF ($N = 30$), Dickson et al. (2013) found an association between NYHA Class III and greater levels of depression ($t = - 2.477$, $p = 0.02$). Wierenga et al. (2017) reported findings related to clinical outcomes based on race. A secondary data analysis showed that NYHA Class (III and IV; 1.24, 95% CI [0.4–3.81], $p = 0.711$) and depressive symptoms ($\beta = 1.02$, 95% CI [1.02, 1.18], $p = 0.018$) were statistically significant predictors of shorter event-free survival in White but not Black individuals. The overall sample ($N = 863$) came from a multisite HF registry, and the Black sample was small ($n = 91$; 10.5%). This relationship might be related to the association between disease severity as measured by the NYHA Class, but more evidence is needed to support this potential relationship.

Among the limited research focused on relationships between anxiety and disease severity (NYHA Class), two studies found higher rates of anxiety based on NYHA Class (Dekker, Lennie, et al., 2014; Freedland et al., 2003). In a meta-analysis, Easton et al. (2016) found that lower NYHA Class was associated with increased anxiety. Few studies have explored the influence of disease severity on depression and anxiety, and the literature review showed only one that specifically tested for racial differences between Black and White individuals. Even fewer researchers have used both the NYHA Class and EF as measures of disease severity. With the employment of two measures, this study had an innovative approach to investigating the influence of disease severity on depression and anxiety. The potential influence of disease

severity on depression and anxiety in Black and White patients with HF was also assessed to identify potential racial differences.

Comorbidities. Comorbidities are illnesses, diseases, or disorders simultaneously present in an individual (American Psychological Association, APA, 2022). In this study, comorbidities were conditions that co-occur with HF (Formiga et al., 2018), as determined by calculating a weighted taxonomy of co-occurring conditions and their prognostic implications (Charlson et al., 1987). Comorbidities are common in patients with HF (Dickson, Buck, & Riegel, 2011; Lang & Mancini, 2007; Wiley et al., 2018; Wong et al., 2011). They are associated with depression (Patel et al., 2018) and anxiety (Dekker et al., 2014) in patients with HF, and Black individuals with HF have more comorbidities than White individuals (Sharma et al., 2014). Measuring the influence of comorbidities on psychological characteristics and QOL is important because comorbidities can result in patients experiencing overlapping symptoms, which can influence study findings (de Groot et al., 2003). Among the common comorbidities observed in patients with HF are depression and anxiety (Celano et al., 2018; Dekker, Lennie, et al., 2014; Konstam et al., 2005; Lang & Mancini, 2007; Sokoreli et al., 2016; Yohannes et al., 2010). There might be racial differences in the number and influence of comorbidities on psychological, sociodemographic, and other clinical characteristics and QOL.

Ahmed et al. (2006) found that greater depression was associated with comorbidities (hypothyroidism, dementia, hypertension, coronary artery disease, diabetes, cardiac dysrhythmias, and acute renal failure) in a sample of 2,040 patients with HF. Although they included 240 Black patients, the researchers did not stratify their findings by race. Assari, Dejman, and Neighbors (2015) found comorbidities associated with depression in White patients with HF, but this association did not emerge in Blacks. From a predominantly Black sample,

Sharma et al. (2009) concluded that comorbidities were associated with more depression in White patients with HF, but this association was not apparent in Blacks ($n = 116$, 86%). However, Freedland et al. (2016) found that depression rates were not affected by comorbidities, including diabetes or renal disease.

Relationships between comorbidities and anxiety are rarely explored. There is very minimal research on the general HF patient population, and none focused on differences between Black and White individuals with HF. Relationships between race, comorbidities, depression, and anxiety in patients with HF have not received sufficient exploration. Despite the limited data, the presence of multiple chronic illnesses might negatively influence Black individuals differently than White individuals. Black individuals with HF might have more comorbidities, such as hypertension, coronary artery disease, and diabetes, than their White counterparts (O'Connor et al., 2019; Parashar et al., 2009). There might be other racial differences related to comorbidities influencing levels of depression and anxiety. Although multiple chronic illnesses could adversely affect depression and anxiety in Black individuals differently than in White individuals, the potential relationships are unclear. Researchers have not presented a good understanding of whether there are racial differences in the influence of comorbidities on depression and anxiety. This study was a means to test for racial differences in the influence of comorbidities on depression and anxiety in Black and White individuals with HF.

Functional Status. Most patients with HF experience decreased functioning and activity limitations, which some investigators and providers might perceive as causing them to underestimate their functional abilities (Gottlieb et al., 2004). Patients whose functional status is not significantly and immediately affected by disease progression and intensified symptoms later experience functional status declines (Wong et al., 2011). These individuals' struggles with

changes in their lives, roles, functions, and independence contribute to depression and anxiety (Herr et al., 2015; Shen et al., 2010). For example, in an all-Black sample of patients with HF ($N = 30$), Dickson, McCarthy, and Katz (2013) identified a significant association between depression levels and lower physical functioning status ($r = -0.418, p = 0.02$). After controlling for other factors, Vaccarino, Kasl, et al. (2001) noted that functional decline at 6 months was higher among patients with more significant depression. Little is known regarding whether functional status declines or depression is the precursor for the other observation.

However, findings related to depression, anxiety, and functional status are limited overall. In addition, there is not a good understanding of whether differences in the relationships between functional status, depression, and anxiety exist between Black and White individuals. Limited studies have suggested relationships between depression, anxiety, and functional status with racial differences between Black and White individuals. For example, investigators have found that patients who are unable to perform ADLs were more depressed, and younger Blacks were more likely to have worse functional status (de Leon et al., 2009; Freedland et al., 2003).

However, there were methodological issues, including the sampling of Black participants. Specifically, de Leon et al. (2009) and Freedland et al. (2003) excluded patients from the studies who were more likely to be Black, which could have confounded the results.

Clarke et al. (2000) found that patients with HF who were depressed and anxious and of a non-White race were more likely to experience severe limitations in ADL functioning after 1 year. However, the findings were not statistically significantly different when controlling for disease severity and comorbidities. A limitation was that Clarke et al. included all minority participants together rather than stratifying findings by race, thus preventing generalizing the findings to Black individuals.

Two other studies among patients with HF found that younger patients had worse general functioning than elder patients (de Leon et al., 2009; Gottlieb et al., 2004), suggesting that depression and anxiety might be even more debilitating for younger patients with HF and could have grave impacts on their functional status. There is a decreasing average HF patient age and an increase in the likelihood that Black individuals with HF are younger (Sharma et al., 2014; Tandon et al., 2019). Because Black individuals with HF are more likely to be younger than White individuals, depression and anxiety could be even more debilitating for younger patients, specifically younger Blacks with HF. As a result, functional status limitations could contribute to more depression and anxiety in Black individuals because of discrepancies between their realized functional status and their functional status expectations based on chronological age. These functional status limitations might contribute to depression and anxiety, and relationships could differ between Black and White individuals (Dickson, McCarthy, & Katz, 2013). However, there is insufficient evidence to support this potential relationship.

Depression and anxiety can result in decreased participation in the HF treatment plan, leading to worse health status and ultimately contributing to diminished functioning among patients with HF (Dickson, McCarthy, et al., 2013; Dickson, McCarthy, & Katz, 2013). Lack of patient participation in the treatment plan results in a cyclical pattern contributing to negative clinical outcomes (Sevilla-Cazes et al., 2018). Although studies regarding functional status, depression, and anxiety are limited—and those considering race even rarer—available evidence suggests that these potential relationships warrant further exploration.

Findings regarding relationships between functional status and depression and anxiety in patients with HF are difficult to compare because researchers have used various measures of functional status. Among the measures used are the NYHA Class (Bean et al., 2009; De Jong et

al., 2011; de Leon et al., 2009; Dekker, Lennie, et al., 2014; Dickson, McCarthy, & Katz, 2013; Freedland et al., 2003; Gottlieb et al., 2004; Rohyans & Pressler, 2009; Sharma et al., 2009; Trivedi et al., 2009) and the Duke Activity Status Index (Dekker, Lennie, et al., 2014; Dickson, McCarthy, & Katz, 2013). However, the NYHA Class is not optimal as a sole measure of disease severity because it is a clinician's assessment of a patient's functional status at a specific time point. Because HF involves frequent fluctuations in severity and patient condition a documented NYHA Class is accurate at the time of assessment and might not remain accurate indefinitely. Additionally, disease severity should include patients' subjective perception that is not captured by and might differ from a healthcare professional's assessment of their condition (Caraballo et al., 2019; Raphael et al., 2007; Severo et al., 2011).

Still, the research shows an association between depression and lower functional status as measured by the NYHA Class (Dekker et al., 2014; Dickson, McCarthy, & Katz, 2013). The present study assessed NYHA Class and EF as measures of disease severity and measured their relationships with depression, anxiety, and QOL. Functional status was also assessed using a valid and reliable measure and its relationship to depression, anxiety, and QOL was investigated.

Functional status is an important variable to measure because declines are common in patients with HF (Salyer, 2015). Declines might directly or indirectly relate to depression and anxiety, influencing patients' energy levels and engagement in managing HF. Thus, improving functional status might reduce depression (X. Zhang et al., 2020). However, most existing studies did not report or compare findings by race. Obtaining a better understanding of the potential relationships and whether there are differences between Black and White patients with HF is needed.

Limited evidence suggests there might be relationships between sociodemographic, clinical, and psychological characteristics in the general HF patient population. However, not enough is known about these potential relationships and the influence of clinical characteristics (disease severity, comorbidities, and functional status) on Black individuals. The presence of differences between Black and White individuals remains unclear due to minimal research comparing racial groups. There is a need for further investigation regarding the potential influences of these clinical characteristics on depression and anxiety and whether any racial differences exist. This study was an exploration of whether there are differences in the influence of sociodemographic characteristics (age, gender, marital status, education, employment status), clinical characteristics (disease severity, comorbidities, and functional status) on psychological characteristics (depression and anxiety) and QOL and a comparison of the potential influences by race in Black and White individuals with HF.

Psychological Characteristics

Chronically ill patients frequently experience depression and anxiety (Assari, Dejman, & Neighbors, 2016; DeJean et al., 2013; Yohannes et al., 2010), symptoms commonly observed in patients with HF. Depression and anxiety can occur independently (Cully et al., 2009; Konstam et al., 2005; Sokoreli et al., 2016; Tsabedze et al., 2021) or together (Celano et al., 2018; Dekker, Lennie, et al., 2014; Yohannes et al., 2010). Although both conditions negatively impact patients with HF, the relationship between depression and anxiety and their comorbidity is not well understood. Little is known about the potential difference in relationships between Black and White patients with HF.

Depression. Depression and depressive symptoms are common in patients with HF (Celano et al., 2018; Ishak et al., 2020; Joynt et al., 2004; Konstam et al., 2005; MacMahon &

Lip, 2002; Rutledge et al., 2006). Although depressive symptoms present similarly to depression, they are less severe and do not meet the diagnostic criteria for depression; however, both conditions negatively impact patients with HF (Alhurani et al., 2015; Celano et al., 2018; Konstam et al., 2005; Nair et al., 2012; Rustad et al., 2013). In this study, “depression” represented both clinical depression and depressive symptoms (see Ahmed et al., 2006).

According to the American Psychological Association (2020), depression, also referred to as major depressive disorder, is a common, serious illness that lasts 2 or more weeks and negatively affects thoughts, feelings, and actions. Depression results in emotional and physical symptoms of varied intensity, including sadness, despondence, hopelessness, helplessness, apathy, loss of interest or pleasure in activities once enjoyable, changes in appetite (weight gain or loss unrelated to dieting), anergia, fatigue, increase in purposeless physical activity (or slowed movements or speech), feeling worthless or guilty, difficulty thinking, concentrating, and making decisions, insomnia, sleeping too much, and suicidal thoughts (American Psychiatric Association, 2020). Depression and depressive symptoms are positively correlated with anxiety (Dickson et al., 2013;).

Depression contributes to adverse clinical outcomes (Freedland, Hesseler, et al., 2016; Sokoreli et al., 2016) yet remains understudied in all patients with HF (Ahmed et al., 2006; Bhatt et al., 2016; Holzapfel et al., 2008). Although Black individuals have the worst clinical outcomes of any racial subpopulation with HF (Benjamin et al., 2017; Go et al., 2013; Roger et al., 2012; Tsao et al., 2022; Virani et al., 2020, 2021), they are underrepresented in depression-related HF research (Evangelista et al., 2009).

Much like the differences observed when comparing Black and White patients in other chronically ill populations (Ani et al., 2009; Assari, Burgard, & Zivin, 2015; Assari, Dejman, &

Neighbors, 2014; Ellis et al., 2015; Ghods et al., 2008; Hill-Joseph, 2019; Hollingshaus & Utz, 2013; Johnson-Lawrence et al., 2013; Li et al., 2008; Mouzon, 2017), differences in depression could become apparent with the exploration of racial differences. However, studies comparing depression rates in Black and White individuals with HF are limited (Evangelista et al., 2009), and findings vary. Some researchers found Black individuals more depressed than White individuals (Dekker, Lennie, et al., 2014; Evangelista et al., 2009). Other studies showed less depression in Black people compared to White (Ahmed et al., 2006; Koenig, 1998; Stamp et al., 2014; Vaccarino, Kasl, et al., 2001; Wierenga et al., 2017) or no statistically significant racial differences in depression between Black and White individuals with HF (Freedland et al., 2003; Gottlieb et al., 2004; Griffin et al., 2007; Piamjariyakul et al., 2013; Rohyans & Pressler, 2009; Xiong et al., 2015).

The limited studies showing Black individuals had more depression than Whites (Dekker et al., 2014; Evangelista, Ter-Galstanyan, Moughrabi, & Moser, 2009) had small Black samples or other methodological issues. More research is needed to better understand whether there are differences in depression in Black and White individuals with HF. For example, in a secondary analysis of an outpatient sample of patients with HF, Dekker et al. (2014) reported that minorities, mostly classified as Black ($N = 200$; 36%), were more depressed than Whites. However, grouping all minorities might have confounded race-related findings.

From a sample of 241 patients with HF (7% Black; $n = 18$) recruited from an outpatient HF clinic, Evangelista et al. (2009) found that Black race was a significant predictor of depression ($\beta = -0.153$, $p = 0.020$). Although there was a trend when comparing depression in Black and White individuals with HF, these relationships were not statistically significant ($p = 0.063$). A post hoc analysis indicated that Blacks were more likely to be depressed than

Hispanics ($F = 3.544$, $p = 0.026$), suggesting that race might influence depression rates in the HF patient population. However, the Black sample in this study was small (7%), and the researchers attributed the lower depression scores observed to their outpatient study setting. Six months or less had elapsed since the participants in their sample had last been hospitalized, which could have resulted in lower rates of depression. This finding is consistent with other studies who have found that hospitalized patients with HF are at risk for higher levels of depression (Albert et al., 2009; Freedland et al., 2003; Polikandrioti, 2019) compared to community-dwelling patients.

Multiple studies have found that Black individuals have less depression than Whites (Ahmed et al., 2006; Koenig, 1998; Stamp et al., 2014; Vaccarino, Kasl, et al., 2001; Wierenga et al., 2017). In a study of 391 inpatients at an academic medical center, Vaccarino, Kasl, et al. (2001) found that Black individuals ($n = 79$) were significantly less depressed than other races (mean = 6.81 ± 2.33 , $p = 0.02$). Ahmed et al. (2006) conducted a retrospective study of 2,720 participants (8.8% Black; $n = 240$) from all 50 states and the District of Columbia using the National Hospital Discharge Survey 2001–2003 data sets. The findings showed that among depressed patients in the randomized postmatched sample ($n = 680$), only 8.7% ($n = 59$) of Black individuals were depressed, as determined by ICD-9-CM coding. However, patients in the prematched sample who were depressed were less likely to be classified as Black ($p = 0.002$). Although Black race was not a significant predictor of depression in the overall post matched sample ($p = 0.876$), the depression prevalence among Black individuals accounted for 24.6% of the Black randomized, postmatched sample ($n = 240$; Ahmed et al., 2006).

Although Ahmed et al. (2006) provided insight into depression rates among Black individuals in large samples from various geographic locales, the measurement of depression could have confounded results. The findings might have varied if they used a screening tool with

demonstrated validity and reliability in measuring depression or different statistical analytic approaches. Using ICD-9-CM coding as a measure of depression limits the findings to depression officially diagnosed and documented by a health care provider. This limitation is problematic because depression is underdiagnosed and undertreated in the HF patient population (Ahmed et al., 2006; Bean et al., 2009; Celano et al., 2018; Dekker, Lennie, et al., 2014) and especially in Black individuals with HF (Bean et al., 2009; Dickson, McCarthy, & Katz, 2013; Evangelista et al., 2009). Relying on ICD-9/10 coding to identify depression likely results in the under identification of patients who are depressed. Additionally, depressive symptoms, although debilitating to patients (Dekker, Lennie, 2014; Dekker, Peden, et al., 2009; Havranek et al., 2004; Lesman-Leegte et al., 2009; Rumsfeld et al., 2003; Vaccarino, Kasl, et al., 2001), are not recognized as a diagnosis that can be captured based on ICD-9/10-CM coding.

Some studies have found no racial differences in depression among Black and White individuals with HF (Freedland et al., 2003; Gottlieb et al., 2004; Griffin et al., 2007; Piamjariyakul et al., 2013; Rohyans & Pressler, 2009; Xiong et al., 2015). For example, in a secondary analysis of data from the SADHART-CHF RCT (randomized controlled trial) that included 155 patients with HF recruited from an outpatient academic HF practice (39% Black; $n = 60$), Xiong et al. (2015) found no baseline racial differences in depression ($\chi^2 = 9.40, p = 0.15$). However, 48 Black patients in the sample (80%) were categorized as mildly ($n = 17$), moderately ($n = 24$), or severely ($n = 7$) depressed. In another study of 155 patients with HF (57% Black; $n = 89$), Gottlieb et al. (2004) found that 48% of the overall sample was depressed, and 53.3% of those depressed were Black ($n = 40$). However, the researchers identified no statistically significant differences in depression rates when comparing Black patients ($n = 89$) and White patients ($n = 65; p = 0.344$).

Among participants recruited from primary care and HF clinics ($N = 150$), Rohyans and Pressler (2009) found that 28% of the overall sample was depressed. However, there were no statistically significant differences in depression in Black individuals (31%; $n = 47$) compared to White individuals (67%; $n = 101$; $t = 1.06$, $p = 0.29$). Finally, in a secondary analysis of data from a randomized controlled trial conducted in a sample of patients with HF, investigators found that there were no baseline differences in depression in Blacks ($n = 77$) when compared to Whites ($n = 93$). However, after a nurse-led group discussion intervention, Black individuals had a statistically and clinically significant reduction in depression of 4.19 points compared to White individuals ($F = 3.99$, $p = 0.047$; Piamjariyakul et al., 2018). These findings suggest that Black individuals with HF who experience depression might benefit from tailored interventions. However, there should be more research regarding potential racial differences in the rates and nature of depression and relationships to anxiety, sociodemographic and clinical characteristics, and QOL in Black and White individuals before determining effective interventions.

Although depression among Black individuals with HF is underdiagnosed and undertreated (Dickson, McCarthy, & Katz, 2013; Lewis, 2011; Mentz et al., 2015), it is often more debilitating when compared to White patients (Mentz et al., 2015). For example, some researchers have found that Black patients with depression were more likely to have worse clinical outcomes, greater mortality, and greater risk for rehospitalization (Mentz et al., 2015). Depression is an undertreated condition in patients with HF, especially in Black patients (Sharma et al., 2009), and Black individuals are less likely to have an antidepressant prescription (Lewis et al., 2011). Although some depressed patients with HF receive antidepressant prescriptions, they remain depressed (Chung et al., 2013; Freedland, Hessler, et al., 2016; Jiang et al., 2001) either because the prescribed treatment is ineffective, or because these patients are not following

the prescribed treatment regimen. Sharma et al. (2009) studied a predominantly Black sample of patients with HF (86%; $N = 116$) recruited from three emergency departments in New York City. The researchers found a depression prevalence of 45% and concluded that depression in Black individuals is frequently untreated yet more debilitating than in White individuals. Dickson, McCarthy, and Katz (2013) conducted a mixed methods study with an all-Black sample of patients with HF ($N = 30$) recruited from a HF clinic and inpatient units at an urban medical center. The findings showed that 40% of the sample was depressed, but none were prescribed antidepressants.

Freedland et al. (2016) conducted a prospective cohort study at an urban academic medical center with a sample of 662 inpatients with HF (40% Black). They found that 39.6% ($n = 42$) of participants with minor depression and 42% ($n = 55$) with major depression were Black. Although these investigators found high rates of depression among Black individuals in their sample, major depression did not significantly influence survival in Black individuals with HF ($p = 0.14$), despite its observable influence in White individuals ($p < 0.0001$). Conversely, minor depression significantly influenced survival in Black individuals with HF ($p = 0.0003$) but did not have the same effect on White individuals ($p = 0.95$). These findings suggest that there might be racial differences and that lower levels of depression can have more pronounced effects on Black individuals with HF. Accordingly, depressive symptoms that do not meet the diagnostic criteria for clinical depression could be especially debilitating to Black individuals with HF and contribute to negative clinical outcomes and even persistent disparities. However, these relationships require further exploration to better understand racial differences in the influences on depression in Blacks compared to Whites and in whether the severity of depression influences Black patients differently than Whites.

Because Freedland et al. (2016) used the DSM-IV/DSM-5 criteria to measure depression, the observed depression rates excluded depressive symptoms and might have been underestimated. Although depressive symptoms result in adverse clinical outcomes for patients with HF (Celano et al., 2018; Freedland et al., 2011), the DSM-IV/DSM-5 is a source to diagnose major depression and not screen for depressive symptoms, leading to an underestimation of depression. Based on these findings, it seems as though depression in Black patients remains underdiagnosed and undertreated, yet it is debilitating to them and results in negative clinical outcomes. This trend might be among the reasons that disparities have persisted among Blacks with HF.

In addition to individual studies, investigators have conducted systematic reviews and meta-analyses on depression incidence among HF patients (Celano et al., 2018; Ishak et al., 2020; Johansson et al., 2006; Joynt et al., 2004; Konstam et al., 2005; MacMahon & Lip, 2002; Rutledge et al., 2006; Sokoreli et al., 2016), yet none reported findings based on race. Perhaps these investigators did not consider race an important construct to include in their analyses, or the sample sizes were too small to analyze (Rutledge et al., 2006). In addition, because few researchers have tested for racial differences across several studies or large enough samples, the literature lacks a reliable summary of potential racial differences in depression between Black and White individuals. This study yielded new knowledge through an investigation of potential differences in depression and anxiety in Black and White individuals with HF to determine whether there are racial differences.

Anxiety. Anxiety disorders include feelings of stress, nervousness, anxiousness, excessive fear, and anticipation of a future concern and incorporate muscle tension and avoidance behavior that impacts functioning (American Psychiatric Association, 2017). Anxiety

symptoms are similar to anxiety disorders but do not meet the diagnostic criteria for the diagnosis; however, both the symptoms and the disorders negatively affect patients with HF (Konstam et al., 2005; Lenze et al., 2000; Rechenberg et al., 2020). For this dissertation, “anxiety” referred to both anxiety that meets the diagnostic criteria and anxious symptoms that do not meet the criteria for generalized anxiety disorder yet are clinically significant and negatively impact patients with HF (see Pelle et al., 2008).

Although anxiety is common in patients with HF, it is significantly understudied among this patient population (Celano et al., 2018; Dekker et al., 2014; Konstam et al., 2005; MacMahon & Lip, 2002; Sokoreli et al., 2016). Few researchers have considered anxiety in Black individuals with HF or whether there are racial differences between Blacks and Whites (Evangelista et al., 2009). The findings from these limited studies have been inconsistent, showing that Black individuals had more anxiety or less anxiety or no statistically significant differences in anxiety between Black and White individuals with HF.

A few researchers have found that Blacks are more likely to be anxious (Dekker, Lennie, et al., 2014; Evangelista et al., 2009) compared to Whites. Dekker, Lennie, et al. (2014) recruited an outpatient sample ($N = 556$; 36% minorities) from three Southeastern and Midwestern academic medical centers and clinics. These investigators reported a 23% ($n = 130$) anxiety prevalence in their overall sample, which was 36% minority ($n = 200$) and mostly Black. The results showed a significantly higher proportion of anxious patients who were minorities (43% vs. 29%, $p < 0.001$). However, because the researchers categorized all minorities together, they did not indicate the specific races of participants and did not isolate Black participants in the analysis. Combining all minorities might have confounded results. Accordingly, readers cannot

determine the specific potential relationship between anxiety and Black individuals with HF and any potential racial differences compared to Whites.

Examining a sample of 241 patients with HF (60.7% White, 22.8% Hispanic, and 7% Black) recruited from an outpatient clinic, Evangelista et al. (2009) found that 40% of the overall sample was anxious. In a post hoc analysis, the investigators concluded that Black individuals ($n = 18$) were more likely to be anxious than Whites ($f = 3.107, p = 0.048$). A limitation of the study was the small Black sample.

Other evidence suggests that Black individuals with HF might have less anxiety than White individuals with HF. Bean et al. (2009) conducted a cross-sectional study with a sample of 97 patients with HF (48 Black [49.5%], 46 White [47.4%], and three Hispanic [3.1%]) recruited from the inpatient setting at a large, urban medical center and cardiac clinic. The results showed that only 8.2% ($n = 8$) of the sample was anxious. Bean et al. also found that 6.2% ($n = 3$) of Black participants and 10.2% ($n = 5$) of Whites were anxious, suggesting that Black individuals might have less anxiety than Whites. However, the overall study sample was small, yielding small racial groups for analysis. The researchers identified the need for more research focused on the Black population to determine whether existing knowledge is generalizable to Black individuals. In Cully et al.'s (2009) study of 158 patients with HF, the observed anxiety prevalence was 25.3%. However, White race was used as the referent group, resulting in only two racial categories and all non-White patients (37%; $n = 27$) categorized together.

Finally, some studies found no statistically significant differences in anxiety when comparing Black and White patients with HF. Evangelista et al. (2009) recruited a sample of 241 patients with HF (7% Black; $n = 18$) from an outpatient HF clinic. Although there was a trend when comparing anxiety in Black and White individuals with HF, these relationships were not

statistically significant ($f = 3.107, p = 0.290$). However, because the Black sample was small, it is unclear if this finding can be generalized to the HF patient population.

De Jong et al. (2011) found that 54% of their sample ($N = 147$) recruited from Midwestern outpatient academic medical centers and clinics had more than double the anxiety levels of healthy adults. Anxiety independently predicted event-free survival after adjusting for age, gender, NYHA Class, EF, ACE inhibitor use, and beta blocker use. Further, patients with greater anxiety levels were more likely to visit the emergency department, be hospitalized, and die. However, the Black patients accounted for 11% of the sample ($n = 16$), and no racial subgroup analysis occurred to assess for racial differences. In summary, research related to racial differences between Black and White patients with HF does not provide enough insight into these phenomena. Anxiety levels in patients with HF could contribute to adverse clinical outcomes and disparities. There is a need to better understand anxiety in Black patients with HF and whether there are racial differences. The proposed study will be an assessment of these relationships.

Comorbid Depression and Anxiety. Limited research suggests that depression and anxiety frequently co-occur (Dekker et al., 2014; Evangelista et al., 2009; Salyer et al., 2019). However, their comorbid occurrence and its influencing factors are unknown. Findings specific to Black individuals with HF are minimal, yet they could be significant contributors to the disparate clinical outcomes in this sub-patient population (Benjamin et al., 2017; Go et al., 2013; Roger et al., 2012; Virani et al., 2020, 2021). Dekker, Lennie, et al. (2014) found that depression and anxiety highly correlated in their sample of 556 patients with HF recruited from academic medical centers and clinics in the U.S. Southeast and Midwest, and that depression was a predictor of anxiety. The results showed that 32% of the sample had both depression and anxiety,

58% ($n = 179$) of the patients who were anxious ($n = 309$) had comorbid depression, and 91% of the patients who were depressed ($n = 179$) were anxious. Minorities comprised 36% ($n = 200$) of the sample, and Dekker et al. concluded that minorities were more likely to have comorbid depression and anxiety.

Because the researchers reported the findings for all minorities together, data specific to Black individuals with HF cannot be isolated. As a result, there is little known about comorbid depression and anxiety in patients with HF. Even less is known specifically about Black patients or whether racial differences in comorbid depression and anxiety exist when comparing Black and White patients. The impact of comorbid depression and anxiety in Black individuals with HF is unclear due to the lack of exploration of the phenomenon. Black individuals with HF might mirror the general HF patient population in that co-occurring depression and anxiety are common, underdiagnosed, and undertreated yet negatively impact clinical outcomes (Celano et al., 2018; Cully et al., 2009a, 2009b; Dekker, Lennie, et al., 2014; Evangelista, et al., 2009; MacMahon & Lip, 2002; Yohannes et al., 2010). A better understanding of comorbid depression and anxiety in patients with HF, the sociodemographic and clinical characteristics that might influence the phenomenon, and whether relationships differ by White race and Black race is needed. Studying the influence of race on anxiety in Black patients with HF is essential to determine the comorbid prevalence in this subpopulation and identify whether racial differences exist.

Depression and anxiety might be more difficult to diagnose in Black individuals with HF than in White individuals because of ethnic/sociocultural differences; reported rates might be inaccurate (Dickson, McCarthy, et al., 2013; Dickson, McCarthy, & Katz, 2013). Providers cannot rely on Black individuals to self-report depression and anxiety or request screening or

treatment because they are less likely to use mental health services than White individuals (Alegría et al., 2008; Cook et al., 2014; Das et al., 2006; Ojeda & McGuire, 2006). Black patients have stated that they seldom reported or admitted their feelings of depression and anxiety to health care providers (Dickson, McCarthy, & Katz, 2013; Gottlieb et al., 2004; Sharma et al., 2009). This lack of disclosure could be because Blacks might perceive depression and anxiety differently than other races. They might not use the term “depression” but consider themselves overwhelmed with sadness or lacking energy (Dickson, McCarthy, & Katz, 2013). Further, some Black individuals with HF might hold negative attitudes about depression and anxiety based on stigma. Withholding these subjective symptoms could signify a cultural difference that contributes to underdiagnosis and undertreatment among Black individuals with HF (Das et al., 2006).

Depression and anxiety could vary by race (Dickson, McCarthy, et al., 2013), but limited knowledge regarding this phenomenon exists. Yet, differences in clinical outcomes between Black and White individuals with HF suggest the need to explore possible racial differences. Several methodological issues, including measurement and sampling, might influence or confound existing findings related to depression and anxiety in Black and White individuals with HF. To better understand whether racial differences in depression and anxiety are present among Black and White individuals with HF, more research is needed that considers race as an important construct with the potential to influence outcome variables (Doll, 2018). This study was among the few investigations of Black/White racial differences in depression and anxiety among patients with HF.

Quality of Life. QOL is a subjective, multidimensional concept encompassing various aspects of well-being and incorporating individual preferences, experiences, and perspectives

(Cella, 1994; Felce & Perry, 1995; Heo et al., 2009). In this study, QOL was characterized as life satisfaction (Ferrans & Powers, 1985) and defined as individuals' sense of well-being based on their satisfaction or dissatisfaction with the aspects of life they deem most important (Ferrans, 1990). QOL in this context is holistic, including various domains, the subjective evaluation of each domain, and the unique level of importance assigned to each domain by the individual (Ferrans & Powers, 1985). The domains contribute to four major dimensions of QOL: health and functioning, psychological/spiritual, social and economic, and family (Ferrans, 1990; Ferrans & Powers, 1985).

Depression and anxiety in chronically ill patients, particularly when undiagnosed and untreated, contribute to decreased life satisfaction, reduced QOL (Lim et al., 2011), including poor health-related quality of life (Balcells et al., 2010; Brown et al., 2010; Chuquilín-Arista et al., 2021; Lim et al., 2011; Liu et al., 2020; Sadlonova et al., 2021; Strine et al., 2008; von Leupoldt et al., 2011) and negative clinical outcomes (Assari, 2014; McLaughlin et al., 2005; Ryu et al., 2016; Sidell, 1997; Whittemore & Dixon, 2008), including greater morbidity, health care utilization (Ani et al., 2009), and mortality (McLaughlin et al., 2005; Sadlonova et al., 2021). Chronically ill Black individuals have worse QOL than their White counterparts (Jackson-Triche et al., 2000).

The chronicity of HF and the nature, intensity, and duration of symptoms, such as the physical and psychosocial symptoms experienced by patients with HF (Alemoush et al., 2021; Bekelman et al., 2007; Hwang et al., 2014; Rechenberg et al., 2020; Salyer et al., 2019), can negatively impact QOL (Bekelman et al., 2007; Cella, 1994; Salyer et al., 2019). Limited evidence suggests that QOL in Black individuals can differ from that observed within the general HF patient population (Warnecke et al., 1996). Although these differences are not well

understood, depression (AbuRuz, 2018; Aggelopoulou et al., 2017; Bekelman et al., 2007; Chung, Moser, Lennie, & Rayens, 2009; Chung, Moser, Lennie, & Frazier, 2013; Cully et al., 2010; Gottlieb et al., 2004; Mentz et al., 2015; Salyer et al., 2019b; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009) and anxiety (AbuRuz, 2018; Aggelopoulou et al., 2017; Chung, Moser, Lennie, & Rayens, 2009; Polikandrioti et al., 2019; Salyer et al., 2019) can negatively impact QOL in patients with HF and might impact Black individuals differently than Whites.

In a study of 695 patients with HF of whom 227 were Black (33%), de Leon et al. (2009) found that greater depression was associated with lower QOL (-0.082 , $SE = 0.0005$, $p < 0.001$) and that Black individuals had better QOL than White individuals (-0.550 , $SE = 0.148$, $p < 0.05$). However, the Black patients in de Leon et al.'s sample reported worse QOL based on their self-perceptions, which differed from the scores derived from the measurement. These findings suggested a possible relationship between depression and QOL and that published findings related to QOL might vary based on conceptualization and measurement. Sampling errors could have influenced the findings, as the researchers excluded 209 patients due to missing data. The excluded patients were more likely to be Black and have lower QOL scores.

Bekelman et al. (2007) found a 32% rate of depression in their sample ($N = 60$), with Black individuals significantly more likely to have worse QOL than White participants ($p = .04$). However, the overall sample size was small, there were only seven (11.7%) Black participants, and the researchers did not stratify depression rates by race. In another study of patients with HF ($N = 97$; 49.5% Black) recruited from both inpatient and outpatient settings at a large medical center, Bean et al. (2009) found that participants with more depression and anxiety reported lower QOL. The researchers observed no statistically significant racial differences ($p > .05$). The researchers might not have found racial differences because of the small sample size ($N = 97$)

and varied sample characteristics, including mean age and disease severity. Bean et al. recommended more research to stratify these phenomena by race.

Despite knowledge that depression (Eisele et al., 2021; Garin et al., 2009; Rustad et al., 2013; Samartzis et al., 2013) and anxiety (Eisele et al., 2021) negatively impact QOL in patients with HF, more research is needed. There is insufficient evidence regarding racial differences between Black individuals compared to White individuals, especially regarding the potential influence of anxiety. Perhaps the lack is due to varying research methods, including sampling and measurement approaches. In this study, depression, anxiety, and QOL were stratified by race in a larger sample to determine whether there are racial differences. Using a valid and reliable research instrument designed to measure QOL in patients with cardiovascular illnesses was a means to improve measurement accuracy. Studies focused on the influence of anxiety on QOL are lacking, and most researchers have not measured both depression and anxiety, despite the conditions' high correlation necessitating joint study in patients with HF (Celano et al., 2018; Chung, Moser, Lennie, & Rayens, 2009; Dekker, Lennie, et al., 2014; Herr et al., 2013; Konstam et al., 2005; Lenze et al., 2000; MacMahon & Lip, 2002). Measuring depression and anxiety simultaneously in this study will add to the existing body of knowledge.

Sociodemographic Characteristics and Quality of Life. Findings regarding the influence of sociodemographic characteristics (race, age, gender, marital status, education, and employment status) on QOL vary. Younger age (Carson et al., 2009; Liu et al., 2021; Reddy et al., 2020) and female gender (Carson et al., 2009; Chana et al., 2019) have been associated with better QOL in some patients with HF. Although most findings have not been stratified by race, in an all-Black study of patients with HF ($N = 1,050$), Carson et al. (2009) found associations between worse QOL and younger age and overall QOL with all-cause mortality. de Leon et al.

(2009) noted that Black participants had lower QOL-physical functioning. Similarly, older age was correlated with poor QOL in a sample of patients from Greece ($N = 231$), but race was not reported (Aggelopoulou et al., 2017). Riegel et al. (2012) found relationships between age, gender, and QOL. They reported that older patients had better QOL, and women had worse QOL than men. However, a third group of investigators noted that age, gender, and marital status did not significantly influence QOL in patients with HF (Luttik et al., 2006). Marital status emerged as a significant predictor of QOL for Tarekegn et al. (2021; $p = 0.022$). However, the researchers conducted the study in Ethiopia and did not report race, perhaps because the sample was racially homogeneous.

Some investigators have found better QOL associated with higher educational levels (Liu et al., 2021), and patients with HF with lower educational status have lower QOL (Barbareschi et al., 2011; Lee et al., 2005). However, these studies occurred in Taiwan, Greece, the Netherlands, and Hong Kong, and race was not reported. AbuRuz (2018) concluded that employed patients with HF had better QOL. The study took place in Jordan, perhaps in a racially heterogeneous sample, and race was not reported. Although there is not much research published to establish the relationship between employment status and QOL, investigators have found that unemployment and poor economic situation are associated with worse QOL (Aggelopoulou et al., 2017). In comparison, economic independence and higher levels of education are associated with greater earning potential, contribute to better financial stability and independence, and are thereby associated with better QOL (Liu et al., 2021). Because low SES is associated with lower QOL (Luttik et al., 2006), education and employment status might influence QOL independently, and there could be differences when comparing Black and White patients with HF. It appears some sociodemographic characteristics might influence QOL. Still, more research is needed,

with an emphasis on identifying whether there are racial differences between Black and White patients with HF.

Clinical Characteristics and Quality of Life. Some investigators have found relationships between clinical characteristics (disease severity, comorbidities, and functional status) and QOL. Multiple studies have shown an association between higher disease severity and lower QOL (AbuRuz, 2018; Carson et al., 2009; Freedland et al., 2021). AbuRuz (2018) found higher EF associated with better QOL, whereas Coelho et al. (2005) found no clear association between EF and QOL. Similarly, a higher NYHA Class was associated with lower QOL in several studies (Coelho et al., 2005; Freedland et al., 2021; Lee et al., 2005). Comorbidities negatively influence QOL in patients with HF (Carson et al., 2009; Freedland et al., 2021; Heo et al., 2012; Reddy et al., 2020), and lower comorbidity burden was associated with better QOL ($r = .365, p < .001$; Heo et al., 2012). Carson et al. (2009) found that some comorbidities (chronic obstructive pulmonary disease and systolic hypotension) were associated with lower QOL in an all-Black sample of patients with HF ($N = 1,050$) derived via stratified sampling from a randomized controlled trial. De Leon et al. (2009) noted that patients with higher symptom burden and lower functional capacity had worse QOL. In a secondary analysis of 408 patients with HF, Reddy et al. (2020) found that comorbidities were associated with worse QOL. However, the researchers did not report race.

Functional status influences various aspects of QOL in patients with HF. Wu et al. (2016) found that functional status was associated with QOL, and patients with better functional status had better QOL ($p < 0.001$). However, most of the sample was White (79%), all minorities were grouped, and no findings were stratified by Black race. Conversely, some researchers found a weak association between functional impairment and QOL (Carels, 2004). De Leon et al. (2009)

found that younger age and Black race were associated with worse QOL-physical functioning, but older age was associated with higher overall QOL scores. This relationship suggests that although patients in the study had lower function, they had greater satisfaction. Despite these findings, Black individuals with HF might experience greater functional impairment (Vacarino, Gahbauer, et al., 2002) and are at risk for lower QOL. This association merits exploration to determine whether racial differences exist.

Summary/Conclusion

Limited research suggests there could be racial differences in the rates of depression and anxiety among Black individuals with HF (Dickson et al., 2013), which is consistent with findings in other chronically ill patient populations. Depression and anxiety in Black individuals with HF are grossly understudied, underdiagnosed, and undertreated, even more than in the general HF patient population. In addition, comorbid depression and anxiety remain largely unstudied, especially in Black individuals. Despite some progress in better understanding psychosocial and psychological functioning in patients with HF, few researchers have focused on anxiety and comorbid depression and anxiety. There has been little progress made in testing whether racial differences are present and whether the existing knowledge can be generalized to Blacks.

Most researchers of depression and anxiety in patients with HF have not included representative Black samples or conducted racial subgroup analyses, even when recruiting adequate Black samples (Bean et al., 2009). As a result, little is known regarding potential differences in the sociodemographic and clinical influences of depression and anxiety on Black individuals with HF, how they might influence QOL, and whether there are racial differences between Black and White patients. Applying and generalizing the limited research to Blacks is

presumptuous because the existing knowledge focuses on depression and anxiety across the entire population of patients with HF without adequately addressing race and potential racial differences. Black patients with HF who experience depression and anxiety might have unique needs and experiences compared to their White counterparts. Providing better care for this sub-patient population requires a better understanding of the rates of depression and anxiety and their comorbid occurrence in Black individuals with HF, the sociodemographic and clinical characteristics associated with depression and anxiety, and how these characteristics influence QOL in Black and White patients with HF.

References

- AbuRuz, M. E. (2018). Anxiety and depression predicted quality of life among patients with heart failure. *Journal of Multidisciplinary Healthcare, 11*, 367–373.
<https://doi.org/10.2147/JMDH.S170327>
- Adelborg, K., Schmidt, M., Sundbøll, J., Pedersen, L., Videbech, P., Bøtker, H. E., Egstrup, K., & Sørensen, H. T. (2016). Mortality risk among heart failure patients with depression: A nationwide population-based cohort study. *Journal of the American Heart Association, 5*(9), Article e004137. <https://doi.org/10.1161/JAHA.116.004137>
- Afzal, A., Ananthasubramaniam, K., Sharma, N., Ai-Malki, Q., Ali, A. S., Jacobsen, G., & Jafri, S. M. (1999). Racial differences in patients with heart failure. *Clinical Cardiology, 22*(12), 791–794. <https://doi.org/10.1002/clc.4960221207>
- Aggelopoulou, Z., Fotos, N. V., Chatziefstratiou, A. A., Giakoumidakis, K., Elefsiniotis, I., & Brokalaki, H. (2017). The level of anxiety, depression and quality of life among patients with heart failure in Greece. *Applied Nursing Research, 34*, 52–56.
<https://doi.org/10.1016/j.apnr.2017.01.003>
- Ahmed, A., Ali, M., Lefante, C. M., Mullick, M. S., & Kinney, F. C. (2006). Geriatric heart failure, depression, and nursing home admission: An observational study using propensity score analysis. *The American Journal of Geriatric Psychiatry, 14*(10), 867–875. <https://doi.org/10.1097/01.jgp.0000209639.30899.72>
- Ahn, S., Kim, S., & Zhang, H. (2017). Changes in depressive symptoms among older adults with multiple chronic conditions: Role of positive and negative social support. *International Journal of Environmental Research and Public Health, 14*(1), Article 16.
<https://doi.org/10.3390/ijerph14010016>

- Akwo, E. A., Kabagambe, E. K., Harrell, J. F. E., Blot, W. J., Bachmann, J. M., Wang, T. J., Gupta, D. K., & Lipworth, L. (2018). Neighborhood deprivation predicts heart failure risk in a low-income population of Blacks and Whites in the Southeastern United States. *Circulation: Cardiovascular Quality and Outcomes*, *11*(1), Article e004052. <https://doi.org/10.1161/CIRCOUTCOMES.117.004052>
- Albert, N. M., Fonarow, G. C., Abraham, W. T., Gheorghiade, M., Greenberg, B. H., Nunez, E., O'Connor, C. M., Stough, W. G., Yancy, C. W., & Young, J. B. (2009). Depression and clinical outcomes in heart failure: an OPTIMIZE-HF analysis. *The American Journal of Medicine*, *122*(4), 366–373. <https://doi.org/10.1016/j.amjmed.2008.09.046>
- Alegría, M., Chatterji, P., Wells, K., Cao, Z., Chen, C.-N., Takeuchi, D., Jackson, J., & Meng, X.-L. (2008). Disparity in depression treatment among racial and ethnic minority populations in the United States. *Psychiatric Services*, *59*(11), 1264–1272. <https://doi.org/10.1176/ps.2008.59.11.1264>
- Alemoush, R. A., Al-Dweik, G., & AbuRuz, M. E. (2021). The effect of persistent anxiety and depressive symptoms on quality of life among patients with heart failure. *Applied Nursing Research*, *62*, Article 151503. <https://doi.org/10.1016/j.apnr.2021.151503>
- Alhurani, A. S., Dekker, R. L., Abed, M. A., Khalil, A., Al Zaghal, M. H., Lee, K. S., Mudd-Martin, G., Biddle, M. J., Lennie, T. A., & Moser, D. K. (2015). The association of comorbid symptoms of depression and anxiety with all-cause mortality and cardiac rehospitalization in patients with heart failure. *Psychosomatics*, *56*(4), 371–380. <https://doi.org/10.1016/j.psych.2014.05.022>

- Almawi, W. Y., Tamim, H., Al-Sayed, N., Arekat, M. R., Al-Khateeb, G. M., Baqer, A., Tutanji, H., & Kamel, C. (2014). Association of comorbid depression, anxiety, and stress disorders with type 2 diabetes in Bahrain, a country with a very high prevalence of type 2 diabetes. *Journal of Endocrinological Investigation*, *31*(11), 1020–1024.
<https://doi.org/10.1007/BF03345642>
- American Heart Association. (2017). *Classes of heart failure*. <https://www.heart.org/en/health-topics/heart-failure/what-is-heart-failure/classes-of-heart-failure>
- American Psychiatric Association. (2017). *What are anxiety disorders?*
<https://www.psychiatry.org/patients-families/anxiety-disorders/what-are-anxiety-disorders>
- American Psychiatric Association. (2020). *What is depression?*
<https://www.psychiatry.org/patients-families/depression/what-is-depression>
- American Psychological Association. (2022). *Dictionary of psychology*.
<https://dictionary.apa.org/educational-attainment-level>
- Anderson, N. B., & Armstead, C. A. (1995). Toward understanding the association of socioeconomic-status and health: A new challenge for the biopsychosocial approach. *Psychosomatic Medicine*, *57*(3), 213–225. <https://doi.org/10.1097/00006842-199505000-00003>
- Andrus, M. R., & Roth, M. T. (2002). Health literacy: A review. *Pharmacotherapy*, *22*(3), 282–302. <https://doi.org/10.1592/phco.22.5.282.33191>

Ani, C., Bazargan, M., Hindman, D., Bell, D., Rodriguez, M., & Baker, R. S. (2009). Comorbid chronic illness and the diagnosis and treatment of depression in safety net primary care settings. *Journal of the American Board of Family Medicine*, 22(2), 123–135.

<https://doi.org/10.3122/jabfm.2009.02.080035>

Assari, S. (2014). Chronic medical conditions and major depressive disorder: Differential role of positive religious coping among African Americans, Caribbean Blacks and non-Hispanic Whites. *International Journal of Preventive Medicine*, 5(4), 405–413.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4018588/pdf/IJPVM-5-405.pdf>

Assari, S. (2016). Race and ethnic differences in additive and multiplicative effects of depression and anxiety on cardiovascular risk. *International Journal of Preventive Medicine*, 7(1),

Article 22. <https://doi.org/10.4103/2008-7802.173931>

Assari, S. (2019). Race, depression, and financial distress in a nationally representative sample of American adults. *Brain Sciences*, 9(2), Article 29.

<https://doi.org/10.3390/brainsci9020029>

Assari, S., Burgard, S., & Zivin, K. (2015). Long-term reciprocal associations between depressive symptoms and number of chronic medical conditions: Longitudinal support for Black–White health paradox. *Journal of Racial and Ethnic Health Disparities*, 2(4),

589–597. <https://doi.org/10.1007/s40615-015-0116-9>

Assari, S., Dejman, M., & Neighbors, H. W. (2015). Ethnic differences in separate and additive effects of anxiety and depression on self-rated mental health among Blacks. *Journal of*

Racial and Ethnic Health Disparities, 3(3), 423–430. <https://doi.org/10.1007/s40615->

[015-0154-3](https://doi.org/10.1007/s40615-015-0154-3)

- Athilingam, P., Jenkins, B., Johansson, M., & Labrador, M. A. (2017). Mobile health intervention to improve self-care in patients with heart failure: Pilot randomized control trial. *JMIR Cardio*, *1*(2), Article e3. <https://doi.org/10.2196/cardio.7848>
- Averbuch, T., Mohamed, M. O., Islam, S., Defilippis, E. M., Breathett, K., Alkhouli, M. A., Michos, E. D., Martin, G. P., Kontopantelis, E., Mamas, M. A., & Van Spall, H. G. C. (2021). The association between socioeconomic status, sex, race/ethnicity and in-hospital mortality among patients hospitalized for heart failure. *Journal of Cardiac Failure*, *28*(5), 697–709. <https://doi.org/10.1016/j.cardfail.2021.09.012>
- Azam, T. U., & Colvin, M. M. (2021). Representation of Black patients in heart failure clinical trials. *Current Opinion in Cardiology*, *36*(3), 329–334. <https://doi.org/10.1097/HCO.0000000000000849>
- Balcells, E., Gea, J., Ferrer, J., Serra, I., Orozco-Levi, M., de Batlle, J., Rodriguez, E., Benet, M., Donaire-Gonzalez, D., Anto, J. M., & Garcia-Aymerich, J. (2010). Factors affecting the relationship between psychological status and quality of life in COPD patients. *Health Qual Life Outcomes*, *8*(1), 108–108. <https://doi.org/10.1186/1477-7525-8-108>
- Barbareschi, G., Sanderman, R., Lesman-Leegte, I., Van Veldhuisen, D. J., & Jaarsma, T. (2011). Educational level and the quality of life of heart failure patients: A longitudinal study. *Journal of Cardiac Failure*, *17*(1), 47–53. <https://doi.org/10.1016/j.cardfail.2010.08.005>
- Bean, M. K., Gibson, D., Flattery, M., Duncan, A., & Hess, M. (2009). Psychosocial factors, quality of life, and psychological distress: Ethnic differences in patients with heart failure. *Progress in Cardiovascular Nursing*, *24*(4), 131–140. <https://doi.org/10.1111/j.1751-7117.2009.00051.x>

- Beatty, & Joffe, R. (2006). An overlooked dimension of diversity: The career effects of chronic illness. *Organizational Dynamics*, 35(2), 182–195.
<https://doi.org/10.1016/j.orgdyn.2006.03.006>
- Bekelman, D. B., Havranek, E. P., Becker, D. M., Kutner, J. S., Peterson, P. N., Wittstein, I. S., Gottlieb, S. H., Yamashita, T. E., Fairclough, D. L., & Dy, S. M. (2007). Symptoms, depression, and quality of life in patients with heart failure. *Journal of Cardiac Failure*, 13(8), 643–648. <https://doi.org/10.1016/j.cardfail.2007.05.005>
- Bell, R. A., Smith, S. L., Arcury, T. A., Snively, B. M., Stafford, J. M., & Quandt, S. A. (2005). Prevalence and correlates of depressive symptoms among rural older African Americans, Native Americans, and Whites with diabetes. *Diabetes Care*, 28(4), 823–829.
<https://doi.org/10.2337/diacare.28.4.823>
- Benatar, D., Bondmass, M., Ghitelman, J., & Avitall, B. (2003). Outcomes of chronic heart failure. *Archives of Internal Medicine*, 163(3), 347–352.
<https://doi.org/10.1001/archinte.163.3.347>
- Benjamin, J. E., Blaha, J. M., Chiuve, E. S., Cushman, R. M., Das, D. S., Deo, R. R., De Ferranti, C. S., Floyd, C. J., Fornage, E. M., Gillespie, H. C., Isasi, T. C., Jiménez, H. M., Jordan, E. L., Judd, W. S., Lackland, K. D., Lichtman, R. J., Lisabeth, J. L., Liu, J. S., Longenecker, A. C., ... Muntner, S. P. (2017). Heart disease and stroke statistics—2017 update: A report from the American Heart Association. *Circulation*, 135(10), e146–e603.
<https://doi.org/10.1161/CIR.0000000000000485>
- Berkman, N. D., Sheridan, S. L., Donahue, K. E., Halpern, D. J., & Crotty, K. (2011). Low health literacy and health outcomes: An updated systematic review. *Annals of Internal Medicine*, 155(2), 97–107. <https://doi.org/10.7326/0003-4819-155-2-201107190-00005>

- Beydoun, M. A., Beydoun, H. A., Mode, N., Dore, G. A., Canas, J. A., Eid, S. M., & Zonderman, A. B. (2016). Racial disparities in adult all-cause and cause-specific mortality among us adults: Mediating and moderating factors. *BMC Public Health*, *16*(1), Article 1113. <https://doi.org/10.1186/s12889-016-3744-z>
- Bhatt, K. N., Kalogeropoulos, A. P., Dunbar, S. B., Butler, J., & Georgiopolou, V. V. (2016). Depression in heart failure: Can PHQ-9 help? *International Journal of Cardiology*, *221*, 246–250. <https://doi.org/10.1016/j.ijcard.2016.07.057>
- Bhopal, R., Gruer, L., Agyemang, C., Davidovitch, N., de-Graft Aikins, A., Krasnik, A., Martinez-Donate, A. P., Miranda, J. J., Pottie, K., Segal, U., Zwi, A., & Kumar, B. (2021). The global society on migration, ethnicity, race and health: Why race can't be ignored even if it causes discomfort. *European Journal of Public Health*, *31*(1), 3–4. <https://doi.org/10.1093/eurpub/ckaa191>
- Bjelland, I., Krokstad, S., Mykletun, A., Dahl, A. A., Tell, G. S., & Tambs, K. (2008). Does a higher educational level protect against anxiety and depression? The HUNT study. *Social*
- Blinderman, Homel, P., Billings, J. A., Portenoy, R. K., & Tennstedt, S. L. (2008). Symptom Distress and Quality of Life in Patients with Advanced Congestive Heart Failure. *Journal of Pain and Symptom Management*, *35*(6), 594–603. <https://doi.org/10.1016/j.jpainsymman.2007.06.007>
- Boersma, P., Black, L. I., & Ward, B. W. (2020). Prevalence of multiple chronic conditions among US adults, 2018. *Preventing Chronic Disease*, *17*, Article e106. <https://doi.org/10.5888/pcd17.200130>
- Braunwald, E. (2013). Research advances in heart failure: a compendium. *Circulation Research*, *113*(6), 633–645. <https://doi.org/10.1161/CIRCRESAHA.113.302254>

- Brouwers, C., Kupper, N., Pelle, A. J., Szabó, B. M., Westerhuis, B. L., & Denollet, J. (2014). Depressive symptoms in outpatients with heart failure: Importance of inflammatory biomarkers, disease severity and personality. *Psychology & Health, 29*(5), 564–582. <https://doi.org/10.1080/08870446.2013.869813>
- Brown, L. F., Kroenke, K., Theobald, D. E., Wu, J., & Tu, W. (2010). The association of depression and anxiety with health-related quality of life in cancer patients with depression and/or pain. *Psycho-Oncology, 19*(7), 734–741. <https://doi.org/10.1002/pon.1627>
- Brunello, G., Fort, M., Schneeweis, N., & Winter-Ebmer, R. (2016). The causal effect of education on health: What is the role of health behaviors? *Health Economics, 25*(3), 314–336. <https://doi.org/10.1002/hec.3141>
- Cajita, M. I., Cajita, T. R., & Han, H.-R. (2016). Health literacy and heart failure: A systematic review. *The Journal of Cardiovascular Nursing, 31*(2), 121–130. <https://doi.org/10.1097/JCN.0000000000000229>
- Caraballo, C., Desai, N. R., Mulder, H., Alhanti, B., Wilson, F. P., Fiuzat, M., Felker, G. M., Piña, I. L., O'Connor, C. M., Lindenfeld, J., Januzzi, J. L., Cohen, L. S., & Ahmad, T. (2019). Clinical implications of the New York Heart Association Classification. *Journal of the American Heart Association, 8*(23), Article e014240. <https://doi.org/10.1161/JAHA.119.014240>
- Carels, R. A. (2004). The association between disease severity, functional status, depression and daily quality of life in congestive heart failure patients. *Quality of Life Research, 13*(1), 63–72. <https://doi.org/10.1023/B:QURE.0000015301.58054.51>

- Carson, P., Tam, S. W., Ghali, J. K., Archambault, W. T., Taylor, A., Cohn, J. N., Braman, V. M., Worcel, M., & Anand, I. S. (2009). Relationship of quality of life scores with baseline characteristics and outcomes in the African-American Heart Failure Trial. *Journal of Cardiac Failure, 15*(10), 835–842. <https://doi.org/10.1016/j.cardfail.2009.05.016>
- Celano, C. M., Villegas, A. C., Albanese, A. M., Gaggin, H. K., & Huffman, J. C. (2018). Depression and anxiety in heart failure: A review. *Harvard Review of Psychiatry, 26*(4), 175–184. <https://doi.org/10.1097/HRP.000000000000162>
- Celik, E., Cay, S., Sensoy, B., Murat, S., Oksuz, F., Cankurt, T., & Mendi, M. A. (2016). Heart failure functional class associated with depression severity but not anxiety severity. *Acta Cardiologica Sinica, 32*(1), 55–61. <https://doi.org/10.6515/ACS20150509A>
- Cella, D. F. (1994). Quality of life: Concepts and definition. *Journal of Pain and Symptom Management, 9*(3), 186–192. [https://doi.org/10.1016/0885-3924\(94\)90129-5](https://doi.org/10.1016/0885-3924(94)90129-5)
- Chana, A., Vaduganathan, M., Lewis, E. F., Claggett, B. L., Rizkala, A. R., Wang, W., Lefkowitz, M. P., Shi, V. C., Anand, I. S., Ge, J., Lam, C. S. P., Maggioni, A. P., Martinez, F., Packer, M., Pfeffer, M. A., Pieske, B., Redfield, M. M., Rouleau, J. L., van Veldhuisen, D. J., ... Solomon, S. D. (2019). Health-related quality of life in heart failure with preserved ejection fraction: The PARAGON-HF trial. *JACC: Heart Failure, 7*(10), 862–874.
- Charles, S. T., & Piazza, J. R. (2009). Age differences in affective well-being: Context matters. *Social and Personality Psychology Compass, 3*(5), 711–724. <https://doi.org/10.1111/j.1751-9004.2009.00202.x>

- Charlson, M. E., Pompei, P., Ales, K. L., & MacKenzie, C. R. (1987). A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *Journal of Chronic Diseases, 40*(5), 373–383. [https://doi.org/10.1016/0021-9681\(87\)90171-8](https://doi.org/10.1016/0021-9681(87)90171-8)
- Chaudhry, S. I., Herrin, J., Phillips, C., Butler, J., Mukerjee, S., Murillo, J., Onwuanyi, A., Seto, T. B., Spertus, J., & Krumholz, H. M. (2011). Racial disparities in health literacy and access to care among patients with heart failure. *Journal of Cardiac Failure, 17*(2), 122–127. <https://doi.org/10.1016/j.cardfail.2010.09.016>
- Chung, M. L., Lennie, T. A., Riegel, B., Wu, J.-R., Dekker, R. L., & Moser, D. K. (2009). Marital status as an independent predictor of event-free survival of patients with heart failure. *American Journal of Critical Care, 18*(6), 562–570. <https://doi.org/10.4037/ajcc2009388>
- Chung, M. L., Moser, D. K., Lennie, T. A., & Frazier, S. K. (2013). Perceived social support predicted quality of life in patients with heart failure, but the effect is mediated by depressive symptoms. *Quality of Life Research, 22*(7), 1555–1563. <https://doi.org/10.1007/s11136-012-0294-4>
- Chung, M. L., Moser, D. K., Lennie, T. A., & Rayens, M. K. (2009). The effects of depressive symptoms and anxiety on quality of life in patients with heart failure and their spouses: Testing dyadic dynamics using actor-partner interdependence model. *Journal of Psychosomatic Research, 67*(1), 29–35. <https://doi.org/10.1016/j.jpsychores.2009.01.009>

- Chuquilín-Arista, F., Álvarez-Avellón, T., & Menéndez-González, M. (2021). Impact of depression and anxiety on dimensions of health-related quality of life in subjects with Parkinson's disease enrolled in an association of patients. *Brain Sciences, 11*(6), Article 771. <https://doi.org/10.3390/brainsci11060771>
- Cirelli, M. A., Lacerda, M. S., Lopes, C. T., de Lima Lopes, J., & de Barros, A. L. B. L. (2018). Correlations between stress, anxiety and depression and sociodemographic and clinical characteristics among outpatients with heart failure. *Archives of Psychiatric Nursing, 32*(2), 235–241. <https://doi.org/10.1016/j.apnu.2017.11.008>
- Clarke, S. P., Frasure-Smith, N., Lespérance, F., & Bourassa, M. G. (2000). Psychosocial factors as predictors of functional status at 1 year in patients with left ventricular dysfunction. *Research in Nursing & Health, 23*(4), 290–300. [https://doi.org/10.1002/1098-240X\(200008\)23:4<290::AID-NUR5>3.0.CO;2-0](https://doi.org/10.1002/1098-240X(200008)23:4<290::AID-NUR5>3.0.CO;2-0)
- Cleland, J. A., Lee, A. J., & Hall, S. (2007). Associations of depression and anxiety with gender, age, health-related quality of life and symptoms in primary care COPD patients. *Family Practice, 24*(3), 217–223. <https://doi.org/10.1093/fampra/cmm009>
- Coelho, R., Ramos, S., Prata, J., Bettencourt, P., Ferreira, A., & Cerqueira-Gomes, M. (2005). Heart failure and health related quality of life. *Clinical Practice and Epidemiology in Mental Health, 1*(1), Article 19. <https://doi.org/10.1186/1745-0179-1-19>
- Cook, B. L., Zuvekas, S. H., Carson, N., Wayne, G. F., Vesper, A., & McGuire, T. G. (2014). Assessing racial/ethnic disparities in treatment across episodes of mental health care. *Health Services Research, 49*(1), 206–229. <https://doi.org/10.1111/1475-6773.12095>

- Corbie-Smith, G. M. D. M., Henderson, G. P., Blumenthal, C. M. P. H., Dorrance, J., & Estroff, S. P. (2008). Conceptualizing race in research. *Journal of the National Medical Association, 100*(10), 1235–1243. [https://doi.org/10.1016/S0027-9684\(15\)31470-X](https://doi.org/10.1016/S0027-9684(15)31470-X)
- Culley, L. (2006). Transcending transculturalism? Race, ethnicity and health-care. *Nursing Inquiry, 13*(2), 144–153. <https://doi.org/10.1111/j.1440-1800.2006.00311.x>
- Cully, J. A., Jimenez, D. E., Ledoux, T. A., & Deswal, A. (2009). Recognition and treatment of depression and anxiety symptoms in heart failure. *Primary Care Companion to the Journal of Clinical Psychiatry, 11*(3), 103–109. <https://doi.org/10.4088/PCC.08m00700>
- Cully, J. A., Johnson, M., Moffett, M. L., Khan, M., & Deswal, A. (2009). Depression and anxiety in ambulatory patients with heart failure. *Psychosomatics, 50*(6), 592–598. <https://doi.org/10.1176/appi.psy.50.6.592>
- Cully, J. A., Phillips, L. L., Kunik, M. E., Stanley, M. A., & Deswal, A. (2010). Predicting quality of life in veterans with heart failure: The role of disease severity, depression, and comorbid anxiety. *Behavioral Medicine, 36*(2), 70–76. <https://doi.org/10.1080/08964280903521297>
- Cuyjet, A. B., & Akinboboye, O. (2014). Acute heart failure (AHF) in the African American patient. *Journal of Cardiac Failure, 20*(7), 533–540.
- Das, A. K., Olfson, M., McCurtis, H. L., & Weissman, M. M. (2006). Depression in African Americans: Breaking barriers to detection and treatment. *Journal of Family Practice, 55*(1), 30–39. <https://doi.org/10.1016/j.cardfail.2014.04.018>
- de Groot, V., Beckerman, H., Lankhorst, G. J., & Bouter, L. M. (2003). How to measure comorbidity: A critical review of available methods. *Journal of Clinical Epidemiology, 56*(3), 221–229. [https://doi.org/10.1016/S0895-4356\(02\)00585-1](https://doi.org/10.1016/S0895-4356(02)00585-1)

- De Jong, M. J., Chung, M. L., Wu, J. R., Riegel, B., Rayens, M. K., & Moser, D. K. (2011). Linkages between anxiety and outcomes in heart failure. *Heart Lung, 40*(5), 393–404.
<https://doi.org/10.1016/j.hrtlng.2011.02.002>
- de Leon, C. F., Grady, K. L., Eaton, C., Rucker-Whitaker, C., Janssen, I., Calvin, J., & Powell, L. H. (2009). Quality of life in a diverse population of patients with heart failure: Baseline findings from the Heart Failure Adherence and Retention Trial (HART). *Journal of Cardiopulmonary Rehabilitation and Prevention, 29*(3), 171–178.
<https://doi.org/10.1097/HCR.0b013e31819a0266>
- Deek, H., Itani, L., & Davidson, P. M. (2020). Literacy critical to heart failure management: A scoping review. *Heart Failure Review, 26*(6), 1413–1419.
<https://doi.org/10.1007/s10741-020-09964-6>
- DeJean, D., Giacomini, M., Vanstone, M., & Brundisini, F. (2013). Patient experiences of depression and anxiety with chronic disease: A systematic review and qualitative meta-synthesis. *Ontario Health Technology Assessment Series, 13*(16), 1–33.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3817854/>
- Dekker, R. L. (2014). Patient perspectives about depressive symptoms in heart failure: A review of the qualitative literature. *The Journal of Cardiovascular Nursing, 29*(1), e9–e15.
<https://doi.org/10.1097/JCN.0b013e318273a5d6>
- Dekker, R. L., Lennie, T. A., Doering, L. V., Chung, M. L., Wu, J. R., & Moser, D. K. (2014). Coexisting anxiety and depressive symptoms in patients with heart failure. *European Journal of Cardiovascular Nursing, 13*(2), 168–176.
<https://doi.org/10.1177/1474515113519520>

- Dekker, R. L., Peden, A. R., Lennie, T. A., Schooler, M. P., & Moser, D. K. (2009). Living with depressive symptoms: Patients with heart failure. *American Journal of Critical Care*, *18*(4), 310–318. <https://doi.org/10.4037/ajcc2009672>
- Dhindsa, D. S., Khambhati, J., Schultz, W. M., Tahhan, A. S., & Quyyumi, A. A. (2020). Marital status and outcomes in patients with cardiovascular disease. *Trends in Cardiovascular Medicine*, *30*(4), 215–220. <https://doi.org/10.1016/j.tcm.2019.05.012>
- Di Marco, F., Verga, M., Reggente, M., Casanova, F. M., Santus, P., Blasi, F., Allegra, L., & Centanni, S. (2006). Anxiety and depression in COPD patients: The roles of gender and disease severity. *Respiratory Medicine*, *100*(10), 1767–1774. <https://doi.org/10.1016/j.rmed.2006.01.026>
- Dickson, V. V., Buck, H., & Riegel, B. (2011). A qualitative meta-analysis of heart failure self-care practices among individuals with multiple comorbid conditions. *Journal of Cardiac Failure*, *17*(5), 413–419. <https://doi.org/10.1016/j.cardfail.2010.11.011>
- Dickson, V. V., Knafel, G. J., & Riegel, B. (2015). Predictors of medication nonadherence differ among Black and White patients with heart failure: Predictors of medication nonadherence. *Research in Nursing & Health*, *38*(4), 289–300. <https://doi.org/10.1002/nur.21663>
- Dickson, V. V., McCarthy, M. M., & Katz, S. M. (2013). How do depressive symptoms influence self-care among an ethnic minority population with heart failure? *Ethnicity & Disease*, *23*(1), 22–28.

- Dickson, V. V., McCarthy, M. M., Howe, A., Schipper, J., & Katz, S. M. (2013). Sociocultural influences on heart failure self-care among an ethnic minority Black population. *Journal of Cardiovascular Nursing*, 28(2), 111–118.
<https://doi.org/10.1097/JCN.0b013e31823db328>
- Dickson, V. V., McCauley, L. A., & Riegel, B. (2008). Work–heart balance: The influence of biobehavioral variables on self-care among employees with heart failure. *Workplace Health & Safety*, 56(2), 63–75. <https://doi.org/10.1177/216507990805600203>
- Doll, K. M. (2018). Investigating Black-White disparities in gynecologic oncology: Theories, conceptual models, and applications. *Gynecologic Oncology*, 149(1), 78–83.
<https://doi.org/10.1016/j.ygyno.2017.10.002>
- Doshi, R. P., Yan, J., & Aseltine, R. H. (2020). Age differences in racial/ethnic disparities in preventable hospitalizations for heart failure in Connecticut, 2009–2015: A population-based longitudinal study. *Public Health Reports*, 135(1), 56–65.
<https://doi.org/10.1177/0033354919884306>
- Dougherty, C. M., Dewhurst, T., Nichol, W. P., & Spertus, J. (1998). Comparison of three quality of life instruments in stable angina pectoris: Seattle Angina Questionnaire, Short Form Health Survey (SF-36), and Quality of Life Index-Cardiac Version III. *Journal of Clinical Epidemiology*, 51(7), 569–575. [https://doi.org/10.1016/S0895-4356\(98\)00028-6](https://doi.org/10.1016/S0895-4356(98)00028-6)
- Drefahl, S. (2012). Do the married really live longer? The role of cohabitation and socioeconomic status. *Journal of Marriage and Family*, 74(3), 462–475.
<https://doi.org/10.1111/j.1741-3737.2012.00968.x>

- Durstenfeld, M. S., Ogedegbe, O., Katz, S. D., Park, H., & Blecker, S. (2016). Racial and ethnic differences in heart failure readmissions and mortality in a large municipal healthcare system. *JACC: Heart Failure*, 4(11), 885–893. <https://doi.org/10.1016/j.jchf.2016.05.008>
- Easton, K., Coventry, P., Lovell, K., Carter, L.-A., & Deaton, C. (2016). Prevalence and measurement of anxiety in samples of patients with heart failure: Meta-analysis. *Journal of Cardiovascular Nursing*, 31(4), 367–379. <https://doi.org/10.1097/JCN.0000000000000265>
- Echouffo-Tcheugui, J. B., Mwasongwe, S. E., Musani, S. K., Hall, M. E., Correa, A., Hernandez, A. F., Golden, S. H., Mentz, R. J., & Bertoni, A. G. (2022). Dysglycemia and incident heart failure among Blacks: The Jackson Heart Study. *The American Heart Journal*, 245, 1–9. <https://doi.org/10.1016/j.ahj.2021.11.003>
- Eisele, M., Rakebrandt, A., Boczor, S., Blozik, E., Träder, J.-M., Störk, S., Herrmann-Lingen, C., & Scherer, M. (2021). The association of general practitioners' awareness of depression and anxiety with change in quality of life in heart failure patients: Results of the prospective observational RECODE-HF cohort study. *Family Practice*, 39(3), 346–353. <https://doi.org/10.1093/fampra/cmab138>
- Ellis, E. M., Orom, H., Giovino, G. A., & Kiviniemi, M. T. (2015). Relations between negative affect and health behaviors by race/ethnicity: Differential effects for symptoms of depression and anxiety. *Health Psychology*, 34(9), 966–969. <https://doi.org/10.1037/hea0000197>
- Evangelista, L. S., Ter-Galstanyan, A., Moughrabi, S., & Moser, D. K. (2009). Anxiety and depression in ethnic minorities with chronic heart failure. *Journal of Cardiac Failure*, 15(7), 572–579. <https://doi.org/10.1016/j.cardfail.2009.03.005>

- Fabbri, M., Murad, M. H., Wennberg, A. M., Turcano, P., Erwin, P. J., Alahdab, F., Berti, A., Manemann, S. M., Yost, K. J., Finney Rutten, L. J., & Roger, V. L. (2020). Health literacy and outcomes among patients with heart failure: A systematic review and meta-analysis. *JACC: Heart Failure*, 8(6), 451–460. <https://doi.org/10.1016/j.jchf.2019.11.007>
- Falk, Doll, S., Gummert, J., Jacquet, L.-M., Glineur, D., Fisher, K., Shipkowitz, T., Crane, D., & Belalcazar, A. (2007). Pilot Study of Chronic Monitoring of Left Ventricular Pressure in Heart Failure Patients. *Journal of Cardiac Failure*, 13(6), S119–S119. <https://doi.org/10.1016/j.cardfail.2007.06.466>
- Felce, D., & Perry, J. (1995). Quality of life: Its definition and measurement. *Research in Developmental Disabilities*, 16(1), 51–74. [https://doi.org/10.1016/0891-4222\(94\)00028-8](https://doi.org/10.1016/0891-4222(94)00028-8)
- Ferguson, L. A., & Pawlak, R. (2011). Health literacy: The road to improved health outcomes. *Journal for Nurse Practitioners*, 7(2), 123–129. <https://doi.org/10.1016/j.nurpra.2010.11.020>
- Ferrans, C. E. (1990). Development of a quality of life index for patients with cancer. *Oncology Nursing Forum*, 17(3 Suppl), 15–19.
- Ferrans, C. E., & Powers, M. J. (1985). Quality of Life Index: Development and psychometric properties. *Advances in Nursing Science*, 8(1), 15–24. <https://doi.org/10.1097/00012272-198510000-00005>

Fleg, J. L., Pina, I. L., Balady, G. J., Chaitman, B. R., Fletcher, B., Lavie, C., Limacher, M. C., Stein, R. A., Williams, M., & Bazzarre, T. (2000). Assessment of functional capacity in clinical and research applications: An advisory from the Committee on Exercise, Rehabilitation, and Prevention, Council on Clinical Cardiology, American Heart Association. *Circulation*, *102*(13), 1591–1597.

<https://doi.org/10.1161/01.CIR.102.13.1591>

Formiga, F., Moreno-Gonzalez, R., Chivite, D., Franco, J., Montero, A., & Corbella, X. (2018). High comorbidity, measured by the Charlson Comorbidity Index, associates with higher 1-year mortality risks in elderly patients experiencing a first acute heart failure hospitalization. *Aging Clinical and Experimental Research*, *30*(8), 927–933.

<https://doi.org/10.1007/s40520-017-0853-1>

Freedland, K. E., Hesseler, M. J., Carney, R. M., Steinmeyer, B. C., Skala, J. A., Davila-Roman, V. G., & Rich, M. W. (2016). Major depression and long-term survival of patients with heart failure. *Psychosomatic Medicine*, *78*(8), 896–903.

<https://doi.org/10.1097/PSY.0000000000000346>

Freedland, K. E., Carney, R. M., & Rich, M. W. (2011). Effect of depression on prognosis in heart failure. *Heart Failure Clinics*, *7*(1), 11–21.

<https://doi.org/10.1016/j.hfc.2010.08.003>

Freedland, K. E., Rich, M. W., & Carney, R. M. (2021). Improving quality of life in heart failure.

Current Cardiology Reports, *23*(11), Article 159. <https://doi.org/10.1007/s11886-021-01588-y>

Freedland, K. E., Rich, M. W., Skala, J. A., Carney, R. M., Davila-Roman, V. G., & Jaffe, A. S. (2003). Prevalence of depression in hospitalized patients with congestive heart failure.

Psychosomatic Medicine, 65(1), 119–128.

<https://doi.org/10.1097/01.PSY.0000038938.67401.85>

García-Olmos, L., Batlle, M., Aguilar, R., Porro, C., Carmona, M., Alberquilla, A., Sánchez-Gómez, L. M., Monge, E., López-Rodríguez, A. B., & Benito, L. (2019). Disability and quality of life in heart failure patients: A cross-sectional study. *Family Practice*, 36(6),

693–698. <https://doi.org/10.1093/fampra/cmz017>

Garin, O., Ferrer, M., Pont, À., Rué, M., Kotzeva, A., Wiklund, I., Van Ganse, E., & Alonso, J. (2009). Disease-specific health-related quality of life questionnaires for heart failure: A systematic review with meta-analyses. *Quality of Life Research*, 18(1), 71–85.

<https://doi.org/10.1007/s11136-008-9416-4>

Geifman, N., Cohen, R., & Rubin, E. (2013). Redefining meaningful age groups in the context of disease. *AGE*, 35(6), 2357–2366. <https://doi.org/10.1007/s11357-013-9510-6>

Gerogianni, S., Babatsikou, F., Gerogianni, G., Grapsa, E., Vasilopoulos, G., Zyga, S., & Koutis, C. (2014). Concerns of patients on dialysis: A research study. *Health Science Journal*, 8(4), 423–437.

Ghods, B. K., Roter, D. L., Ford, D. E., Larson, S., Arbelaez, J. J., & Cooper, L. A. (2008). Patient–physician communication in the primary care visits of African Americans and Whites with depression. *Journal of General Internal Medicine*, 23(5), 600–606.

<https://doi.org/10.1007/s11606-008-0539-7>

- Glynn, P., Lloyd-Jones, D. M., Feinstein, M. J., Carnethon, M., & Khan, S. S. (2019). Disparities in cardiovascular mortality related to heart failure in the United States. *Journal of the American College of Cardiology*, *73*(18), 2354–2355. <https://doi.org/10.1016/j.jacc.2019.02.042>
- Go, A. S., Mozaffarian, D., Roger, V. L., Benjamin, E. J., Berry, J. D., Blaha, M. J., Dai, S., Ford, E. S., Fox, C. S., Franco, S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Huffman, M. D., Judd, S. E., Kissela, B. M., Kittner, S. J., ... Turner, M. B. (2013). Heart disease and stroke statistics—2014 update: A report from the American Heart Association. *Circulation*, *129*(3). <https://doi.org/10.1161/01.cir.0000441139.02102.80>
- Gordon, H. S., Nowlin, P. R., Maynard, D., Berbaum, M. L., & Deswal, A. (2010). Mortality after hospitalization for heart failure in Blacks compared to Whites. *The American Journal of Cardiology*, *105*(5), 694–700. <https://doi.org/10.1016/j.amjcard.2009.10.051>
- Gorini, A., Giuliani, M., Raggio, L., Barbieri, S., & Tremoli, E. (2020). Depressive and anxiety symptoms screening in cardiac inpatients: A virtuous Italian approach to psychocardiology. *International Journal of Environmental Research and Public Health*, *17*(14), Article 5007. <https://doi.org/10.3390/ijerph17145007>
- Gottlieb, S. S., Khatta, M., Friedmann, E., Einbinder, L., Katzen, S., Baker, B., Marshall, J., Minshall, S., Robinson, S., Fisher, M. L., Potenza, M., Sigler, B., Baldwin, C., & Thomas, S. A. (2004). The influence of age, gender, and race on the prevalence of depression in heart failure patients. *Journal of the American College of Cardiology*, *43*(9), 1542–1549. <https://doi.org/10.1016/j.jacc.2003.10.064>

- Goyal, A., Lahan, S., Bhyan, P., Ranka, S., Pothuru, S., Mehta, H., Dalia, T., Patel, N., Acharya, P., Truyen, T. T. T. T., & Mazek, H. (2022). Mortality trends in heart failure and differences by race and sex in the United States: 1999–2019. *Journal of the American College of Cardiology*, 79(9_Supplement), Article 275.
- Granger, B. B., Sandelowski, M., Tahshjain, H., Swedberg, K., & Ekman, I. (2009). A qualitative descriptive study of the work of adherence to a chronic heart failure regimen: patient and physician perspectives. *Journal of Cardiovascular Nursing*, 24(4), 308–315.
<https://doi.org/10.1097/JCN.0b013e3181a4be30>
- Griffin, M. T. Q., Lee, Y.-H., Salman, A., Seo, Y., Marin, P. A., Starling, R. C., & Fitzpatrick, J. J. (2007). Spirituality and well being among elders: Differences between elders with heart failure and those without heart failure. *Clinical Interventions in Aging*, 2(4), 669–675.
<https://doi.org/10.2147/cia.s874>
- Grundström, J., Kontinen, H., Berg, N., & Kiviruusu, O. (2021). Associations between relationship status and mental well-being in different life phases from young to middle adulthood. *SSM: Population Health*, 14, Article 100774.
<https://doi.org/10.1016/j.ssmph.2021.100774>
- Havranek, E. P., Spertus, J. A., Masoudi, F. A., Jones, P. G., & Rumsfeld, J. S. (2004). Predictors of the onset of depressive symptoms in patients with heart failure. *Journal of the American College of Cardiology*, 44(12), 2333–2338.
<https://doi.org/10.1016/j.jacc.2004.09.034>
- HealthyPeople.gov. (n.d.). *Employment*. Office of Disease Prevention and Health Promotion.
<https://www.healthypeople.gov/2020/topics-objectives/topic/social-determinants-health/interventions-resources/employment>

- Heiat, A., Gross, C. P., & Krumholz, H. M. (2002). Representation of the elderly, women, and minorities in heart failure clinical trials. *Archives of Internal Medicine*, *162*(15), 1682–1688. <https://doi.org/10.1001/archinte.162.15.1682>
- Heo, S., Lennie, T. A., Okoli, C., & Moser, D. K.. (2009). Quality of life in patients with heart failure: Ask the patients. *Heart & Lung*, *38*(2), 100–108. <https://doi.org/10.1016/j.hrtlng.2008.04.002>
- Heo, S., Moser, D. K., Chung, M. L., & Lennie, T. A. (2012). Social status, health-related quality of life, and event-free survival in patients with heart failure. *European Journal of Cardiovascular Nursing*, *11*(2), 141–149. <https://doi.org/10.1016/j.ejcnurse.2010.10.003>
- Heo, S., Shin, M.-S., Hwang, S. Y., An, M., Park, J.-K., Kim, S., Shim, J. L., & Kim, J. (2019). Sex differences in heart failure symptoms and factors associated with heart failure symptoms. *The Journal of Cardiovascular Nursing*, *34*(4), 306–312. <https://doi.org/10.1097/JCN.0000000000000579>
- Herr, J. K., Salyer, J., Flattery, M., Goodloe, L., Lyon, D. E., Kabban, C. S., & Clement, D. G. (2015). Heart failure symptom clusters and functional status: A cross-sectional study. *Journal of Advanced Nursing*, *71*(6), 1274–1287. <https://doi.org/10.1111/jan.12596>
- Herr, J. K., Salyer, J., Lyon, D. E., Goodloe, L., Schubert, C., & Clement, D. G. (2013). Heart failure symptom relationships: A systematic review. *Journal of Cardiovascular Nursing*, *29*(5), 416–422. <https://doi.org/10.1097/JCN.0b013e31829b675e>
- Hill, C. V., Perez-Stable, E. J., Anderson, N. A., & Bernard, M. A. (2015). The National Institute on Aging Health Disparities Research Framework. *Ethnicity & Disease*, *25*(3), 245–254. <https://doi.org/10.18865/ed.25.3.245>

- Hill-Joseph, E. A. (2019). Coping while Black: Chronic illness, mastery, and the Black-White health paradox. *Journal of Racial and Ethnic Health Disparities*, 6(5), 935–943.
<https://doi.org/10.1007/s40615-019-00594-9>
- Hollingshaus, M. S., & Utz, R. L. (2013). Depressive symptoms following the diagnosis of major chronic illness. *Society and Mental Health*, 3(1), 22–39.
<https://doi.org/10.1177/2156869312464788>
- Holzappel, N., Muller-Tasch, T., Wild, B., Junger, J., Zugck, C., Remppis, A., Herzog, W., & Lowe, B. (2008). Depression profile in patients with and without chronic heart failure. *Journal of Affective Disorders*, 105(1–3), 53–62.
<https://doi.org/10.1016/j.jad.2007.04.009>
- Holzappel, N., Muller-Tasch, T., Wild, B., Junger, J., Zugck, C., Remppis, A., Herzog, W., & Lowe, B. (2008). Depression profile in patients with and without chronic heart failure. *Journal of Affective Disorders*, 105(1–3), 53–62.
<https://doi.org/10.1016/j.jad.2007.04.009>
- Hopper, I., Kotecha, D., Chin, K. L., Mentz, R. J., & von Lueder, T. G. (2016). Comorbidities in heart failure: Are there gender differences? *Current Heart Failure Reports*, 13(1), 1–12.
<https://doi.org/10.1007/s11897-016-0280-1>
- Hwang, S.-L., Liao, W.-C., & Huang, T.-Y. (2014). Predictors of quality of life in patients with heart failure. *Japan Journal of Nursing Science*, 11(4), 290–298.
<https://doi.org/10.1111/jjns.12034>
- IBM Corp. (2020). IBM SPSS Statistics for Windows, Version 27.0.

- Ishak, W. W., Edwards, G., Herrera, N., Lin, T., Hren, K., Peterson, M., Ngor, A., Liu, A., Kimchi, A., & Spiegel, B. (2020). Depression in heart failure: A systematic review. *Innovations in Clinical Neuroscience*, *17*(4-6), 27–38.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7413333/>
- Jaarsma, T., Stromberg, A., Ben Gal, T., Cameron, J., Driscoll, A., Duengen, H. D., Inkrot, S., Huang, T. Y., Huyen, N. N., Kato, N., Koberich, S., Lupon, J., Moser, D. K., Pulignano, G., Rabelo, E. R., Suwanno, J., Thompson, D. R., Vellone, E., Alvaro, R., ... Riegel, B. (2013). Comparison of self-care behaviors of heart failure patients in 15 countries worldwide. *Patient Education and Counseling*, *92*(1), 114–120.
<https://doi.org/10.1016/j.pec.2013.02.017>
- Jackson, J. L., & Emery, C. F. (2011). Illness knowledge moderates the influence of coping style on quality of life among women with congestive heart failure. *Heart & Lung*, *40*(2), 122–129. <https://doi.org/10.1016/j.hrtlng.2009.12.005>
- Jiang, Krishnan, R., Kuchibhatla, M., Cuffe, M. S., Martsberger, C., Arias, R. M., & O'Connor, C. M. (2011). Characteristics of Depression Remission and Its Relation With Cardiovascular Outcome Among Patients With Chronic Heart Failure (from the SADHART-CHF Study). *The American Journal of Cardiology*, *107*(4), 545–551.
<https://doi.org/10.1016/j.amjcard.2010.10.013>
- Jiang, W., Alexander, J., Christopher, E., Kuchibhatla, M., Gauden, L. H., Cuffe, M. S., Blazing, M. A., Davenport, C., Califf, R. M., Krishnan, R. R., & O'Connor, C. M. (2001). Relationship of depression to increased risk of mortality and rehospitalization in patients with congestive heart failure. *Archives of Internal Medicine*, *161*(15), 1849–1856. <https://doi.org/10.1001/archinte.161.15.1849>

- Johansson, P., Dahlström, U., & Broström, A. (2006a). Consequences and predictors of depression in patients with chronic heart failure: Implications for nursing care and future research. *Progress in Cardiovascular Nursing*, 21(4), 202–211.
<https://doi.org/10.1111/j.0889-7204.2006.05415.x>
- Johansson, P., Dahlström, U., & Broström, A. (2006b). The measurement and prevalence of depression in patients with chronic heart failure. *Progress in Cardiovascular Nursing*, 21(1), 28–36. <https://doi.org/10.1111/j.0197-3118.2006.04644.x>
- Johnson-Lawrence, V., Griffith, D. M., & Watkins, D. C. (2013). The effects of race, ethnicity, and mood/anxiety disorders on the chronic physical health conditions of men from a national sample. *American Journal of Men's Health*, 7(4_suppl), 58S–67S.
<https://doi.org/10.1177/1557988313484960>
- Jones, G. C., & Crews, J. E. (2013). Health disparities among workers and nonworkers with functional limitations: Implications for improving employment in the United States. *Disabil Rehabil*, 35(17), 1479–1490. <https://doi.org/10.3109/09638288.2012.740137>
- Joynt, K. E., Whellan, D. J., & O'Connor, C. M. (2004). Why is depression bad for the failing heart? A review of the mechanistic relationship between depression and heart failure. *Journal of Cardiac Failure*, 10(3), 258–271.
<https://doi.org/10.1016/j.cardfail.2003.09.008>
- Kalsoom, U. (2020). Gender role in anxiety, depression and quality of life in chronic kidney disease patients. *Pakistan Journal of Medical Sciences*, 36(2), 251–254.
<https://doi.org/10.12669/pjms.36.2.869>

- Katon, W., Lin, E. H., & Kroenke, K. (2007). The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *General Hospital Psychiatry, 29*(2), 147–155. <https://doi.org/10.1016/j.genhosppsych.2006.11.005>
- Katz, S. D. (2013). *Heart failure: A practical guide for diagnosis and management*. Oxford University Press.
- Kewcharoen, J., Thangjui, S., Kanitsoraphan, C., Techorueangwiwat, C., Mekraksakit, P., & Vutthikraivit, W. (2021). The effects of marital status on outcome of heart failure population: A systematic review and meta-analysis. *Acta cardiologica, 76*(1), 11–19. <https://doi.org/10.1080/00015385.2019.1699281>
- Khayyat, I., Huynh, T., & Khayyat, H. (2021). Black patients' representation in major randomized controlled trials of heart failure with preserved ejection fraction. *Canadian Journal of Cardiology, 37*(10), S67–S68. <https://doi.org/10.1016/j.cjca.2021.07.139>
- Kickbusch, I. S. (2001). Health literacy: Addressing the health and education divide. *Health Promotion International, 16*(3), 289–297. <https://doi.org/10.1093/heapro/16.3.289>
- Kim, K. U., Park, H. K., Jung, H. Y., Ahn, J. J., Moon, E., Kim, Y. S., Lee, M. K., & Lee, H. (2014). Association of depression with disease severity in patients with chronic obstructive pulmonary disease. *Lung, 192*(2), 243–249. <https://doi.org/10.1007/s00408-013-9547-4>
- Koenig, H. G. (1998). Depression in hospitalized older patients with congestive heart failure. *General Hospital Psychiatry, 20*(1), 29–43. [https://doi.org/10.1016/S0163-8343\(98\)80001-7](https://doi.org/10.1016/S0163-8343(98)80001-7)

Konstam, V., Moser, D. K., & De Jong, M. J. (2005). Depression and anxiety in heart failure. *Journal of Cardiac Failure, 11*(6), 455–463.

<https://doi.org/10.1016/j.cardfail.2005.03.006>

Kusek, J. W., Greene, P., Wang, S.-R., Beck, G., West, D., Jamerson, K., Agodoa, L. Y., Faulkner, M., & Level, B. (2002). Cross-sectional study of health-related quality of life in African Americans with chronic renal insufficiency: The African American Study of Kidney Disease and Hypertension Trial. *American Journal of Kidney Diseases, 39*(3), 513–524. <https://doi.org/10.1053/ajkd.2002.31401>

Lam, J., & Perales, F. (2017, January). *Chronic illness and mental strain: The moderating role of marital status over the disease cycle*. ARC Center of Excellence for Children and Families Over the Life Course.

Landsbergis, P. A., Choi, B., Dobson, M., Sembajwe, G., Slatin, C., Delp, L., Siqueira, C. E., Schnall, P., & Baron, S. (2018). The key role of work in population health inequities. *American Journal of Public Health, 108*(3), 296–297.

<https://doi.org/10.2105/AJPH.2017.304288>

Lang, C. C., & Mancini, D. M. (2007). Non-cardiac comorbidities in chronic heart failure. *Heart, 93*(6), 665–671. <https://doi.org/10.1136/hrt.2005.068296>

Lee, D. T. F., Yu, D. S. F., Woo, J., & Thompson, D. R. (2005). Health-related quality of life in patients with congestive heart failure. *European Journal of Heart Failure, 7*(3), 419–422.

<https://doi.org/10.1016/j.ejheart.2004.08.004>

- Lee, K. S., Lennie, T. A., Yoon, J. Y., Wu, J.-R., & Moser, D. K. (2017). Living Arrangements Modify The Relationship Between Depressive Symptoms And Self-Care In Patients With Heart Failure. *The Journal of Cardiovascular Nursing*, 32(2), 171–179.
<https://doi.org/10.1097/JCN.0000000000000327>
- Lekavich, C. L., & Barksdale, D. J. (2016). A critical evaluation of the representation of Black patients with heart failure and preserved ejection fraction in clinical trials: A literature review. *The Journal of Cardiovascular Nursing*, 31(3), 202–208.
<https://doi.org/10.1097/JCN.0000000000000237>
- Lenze, E. J., Mulsant, B. H., Shear, M. K., Schulberg, H. C., Dew, M. A., Begley, A. E., Pollock, B. G., & Reynolds, C. F. (2000). Comorbid anxiety disorders in depressed elderly patients. *American Journal of Psychiatry*, 157(5), 722–728.
<https://doi.org/10.1176/appi.ajp.157.5.722>
- Lesman-Leegte, I., van Veldhuisen, D. J., Hillege, H. L., Moser, D., Sanderman, R., & Jaarsma, T. (2009). Depressive symptoms and outcomes in patients with heart failure: Data from the COACH study. *European Journal of Heart Failure*, 11(12), 1202–1207.
<https://doi.org/10.1093/eurjhf/hfp155>
- Lewis, T. T., Guo, H., Lunos, S., Mendes de Leon, C. F., Skarupski, K. A., Evans, D. A., & Everson-Rose, S. A. (2011). Depressive symptoms and cardiovascular mortality in older African-American and White adults: Evidence for a differential association by race. *Circulation Cardiovascular Quality and Outcomes*, 4(3), 293–299.
<https://doi.org/10.1161/CIRCOUTCOMES.110.957548>

- Lewsey, S. C., & Breathett, K. (2021). Racial and ethnic disparities in heart failure: Current state and future directions. *Current Opinion in Cardiology*, *36*(3), 320–328.
<https://doi.org/10.1097/HCO.0000000000000855>
- Li, C., Barker, L., Ford, E. S., Zhang, X., Strine, T. W., & Mokdad, A. H. (2008). Diabetes and anxiety in US adults: Findings from the 2006 Behavioral Risk Factor Surveillance System. *Diabetic Medicine*, *25*(7), 878–881. <https://doi.org/10.1111/j.1464-5491.2008.02477.x>
- Liew, H.-P. (2012). Depression and chronic illness: A test of competing hypotheses. *Journal of Health Psychology*, *17*(1), 100–109. <https://doi.org/10.1177/1359105311409788>
- Lim, L., Jin, A.-Z., & Ng, T.-P. (2011). Anxiety and depression, chronic physical conditions, and quality of life in an urban population sample study. *Social Psychiatry and Psychiatric Epidemiology*, *47*(7), 1047–1053. <https://doi.org/10.1007/s00127-011-0420-6>
- Liu, M.-H., Chiou, A.-F., Wang, C.-H., Yu, W.-P., & Lin, M.-H. (2021). Relationship of symptom stress, care needs, social support, and meaning in life to quality of life in patients with heart failure from the acute to chronic stages: A longitudinal study. *Health Quality of Life Outcomes*, *19*(1), Article 252. <https://doi.org/10.1186/s12955-021-01885-8>
- Liu, X., Haagsma, J., Sijbrands, E., Buijks, H., Boogaard, L., Mackenbach, J., Erasmus, V., & Polinder, S. (2020). Anxiety and depression in diabetes care: Longitudinal associations with health-related quality of life. *Scientific Reports*, *10*(1), Article 8307.
<https://doi.org/10.1038/s41598-020-57647-x>

- Luttik, M. L., Jaarsma, T., Veeger, N., & van Veldhuisen, D. J. (2006). Marital status, quality of life, and clinical outcome in patients with heart failure. *Heart & Lung, 35*(1), 3–8.
<https://doi.org/10.1016/j.hrtlng.2005.08.001>
- MacMahon, K. M., & Lip, G. Y. (2002). Psychological factors in heart failure: A review of the literature. *Archives of Internal Medicine, 162*(5), 509–516.
<https://doi.org/10.1001/archinte.162.5.509>
- Manfredini, R., De Giorgi, A., Tiseo, R., Boari, B., Cappadona, R., Salmi, R., Gallerani, M., Signani, F., Manfredini, F., Mikhailidis, D. P., & Fabbian, F. (2017). Marital status, cardiovascular diseases, and cardiovascular risk factors: A review of the evidence. *Journal of Women's Health, 26*(6), 624–632. <https://doi.org/10.1089/jwh.2016.6103>
- Manzoli, L., Villari, P., Pirone, G. M., & Boccia, A. (2007). Marital status and mortality in the elderly: A systematic review and meta-analysis. *Social Science & Medicine, 64*(1), 77–94. <https://doi.org/10.1016/j.socscimed.2006.08.031>
- McCann, K., & Boore, J. R. (2000). Fatigue in persons with renal failure who require maintenance haemodialysis. *Journal of Advanced Nursing, 32*(5), 1132–1142.
<https://doi.org/10.1046/j.1365-2648.2000.01584.x>
- McCarthy, M., Katz, S. D., Schipper, J., & Dickson, V. V. (2015). “I just can’t do it anymore”: Patterns of African Americans with heart failure: A mixed method study. *Healthcare, 3*(4), 973–986. <https://doi.org/10.3390/healthcare3040973>
- McLaughlin, T. J., Aupont, O., Bambauer, K. Z., Stone, P., Mullan, M. G., Colagiovanni, J., Polishuk, E., Johnstone, M., & Locke, S. E. (2005). Improving psychologic adjustment to chronic illness in cardiac patients: The role of depression and anxiety. *Journal of General Internal Medicine, 20*(12), 1084–1090. <https://doi.org/10.1111/j.1525-1497.2005.00256.x>

- Mentz, R. J., Babyak, M. A., Bittner, V., Fleg, J. L., Keteyian, S. J., Swank, A. M., Pina, I. L., Kraus, W. E., Whellan, D. J., O'Connor, C. M., & Blumenthal, J. A. (2015). Prognostic significance of depression in Blacks with heart failure: Insights from Heart Failure: A Controlled Trial Investigating Outcomes of Exercise Training. *Circulation: Heart Failure*, 8(3), 497–503. <https://doi.org/10.1161/CIRCHEARTFAILURE.114.001995>
- Merriam-Webster. (n.d.-a). *Age*. <https://www.merriam-webster.com/dictionary/age>
- Merriam-Webster. (n.d.-b). *Marital status*. <https://www.merriam-webster.com/dictionary/marital%20status>
- Minhas, A. M. K., Qavi, A. H., Ijaz, S. H., Khan, M. Z., Fudim, M., & Nasir, K. (2022). Ethnic and racial disparities in heart failure related mortality rates among older adults: Data from CDC Wonder 1999–2019. *Journal of Cardiac Failure*, 28(5), S110–S111. <https://doi.org/10.1016/j.cardfail.2022.03.282>
- Morrow, D., Clark, D., Tu, W., Wu, J., Weiner, M., Steinley, D., & Murray, M. D. (2006). Correlates of health literacy in patients with chronic heart failure. *The Gerontologist*, 46(5), 669–676. <https://doi.org/10.1093/geront/46.5.669>
- Mouzon, D. M. (2017). Religious involvement and the Black–White paradox in mental health. *Race and Social Problems*, 9(1), 63–78. <https://doi.org/10.1007/s12552-017-9198-9>
- Mwansa, H., Lewsey, S., Mazimba, S., & Breathett, K. (2021). Racial/ethnic and gender disparities in heart failure with reduced ejection fraction. *Current Heart Failure Reports*, 18(2), 41–51. <https://doi.org/10.1007/s11897-021-00502-5>
- Nair, N., Farmer, C., Gongora, E., & Dehmer, G. J. (2012). Commonality between depression and heart failure. *American Journal of Cardiology*, 109(5), 768–772. <https://doi.org/10.1016/j.amjcard.2011.10.039>

National Institute for Nursing Research, n.d. National Institute for Nursing Research's (NINR) Strategic Plan's (NINR) Strategic Plan. Retrieved from,

<https://www.ninr.nih.gov/aboutninr/ninr-mission-and-strategic-plan>

Nayak, A., Hicks, A. J., & Morris, A. A. (2020). Understanding the complexity of heart failure risk and treatment in Black patients. *Circulation. Heart Failure*, 13(8), Article e007264.

<https://doi.org/10.1161/CIRCHEARTFAILURE.120.007264>

NIMHD Minority Health and Health Disparities. (2017). *NIMHD Minority Health and Health Disparities research framework*. <https://nimhd.nih.gov/researchFramework>

O'Connor, C., Fiuzat, M., Mulder, H., Coles, A., Ahmad, T., Ezekowitz, J. A., Adams, K. F., Piña, I. L., Anstrom, K. J., Cooper, L. S., Mark, D. B., Whellan, D. J., Januzzi, J. L., Leifer, E. S., & Felker, G. M. (2019). Clinical factors related to morbidity and mortality in high-risk heart failure patients: The GUIDE-IT predictive model and risk score.

European Journal of Heart Failure, 21(6), 770–778. <https://doi.org/10.1002/ejhf.1450>

O'Neill, E. S., & Morrow, L. L. (2001). The symptom experience of women with chronic illness: Women's symptom experience. *Journal of Advanced Nursing*, 33(2), 257–268.

<https://doi.org/10.1111/j.1365-2648.2001.01653.x>

OECD. (n.d.). *Glossary of statistical terms*. <https://stats.oecd.org/glossary/index.htm>

Ohlsson, A., Eckerdal, N., Lindahl, B., Hanning, M., & Westerling, R. (2021). Non-employment and low educational level as risk factors for inequitable treatment and mortality in heart

failure: A population-based cohort study of register data. *BMC Public Health*, 21(1),

Article 1040. <https://doi.org/10.1186/s12889-021-10919-1>

- Ojeda, V. D., & McGuire, T. G. (2006). Gender and racial/ethnic differences in use of outpatient mental health and substance use services by depressed adults. *Psychiatric Quarterly*, 77(3), 211–222. <https://doi.org/10.1007/s11126-006-9008-9>
- Parashar, S., Katz, R., Smith, N. L., Arnold, A. M., Vaccarino, V., Wenger, N. K., & Gottdiener, J. S. (2009). Race, gender, and mortality in adults ≥ 65 years of age with incident heart failure (from the Cardiovascular Health Study). *The American Journal of Cardiology*, 103(8), 1120–1127. <https://doi.org/10.1016/j.amjcard.2008.12.043>
- Patel, R. S., Shrestha, S., Saeed, H., Raveendranathan, S., Isidahome, E. E., Ravat, V., Fakorede, M. O., & Patel, V. (2018). Comorbidities and consequences in hospitalized heart failure patients with depression. *Cureus*, 10(8), Article e3193. <https://doi.org/10.7759/cureus.3193>
- Pelle, A. J., Gidron, Y. Y., Szabó, B. M., & Denollet, J. (2008). Psychological predictors of prognosis in chronic heart failure. *Journal of Cardiac Failure*, 14(4), 341–350. <https://doi.org/10.1016/j.cardfail.2008.01.004>
- Peterson, P. N., Shetterly, S. M., Clarke, C. L., Bekelman, D. B., Chan, P. S., Allen, L. A., Matlock, D. D., Magid, D. J., & Masoudi, F. A. (2011). Health literacy and outcomes among patients with heart failure. *JAMA*, 305(16), 1695–1701. <https://doi.org/10.1001/jama.2011.512>
- Phillips, S. P. (2005). Defining and measuring gender: A social determinant of health whose time has come. *International Journal for Equity in Health*, 4(1), Article 11. <https://doi.org/10.1186/1475-9276-4-11>

- Piamjariyakul, U., Smith, C. E., Wick, J., & Fitzgerald, S. (2013). Nurse-led group clinic intervention reduced depression among African American patients with heart failure. *Clinical and Translational Science*, 6(2), 128–132.
<https://doi.org/10.1016/j.hrtlng.2018.02.005>
- Piamjariyakul, U., Thompson, N. C., Russell, C., & Smith, C. E. (2018). The effect of nurse-led group discussions by race on depressive symptoms in patients with heart failure. *Heart & Lung*, 47(3), 211–215. <https://doi.org/10.1016/j.hrtlng.2018.02.005>
- Pindikura, R. K., Gunturu, A. G., & Kadiveti, U. K. (2019). Depression, anxiety, and their association with sociodemographic factors in patients undergoing haemodialysis: A cross-sectional study. *Open Journal of Psychiatry & Allied Sciences*, 10(1), 44–48.
<https://doi.org/10.5958/2394-2061.2019.00011.9>
- Polikandrioti, M., Panoutsopoulos, G., Tsami, A., Gerogianni, G., Saroglou, S., Thomai, E., & Leventzonis, I. (2019). Assessment of quality of life and anxiety in heart failure outpatients. *Archives of Medical Sciences: Atherosclerotic Diseases*, 4, Article e38.
<https://doi.org/10.5114/amsad.2019.84444>
- Potluri, R., Carter, P., Patel, B., More, R., & Lavu, D. (2019). The impact of gender and marital status on long term mortality in patients with cardiovascular disease: Insights utilizing big data from the ACALM study. *Journal of the American College of Cardiology*, 73(9), Article 1832. [https://doi.org/10.1016/S0735-1097\(19\)32438-6](https://doi.org/10.1016/S0735-1097(19)32438-6)
- Pumar, M. I., Gray, C. R., Walsh, J. R., Yang, I. A., Rolls, T. A., & Ward, D. L. (2014). Anxiety and depression—Important psychological comorbidities of COPD. *Journal of Thoracic Disease*, 6(11), 1615–1631. <https://doi.org/10.3978%2Fj.issn.2072-1439.2014.09.28>

- Raphael, C., Briscoe, C., Davies, J., Whinnett, Z. I., Manisty, C., Sutton, R., Mayet, J., & Francis, D. P. (2007). Limitations of the New York Heart Association functional classification system and self-reported walking distances in chronic heart failure. *Heart*, 93(4), 476–482.
- Rathore, S. S., Foody, J. M., Wang, Y., Smith, G. L., Herrin, J., Masoudi, F. A., Wolfe, P., Havranek, E. P., Ordin, D. L., & Krumholz, H. M. (2003). Race, quality of care, and outcomes of elderly patients hospitalized with heart failure. *JAMA*, 289(19), 2517–2524. <https://doi.org/10.1001/jama.289.19.2517>
- Read, J., Sharpe, L., Modini, M., & Dear, B. F. (2017). Multimorbidity and depression: A systematic review and meta-analysis. *Journal of Affective Disorders*, 221, 36–46. <https://doi.org/10.1016/j.jad.2017.06.009>
- Rechenberg, K., Cousin, L., & Redwine, L. (2020). Mindfulness, anxiety symptoms, and quality of life in heart failure. *Journal of Cardiovascular Nursing*, 35(4), 358–363. <https://doi.org/10.1097/JCN.0000000000000630>
- Reddy, Y. N. V., Rikhi, A., Obokata, M., Shah, S. J., Lewis, G. D., AbouEzzedine, O. F., Dunlay, S., McNulty, S., Chakraborty, H., Stevenson, L. W., Redfield, M. M., & Borlaug, B. A. (2020). Quality of life in heart failure with preserved ejection fraction: Importance of obesity, functional capacity, and physical inactivity. *European Journal of Heart Failure*, 22(6), 1009–1018. <https://doi.org/10.1002/ejhf.1788>
- Reza, N., Gruen, J., & Bozkurt, B. (2022). Representation of women in heart failure clinical trials: Barriers to enrollment and strategies to close the gap. *American Heart Journal Plus: Cardiology Research and Practice*, 13, Article 100093. <https://doi.org/10.1016/j.ahjo.2022.100093>

- Riegel, Lee, C. S., Ratcliffe, S. J., De Geest, S., Potashnik, S., Patey, M., Sayers, S. L., Goldberg, L. R., & Weintraub, W. S. (2012). Predictors of Objectively Measured Medication Nonadherence in Adults With Heart Failure. *Circulation. Heart Failure*, 5(4), 430–436. <https://doi.org/10.1161/CIRCHEARTFAILURE.111.965152>
- Riegel, B., Driscoll, A., Suwanno, J., Moser, D. K., Lennie, T. A., Chung, M. L., Wu, J. R., Dickson, V. V., Carlson, B., & Cameron, J. (2009). Heart failure self-care in developed and developing countries. *Journal of Cardiac Failure*, 15(6), 508–516. <https://doi.org/10.1016/j.cardfail.2009.01.009>
- Riegel, B., Moser, D. K., Rayens, M. K., Carlson, B., Pressler, S. J., Shively, M., Albert, N. M., Armola, R. R., Evangelista, L., Westlake, C., & Sethares, K. (2008). Ethnic differences in quality of life in persons with heart failure. *Journal of Cardiac Failure*, 14(1), 41–47. <https://doi.org/10.1016/j.cardfail.2007.09.008>
- Roger, V. L., Go, A. S., Lloyd-Jones, D. M., Benjamin, E. J., Berry, J. D., Borden, W. B., Bravata, D. M., Dai, S., Ford, E. S., Fox, C. S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Kissela, B. M., Kittner, S. J., Lackland, D. T., Lichtman, J. H., ... Turner, M. B. (2012). Heart disease and stroke statistics—2012 update: A report from the American Heart Association. *Circulation*, 125(1), e2–e220. <https://doi.org/10.1161/CIR.0b013e31823ac046>
- Rohrbaugh, M. J., Shoham, V., & Coyne, J. C. (2006). Effect of marital quality on eight-year survival of patients with heart failure. *American Journal of Cardiology*, 98(8), 1069–1072. <https://doi.org/10.1016/j.amjcard.2006.05.034>

- Rohyans, L. M., & Pressler, S. J. (2009). Depressive symptoms and heart failure: Examining the sociodemographic variables. *Clinical Nurse Specialist: The Journal for Advanced Nursing Practice*, 23(3), 138–144. <https://doi.org/10.1097/NUR.0b013e3181a443b4>
- Rørth, R., Fosbøl, E. L., Mogensen, U. M., Kragholm, K., Numé, A. K., Gislason, G. H., Jhund, P. S., Petrie, M. C., McMurray, J. J. V., Torp-Pedersen, C., Køber, L., & Kristensen, S. L. (2018). Employment status at time of first hospitalization for heart failure is associated with a higher risk of death and rehospitalization for heart failure. *European Journal of Heart Failure*, 20(2), 240–247. <https://doi.org/10.1002/ejhf.1046>
- Ross, C. E., & Mirowsky, J. (1999). Refining the association between education and health: The effects of quantity, credential, and selectivity. *Demography*, 36(4), 445–460. <https://doi.org/10.2307/2648083>
- Ross, C. E., & Wu, C. L. (1995). The links between education and health. *American Sociological Review*, 60(5), 719–745. <https://doi.org/10.2307/2096319>
- Rumsfeld, J. S., Havranek, E., Masoudi, F. A., Peterson, E. D., Jones, P., Tooley, J. F., Krumholz, H. M., Spertus, J. A., & Cardiovascular Outcomes Research Consortium. (2003). Depressive symptoms are the strongest predictors of short-term declines in health status in patients with heart failure. *Journal of the American College of Cardiology*, 42(10), 1811–1817. <https://doi.org/10.1016/j.jacc.2003.07.013>
- Rustad, J. K., Stern, T. A., Hebert, K. A., & Musselman, D. L. (2013). Diagnosis and treatment of depression in patients with congestive heart failure: A review of the literature. *The Primary Care Companion for CNS Disorders*, 15(4). <https://doi.org/10.4088/PCC.13r01511>

- Rutledge, T., Reis, V. A., Linke, S. E., Greenberg, B. H., & Mills, P. J. (2006). Depression in heart failure: A meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. *Journal of the American College of Cardiology*, *48*(8), 1527–1537. <https://doi.org/10.1016/j.jacc.2006.06.055>
- Ryu, E., Chamberlain, A. M., Pendegraft, R. S., Petterson, T. M., Bobo, W. V., & Pathak, J. (2016). Quantifying the impact of chronic conditions on a diagnosis of major depressive disorder in adults: A cohort study using linked electronic medical records. *BMC Psychiatry*, *16*, Article 114. <https://doi.org/10.1186/s12888-016-0821-x>
- Sadlonova, M., Wasser, K., Nagel, J., Weber-Krüger, M., Gröschel, S., Uphaus, T., Liman, J., Hamann, G. F., Kermer, P., Gröschel, K., Herrmann-Lingen, C., & Wachter, R. (2021). Health-related quality of life, anxiety and depression up to 12 months post-stroke: Influence of sex, age, stroke severity and atrial fibrillation – A longitudinal subanalysis of the Find-AF_{RANDOMISED} trial. *Journal of Psychosomatic Research*, *142*, Article 110353. <https://doi.org/10.1016/j.jpsychores.2020.110353>
- Salyer, J., Flattery, M., & Lyon, D. E. (2019). Heart failure symptom clusters and quality of life. *Heart & Lung*, *48*(5), 366–372. <https://doi.org/10.1016/j.hrtlng.2019.05.016>
- Samartzis, L., Dimopoulos, S., Tziogourou, M., & Nanas, S. (2013). Effect of psychosocial interventions on quality of life in patients with chronic heart failure: A meta-analysis of randomized controlled trials. *Journal of Cardiac Failure*, *19*(2), 125–134. <https://doi.org/10.1016/j.cardfail.2012.12.004>
- Schöllgen, I., Morack, J., Infurna, F. J., Ram, N., & Gerstorf, D. (2016). Health sensitivity: Age differences in the within-person coupling of individuals' physical health and well-being. *Developmental Psychology*, *52*(11), 1944–1953. <https://doi.org/10.1037/dev0000171>

- Schultz, W. M., Hayek, S. S., Samman Tahhan, A., Ko, Y. A., Sandesara, P., Awad, M., Mohammed, K. H., Patel, K., Yuan, M., & Zheng, S. (2017). Marital status and outcomes in patients with cardiovascular disease. *Journal of the American Heart Association*, 6(12), Article e005890. <https://doi.org/10.1161/JAHA.117.005890>
- Seo, Y., Wang, J., Barnes, D., & Barshikar, S. (2021). Heart failure and disability in community-dwelling older adults in the United States—1999 to 2018: Data from National Health and Nutrition Examination Survey. *Clinical Nursing Research*, 571–578. <https://doi.org/10.1177/10547738211061620>
- Serdar, C. C., Cihan, M., Yücel, D., & Serdar, M. A. (2021). Sample size, power and effect size revisited: Simplified and practical approaches in pre-clinical, clinical and laboratory studies. *Biochemia Medica*, 31(1), 27–53. <https://doi.org/10.11613/BM.2021.010502>
- Sethares, K. A., & Chin, E. (2021). Age and gender differences in physical heart failure symptom clusters. *Heart & Lung*, 50(6), 832–837. <https://doi.org/10.1016/j.hrtlng.2021.07.001>
- Severo, M., Gaio, R., Lourenço, P., Alvelos, M., Bettencourt, P., & Azevedo, A. (2011). Indirect calibration between clinical observers-application to the New York Heart Association functional classification system. *BMC Research Notes*, 4(1), Article 276. <https://doi.org/10.1186/1756-0500-4-276>
- Sevilla-Cazes, J., Ahmad, F. S., Bowles, K. H., Jaskowiak, A., Gallagher, T., Goldberg, L. R., Kangovi, S., Alexander, M., Riegel, B., Barg, F. K., & Kimmel, S. E. (2018). Heart failure home management challenges and reasons for readmission: A qualitative study to understand the patient's perspective. *Journal of General Internal Medicine*, 33(10), 1700–1707. <https://doi.org/10.1007/s11606-018-4542-3>

- Sharma, A., Colvin-Adams, M., & Yancy, C. W. (2014). Heart failure in African Americans: Disparities can be overcome. *Cleveland Clinical Journal of Medicine*, *81*(5), 301–311. http://www.abccardio.org/wp-content/uploads/2016/11/Yancy_Colvin.pdf
- Sharma, V., Zehtabchi, S., Rojas, N., & Birkhahn, R. (2009). Ethnic variations in quality of life and depressive symptoms among Black Americans with acute decompensated heart failure. *Journal of the National Medical Association*, *101*(10), 985–991. [https://doi.org/10.1016/S0027-9684\(15\)31064-6](https://doi.org/10.1016/S0027-9684(15)31064-6)
- Shavers, V. L. (2007). Measurement of socioeconomic status in health disparities research. *Journal of the National Medical Association*, *99*(9), 1013–1023. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2575866/>
- Shen, B.-J., Eisenberg, S. A., Maeda, U., Farrell, K. A., Schwarz, E. R., Penedo, F. J., Bauerlein, E. J., & Mallon, S. (2010). Depression and anxiety predict decline in physical health functioning in patients with heart failure. *Annals of Behavioral Medicine*, *41*(3), 373–382. <https://doi.org/10.1007/s12160-010-9251-z>
- Sidell, N. L. (1997). Adult adjustment to chronic illness: A review of the literature. *Health & Social Work*, *22*(1), 5–11. <https://doi.org/10.1093/hsw/22.1.5>
- Smith, G. L., Shlipak, M. G., Havranek, E. P., Masoudi, F. A., McClellan, W. M., Foody, J. M., Rathore, S. S., & Krumholz, H. M. (2005). Race and renal impairment in heart failure: Mortality in Blacks versus Whites. *Circulation*, *111*(10), 1270–1277. <https://doi.org/10.1161/01.CIR.0000158131.78881.D5>
- Sokoreli, I., Vries, J., Pauws, S., & Steyerberg, E. (2016). Depression and anxiety as predictors of mortality among heart failure patients: Systematic review and meta-analysis. *Heart Failure Review*, *21*(1), 49–63. <https://doi.org/10.1007/s10741-015-9517-4>

- Spahillari, A., Talegawkar, S., Correa, A., Carr, J. J., Terry, J. G., Lima, J., Freedman, J. E., Das, S., Kociol, R., de Ferranti, S., Mohebbi, D., Mwasongwe, S., Tucker, K. L., Murthy, V. L., & Shah, R. V. (2017). Ideal cardiovascular health, cardiovascular remodeling, and heart failure in Blacks: The Jackson Heart Study. *Circulation: Heart Failure*, *10*(2). <https://doi.org/10.1161/CIRCHEARTFAILURE.116.003682>
- Stamp, K. D., Dunbar, S. B. R., Clark, P. C., Reilly, C. M., Gary, R. A., Higgins, M., & Kaslow, N. (2014). Family context influences psychological outcomes of depressive symptoms and emotional quality of life in patients with heart failure. *Journal of Cardiovascular Nursing*, *29*(6), 517–527. <https://doi.org/10.1097/JCN.0000000000000097>
- Steca, P., Greco, A., Monzani, D., Politi, A., Gestra, R., Ferrari, G., Malfatto, G., & Parati, G. (2013). How does illness severity influence depression, health satisfaction and life satisfaction in patients with cardiovascular disease? The mediating role of illness perception and self-efficacy beliefs. *Psychology & Health*, *28*(7), 765–783.
- Strine, T. W., Chapman, D. P., Balluz, L. S., Moriarty, D. G., & Mokdad, A. H. (2008). The associations between life satisfaction and health-related quality of life, chronic illness, and health behaviors among U.S. community-dwelling adults. *Journal of Community Health*, *33*(1), 40–50. <https://doi.org/10.1007/s10900-007-9066-4>
- Stronks, K., van de Mheen, H., van den Bos, J., & Mackenbach, J. P. (1997). The interrelationship between income, health and employment status. *International Journal of Epidemiology*, *26*(3), 592–600. <https://doi.org/10.1093/ije/26.3.592>

- Su, A., Al'Aref, S. J., Beecy, A. N., Min, J. K., & Karas, M. G. (2019). Clinical and socioeconomic predictors of heart failure readmissions: A review of contemporary literature. *Mayo Clinic Proceedings*, *94*(7), 1304–1320.
<https://doi.org/10.1016/j.mayocp.2019.01.017>
- Sullivan, L. T., Randolph, T., Merrill, P., Jackson, L. R., Egwim, C., Starks, M. A., & Thomas, K. L. (2018). Representation of Black patients in randomized clinical trials of heart failure with reduced ejection fraction. *The American Heart Journal*, *197*, 43–52.
<https://doi.org/10.1016/j.ahj.2017.10.025>
- Supino, P. G., Borer, J. S., Franciosa, J. A., Preibisz, J. J., Hochreiter, C., Isom, O. W., Krieger, K. H., Girardi, L. N., Bouraad, D. & Forur, L., 2009. Acceptability and psychometric properties of the Minnesota Living With Heart Failure Questionnaire among patients undergoing heart valve surgery: Validation and comparison with SF-36. *Journal of Cardiac Failure*, *15*(3), 267–277. <https://doi.org/10.1016/j.cardfail.2008.10.003>
- Tamaki, S., Yamada, T., Morita, T., Furukawa, Y., Iwasaki, Y., Kawasaki, M., Kikuchi, A., Kawai, T., Seo, M., Ikeda, I., Fukuhara, E., Abe, M., Nakamura, J., Kayama, K., Tanabe, K., Kawahira, M., & Fukunami, M. (2018). Prognostic value of charlson comorbidity index in patients admitted for acute decompensated heart failure with reduced or preserved left ventricular ejection fraction: A prospective study. *Journal of the American College of Cardiology*, *71*(11), Article A829. [https://doi.org/10.1016/S0735-1097\(18\)31370-6](https://doi.org/10.1016/S0735-1097(18)31370-6)

- Tandon, V. V., Stringer, B. B., Conner, C. C., Gabriel, A. A., Tripathi, B. B., Balakumaran, K. K., & Chen, K. K. (2020). An observation of racial and gender disparities in congestive heart failure admissions using the national inpatient sample. *Cureus*, *12*(10), Article e10914. <https://doi.org/10.7759/cureus.10914>
- Tandon, V., Gabriel, A., Balakumaran, K., Tandon, A., & Conner, C. (2019). Healthcare disparities in heart failure between African-Americans and Caucasian populations in the United States: A national review. *Chest*, *156*(4), Article A427. <https://doi.org/10.1016/j.chest.2019.08.457>
- Tansey, P. (2010). Counting the cost of heart failure to the patient, the nurse and the NHS. *British Journal of Nursing*, *19*(22), 1396–1401. <https://doi.org/10.12968/bjon.2010.19.22.1396>
- Tarekegn, G. E., Gezie, L. D., Birhan, T. Y., & Ewnetu, F. (2021). Health-related quality of life among heart failure patients attending an outpatient clinic in the University of Gondar Comprehensive Specialized Hospital Northwest, Ethiopia, 2020: Using structural equation modeling approach. *Patient Related Outcome Measures*, *12*, 279–290. <https://doi.org/10.2147%2FPROM.S322421>
- Thurston, R. C., Blumenthal, J. A., Babyak, M. A., & Sherwood, A. (2006). Association between hot flashes, sleep complaints, and psychological functioning among healthy menopausal women. *International Journal of Behavioral Medicine*, *13*(2), 163–172. https://doi.org/10.1207/s15327558ijbm1302_8

- Trivedi, R. B., Blumenthal, J. A., O'Connor, C., Adams, K., Hinderliter, A., Dupree, C., Johnson, K., & Sherwood, A. (2009). Coping styles in heart failure patients with depressive symptoms. *Journal of Psychosomatic Research*, *67*(4), 339–346. <https://doi.org/10.1016/j.jpsychores.2009.05.014>
- Tsabedze, N., Kinsey, J.-L. H., Mpanya, D., Mogashoa, V., Klug, E., & Manga, P. (2021). The prevalence of depression, stress and anxiety symptoms in patients with chronic heart failure. *International Journal of Mental Health Systems*, *15*(1), Article 44. <https://doi.org/10.1186/s13033-021-00467-x>
- Tsao, C. W., Aday, A. W., Almarzooq, Z. I., Alonso, A., Beaton, A. Z., Bittencourt, M. S., Boehme, A. K., Buxton, A. E., Carson, A. P., Commodore-Mensah, Y., Elkind, M. S. V., Evenson, K. R., Eze-Nliam, C., Ferguson, J. F., Generoso, G., Ho, J. E., Kalani, R., Khan, S. S., Kissela, B. M., ... Martin, S. S. (2022). Heart disease and stroke statistics—2022 update: A report from the American Heart Association. *Circulation*, *145*(8), e153–e639. <https://doi.org/10.1161/CIR.0000000000001052>
- Vaccarino, V., Gahbauer, E., Kasl, S. V., Charpentier, P. A., Acampora, D., & Krumholz, H. M. (2002). Differences between African Americans and Whites in the outcome of heart failure: Evidence for a greater functional decline in African Americans. *American Heart Journal*, *143*(6), 1058–1067. <https://doi.org/10.1067/mhj.2002.122123>
- Vaccarino, V., Kasl, S. V., Abramson, J., & Krumholz, H. M. (2001). Depressive symptoms and risk of functional decline and death in patients with heart failure. *Journal of the American College of Cardiology*, *38*(1), 199–205. [https://doi.org/10.1016/S0735-1097\(01\)01334-1](https://doi.org/10.1016/S0735-1097(01)01334-1)

- Van Nuys, K. E., Xie, Z., Tysinger, B., Hlatky, M. A., & Goldman, D. P. (2018). Innovation in heart failure treatment: Life expectancy, disability, and health disparities. *JACC: Heart Failure*, 6(5), 401–409. <https://doi.org/10.1016/j.jchf.2017.12.006>
- Virani, S. S., Alonso, A., Aparicio, H. J., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Cheng, S., Delling, F. N., Elkind, M. S. V., Evenson, K. R., Ferguson, J. F., Gupta, D. K., Khan, S. S., Kissela, B. M., Knutson, K. L., Lee, C. D., Lewis, T. T., ... Tsao, C. W. (2021). Heart disease and stroke statistics—2021 update: A report from the American Heart Association. *Circulation*, 143(8), e254–e743. <https://doi.org/10.1161/CIR.0000000000000950>
- Virani, S. S., Alonso, A., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Chang, A. R., Cheng, S., Delling, F. N., Djousse, L., Elkind, M. S. V., Ferguson, J. F., Fornage, M., Khan, S. S., Kissela, B. M., Knutson, K. L., Kwan, T. W., Lackland, D. T., ... Tsao, C. W. (2020). Heart Disease and Stroke Statistics—2020 update: A report from the American Heart Association. *Circulation*, 141(9), e139–e151. <https://doi.org/10.1161/CIR.0000000000000757>
- von Leupoldt, A. P., Taube, K. M. D., Lehmann, K., Fritzsche, A. P., & Magnussen, H. M. D. (2011). The impact of anxiety and depression on outcomes of pulmonary rehabilitation in patients with COPD. *Chest*, 140(3), 730–736. <https://doi.org/10.1378/chest.10-2917>
- Waldron, I., Weiss, C. C., & Hughes, M. E. (1997). Marital status effects on health: Are there differences between never married women and divorced and separated women? *Social Science & Medicine*, 45(9), 1387–1397. [https://doi.org/10.1016/S0277-9536\(97\)00065-8](https://doi.org/10.1016/S0277-9536(97)00065-8)

- Wang, Z.-F., Cheng, Y.-C., Zhang, N.-H., Luo, R., Guo, K.-l., Ge, S.-W., & Xu, G. (2021). Effect of marital status on depression and mortality among patients with chronic kidney disease from National Health and Nutrition Examination Survey 2005–2014. *Kidney Diseases*, 7(5), 391–400. <https://doi.org/10.1159/000515440>
- Warnecke, R. B., Ferrans, C. E., Johnson, T. P., Chapa-Resendez, G., O'Rourke, D. P., Chávez, N., Dudas, S., Smith, E. D., Martínez Schallmoser, L., Hand, R. P., & Lad, T. (1996). Measuring quality of life in culturally diverse populations. *Monographs*(20), 29–38.
- Warner, C. B., Roberts, A. R., Jeanblanc, A. B., & Adams, K. B. (2019). Coping resources, loneliness, and depressive symptoms of older women with chronic illness. *Journal of Applied Gerontology*, 38(3), 295–322. <https://doi.org/10.1177/0733464816687218>
- Watkins, D. C., Assari, S., & Johnson-Lawrence, V. (2015). Race and ethnic group differences in comorbid major depressive disorder, generalized anxiety disorder, and chronic medical conditions. *Journal of Racial and Ethnic Health Disparities*, 2(3), 385–394. <https://doi.org/10.1007/s40615-015-0085-z>
- Watkins, T., Mansi, M., Thompson, J., Mansi, I., & Parish, R. (2013). Effect of marital status on clinical outcome of heart failure. *Journal of Investigative Medicine*, 61(5), 835–841. <https://doi.org/10.2310/JIM.0b013e31828c823e>
- Westlake, C., Sethares, K., & Davidson, P. (2013). How can health literacy influence outcomes in heart failure patients? Mechanisms and interventions. *Current Heart Failure Reports*, 10(3), 232–243. <https://doi.org/10.1007/s11897-013-0147-7>
- Whittemore, R., & Dixon, J. (2008). Chronic illness: The process of integration. *Journal of Clinical Nursing*, 17(7b), 177–187. <https://doi.org/10.1111/j.1365-2702.2007.02244.x>

- Wierenga, K. L., Dekker, R. L., Lennie, T. A., Chung, M. L., & Dracup, K. (2017). African American race is associated with poorer outcomes in heart failure patients. *Western Journal of Nursing Research*, 39(4), 524–538.
<https://doi.org/10.1177/0193945916661277>
- Wiley, J. F., Chan, Y.-K., Ahamed, Y., Ball, J., Carrington, M. J., Riegel, B., & Stewart, S. (2018). Multimorbidity and the risk of all-cause 30-day readmission in the setting of multidisciplinary management of chronic heart failure: A retrospective analysis of 830 hospitalized patients in Australia. *The Journal of Cardiovascular Nursing*, 33(5), 437–445. <https://doi.org/10.1097/JCN.0000000000000391>
- Williams, D. R., Takeuchi, D. T., & Adair, R. K. (1992). Marital status and psychiatric disorders among Blacks and Whites. *Journal of Health and Social Behavior*, 33(2), 140–157.
<https://doi.org/10.2307/2137252>
- Winkleby, M. A., Jatulis, D. E., Frank, E., & Fortmann, S. P. (1992). Socioeconomic status and health: How education, income, and occupation contribute to risk factors for cardiovascular disease. *American Journal of Public Health*, 82(6), 816–820.
<https://doi.org/10.2105/AJPH.82.6.816>
- Wong, C. Y., Chaudhry, S. I., Desai, M. M., & Krumholz, H. M. (2011). Trends in comorbidity, disability, and polypharmacy in heart failure. *The American Journal of Medicine*, 124(2), 136–143. <https://doi.org/10.1016/j.amjmed.2010.08.017>
- World Health Organization. (n.d.). *Gender and health*. https://www.who.int/health-topics/gender#tab=tab_1

- Wu, J. R., Lennie, T. A., Frazier, S. K., & Moser, D. K. (2016). Health-related quality of life, functional status and cardiac event-free survival in patients with heart failure. *The Journal of cardiovascular nursing*, 31(3), 236. Xiong, G. L., Prybol, K., Boyle, S. H., Hall, R., Streilein, R. D., Steffens, D. C., Krishnan, R., Rogers, J. G., O'Connor, C. M., & Jiang, W. (2015). Inflammation markers and major depressive disorder in patients with chronic heart failure: Results from the Sertraline Against Depression and Heart Disease in Chronic Heart Failure Study. *Psychosomatic Medicine*, 77(7), 808–815. <https://doi.org/10.1097/PSY.0000000000000216>
- Yancy, C. W. (2001). Heart failure in Blacks: Etiologic and epidemiologic differences. *Current Cardiology Reports*, 3(3), 191–197. <https://doi.org/10.1007/s11886-001-0022-0>
- Yohannes, A. M., Willgoss, T. G., Baldwin, R. C., & Connolly, M. J. (2010). Depression and anxiety in chronic heart failure and chronic obstructive pulmonary disease: Prevalence, relevance, clinical implications and management principles. *International Journal of Geriatric Psychiatry*, 25(12), 1209–1221. <https://doi.org/10.1002/gps.2463>
- Zhang, A.-Z., Wang, Q.-C., Huang, K.-M., Huang, J.-G., Zhou, C.-H., Sun, F.-Q., Wang, S.-W., & Wu, F.-T. (2016). Prevalence of depression and anxiety in patients with chronic digestive system diseases: A multicenter epidemiological study. *World Journal of Gastroenterology*, 22(42), 9437–9444. <https://doi.org/10.3748/wjg.v22.i42.9437>
- Zhang, X., Zou, H., Hou, D., He, D., & Fan, X. (2020). Functional status mediates the association of nutritional status with depressive symptoms in patients with heart failure. *Journal of Advanced Nursing*, 76(12), 3363–3371. <https://doi.org/10.1111/jan.14522>

Ziaean, B., Kominski, G. F., Ong, M. K., Mays, V. M., Brook, R. H., & Fonarow, G. C. (2017).

National differences in trends for heart failure hospitalizations by sex and race/ethnicity.

Circulation: Cardiovascular Quality and Outcomes, 10(7).

<https://doi.org/10.1161/CIRCOUTCOMES.116.003552>

CHAPTER 3: METHODOLOGY

Introduction

The purpose of this descriptive, comparative study was to explore the relationships between sociodemographic characteristics (race, age, gender, marital status, education, and employment status), clinical characteristics (disease severity, comorbidities, and functional status), psychological characteristics (depression and anxiety), and quality of life (QOL) in patients with heart failure (HF) and to determine if there are racial differences between Blacks and Whites. The research question was: What are the differences between Blacks and Whites with HF with respect to sociodemographic and clinical characteristics and quality of life? The specific aims were to (a) describe the differences and examine the relationships between sociodemographic, clinical, and psychological characteristics, and QOL in Blacks and Whites persons with HF and (b) examine the influence of sociodemographic, clinical, and psychological characteristics on QOL in Black and White persons with HF.

Like several prior studies, Herr et al. (2015; the parent study) did not aim to identify differences in depression and anxiety by race in patients with HF. Hence, a secondary analysis of previously collected data was employed. Exploring whether there are racial differences in depression and anxiety based on these variables in a geographically homogeneous sample will add to the limited research that suggests that there might be racial differences and that other sociodemographic, clinical, and psychological characteristics might also influence these relationships. Findings from this study could serve as foundational research for future studies that explore racial differences in depression and anxiety in Black patients with HF. Ultimately, learning more about racial differences, if they exist, could help to decrease clinical outcome disparities among Blacks. Following is a discussion of the research approach and the rationale

for its choice, followed by the sample, setting, data collection procedures, variables and instruments used for measurement, and a description of the statistical analysis plan.

Rationale for Conducting a Secondary Data Analysis

Given that all data and perspectives are usually not explored in primary research studies, investigators considering secondary data analysis apply theoretical knowledge and conceptual skills to design studies with new research questions that are supportable by existing knowledge and gaps in knowledge (Johnston, 2017). Analyzing existing data helps researchers reduce costs, improve feasibility, and inflict less participant burden (Dunn et al., 2015). Secondary data analysis also allows researchers to answer important questions that generate new knowledge while posing less risk to vulnerable, hard-to-reach, and over-researched groups such as Blacks and patients with cardiovascular diseases (Dunn et al., 2015; O'Connor, 2020).

Setting, Sample, Data Collection, and Human Subjects Protection

The present study was a secondary analysis of the data from Herr et al.'s (2015) cross-sectional study. The parent study occurred within a heart failure/transplant program at Virginia Commonwealth University Medical Center in Richmond, Virginia, to identify symptom clusters and evaluate relationships between identified clusters and functional status (Herr et al., 2015). A subsequent study of the parent study investigated HF symptom clusters and their influence on QOL (Salyer et al., 2019). While conducting the parent study, Herr et al. recruited a convenience sample ($N = 117$) during a scheduled follow-up clinic visit. Potential participants entered a private examination room, where advanced practice nurses and cardiologists screened them for inclusion using the established participant eligibility criteria: (a) confirmed diagnosis of HF, (b) age 18 or older, (c) NYHA Class II to IV, and (d) receiving standard treatment for HF (beta blocker, diuretics, or ACE inhibitor therapy). Exclusion criteria were: (a) diagnosis of dementia,

(b) the presence of a ventricular assist device, or (c) the presence of a total artificial heart. After confirming participant eligibility, a research assistant explained the study to potential participants and gave each eligible patient a study packet that included a cover letter, consent form, and study questionnaires. After obtaining informed consent, study staff reviewed the questionnaire packet with each participant and answered questions. Participants then spent about 30 minutes completing the self-report questionnaires, which they returned to the staff (Herr et al., 2015).

The medical records provided participants' ages. Five the 151 eligible patients declined to participate. Of the questionnaires provided to 146 eligible participants, 29 were incomplete. Only complete questionnaires were included in the analysis, leaving a sample of 117 participants' data in the parent study (Herr et al., 2015). To address the aims of the current study, the sample for this secondary analysis ($N = 114$) includes the completed data of patients who racially self-identified as Black ($n = 59$) or White ($n = 55$). Patients who self-identified as American Indian or Alaska Native ($n = 3$) were excluded from this analysis.

Virginia Commonwealth University's Institutional Review Board (IRB) provided approval prior to conducting the parent study (Herr et al., 2015; Salyer et al., 2019). The primary investigators deidentified the data set used for this secondary study. The secondary study proceeded after obtaining Virginia Commonwealth University IRB approval.

Study Variables and Measures

Sociodemographic Characteristics

Sociodemographic data were self-reported by participants and collected using a researcher-developed questionnaire. The sociodemographic characteristics examined in this study were race, age, gender, marital status, education, and employment status. In this study, race was a categorical variable with two levels (Black/White), age was a continuous variable, gender

was a categorical variable with two levels (male/female), and marital status was a categorical variable with three levels (never married, previously married, and currently married). Education was a categorical variable with three levels (did not complete high school, high school graduate, and college graduate), and employment status was a categorical variable with eight levels (unemployed, employed part-time, employed full time, homemaker, receiving disability, retired, student, and volunteer).

Clinical Characteristics

Clinical characteristics came from each participant's medical record. The three clinical characteristics investigated in this study were disease severity, comorbidities, and functional status. Left ventricular EF and NYHA Class were the measures of HF disease severity used, with the EF serving as the primary measure of disease severity with a predictive value. EF is a continuous variable expressed as a percentage, which ranges from 0–100%. A normal EF ranges from 50–70%, and an EF of less than 40% is consistent with a diagnosis of HF (American Heart Association, 2017b). The NYHA Class was the second measure of HF disease severity, used descriptively. NYHA Class includes four classes (Levels I–IV) that coincide with patients' symptom severity and related activity limitations. Class I has with no physical activity limitations, and Class II involves slight physical activity limitations. Patients classified at Level II are comfortable at rest, but ordinary activity results in symptoms such as fatigue, palpitations, and dyspnea. Class III involves marked physical activity limitations. Although patients in this group are comfortable at rest, limited activity causes the same as Class II. Class IV involves severe limitations, with patients experiencing HF symptoms at rest (American Hospital Association, 2017a). In this study, NYHA Class was a categorical variable with three levels (II–IV). There were no NYHA Class I patients included.

The Charlson Comorbidity Index (Charlson et al., 1987) was the measurement used to summarize co-occurring conditions. The method provides a weighted taxonomy of comorbid disorders and accompanying prognostic consequences to estimate the risk of death based on an individual's number and type of comorbidities. The Charlson Comorbidity Index received extensive use in research conducted on Black individuals with chronic illnesses (Braithwaite et al., 2009; Hill-Briggs et al., 2002; Rattanasompattikul et al., 2011; Schoenthaler et al., 2009) and among the HF patient population (Dickson et al., 2011; Formiga et al., 2018; Shuvy et al., 2020; Siabani et al., 2016; Tamaki et al., 2018).

Measuring functional status, characterized as physical functioning, entailed using a battery of items from the physical functioning subscale of the Medical Outcomes Survey (MOS) 36-Item Short Form (SF-36). The SF-36 questionnaire was an adaptation of the longer instruments used to collect data in the MOS, which was an observational study of variations on physician practice styles and patient outcomes in different healthcare delivery systems. In its original form, MOS was appropriate for respondents 14 years of age or older and is either self-administered or administered by a trained interviewer.

In this study, the conceptualization of functional status was as functional limitations. The physical functioning items ($n = 10$) assess the degree to which an individual's health status limits the ability to perform basic physical activities of daily living (ADLs), such as bathing, dressing, lifting, or carrying groceries. Respondents choose from three responses (1 = *limited a lot*, 2 = *limited a little*, and 3 = *not limited at all*), with a score for the physical functioning subscale subsequently generated by recoding and then averaging the 10 SF-36 items that comprise the physical functioning subscale. The physical functioning subscale is a valid and reliable measure of physical functioning, with a Cronbach's alpha of 0.93 (Hays et al., 1995).

The SF-36, including the physical functioning subscale, has received extensive use to measure functioning in chronically ill patients (Bjorner et al., 2013; Del Core et al., 2018; Fukuhara et al., 2006; Padilla et al., 2008; Yarlal et al., 2011), Black individuals with chronic illnesses (Hill-Briggs et al., 2002; Kusek et al., 2002; Montague & Perchonok, 2012; Porter et al., 2014), patients with cardiovascular illnesses (Aydemir et al., 2005; Konerman et al., 2011; Stefanick et al., 2016), including those with HF (Edelmann et al., 2011; Supino et al., 2009), and Black individuals with HF (de Leon et al., 2009; Gottlieb et al., 2004; Jayadevappa et al., 2007).

Psychological Characteristics

The Hospital Anxiety and Depression Scale (HADS) was the instrument used to measure depression and anxiety. The HADS is a well-established measure to screen for clinically significant depression and anxiety that takes 2 to 5 minutes to complete. The 14-item questionnaire with two subscales, depression and anxiety. The depression and anxiety subscales include seven items each and with ratings on a 4-point Likert scale from 0 (*absence/occasionally/very seldom*) to 3 (*occurs most of the time/very definitely/as much as I ever did*), with total possible scores for each subscale ranging from 0–21. Higher total scores for either subscale indicate greater severity of depression and/or anxiety with a total score of 0–7 considered normal, a total score of 8–10 indicating borderline (mild to moderate) depression or anxiety, and a total score of 11 or greater on either subscale suggesting abnormal levels of depression or anxiety (Snaith, 2003; Zigmond & Snaith, 1983).

The HADS is a valid and reliable measure of depression and anxiety with a Cronbach's alpha of 0.81 (Stafford et al., 2007). The tool is useful to measure depression and anxiety in patients with chronic illnesses (Bener et al., 2011; Loosman et al., 2010; Marinus et al., 2002), chronically ill Blacks (Nelson et al., 2010; Snigdha Alur-Gupta et al., 2021; Weiss et al., 2011)

and patients with cardiovascular illnesses (Matsuda et al., 2016; Roberts et al., 2001), including those with HF (Cameron et al., 2008; Cubbon Gale et al., 2011; Gaston-Johansson et al., 2013; Roberts et al., 2001; Yuen et al., 2011), and Black individuals with HF (Bakitas et al., 2020).

Quality of Life

In this study, the Quality of Life Index Cardiac Version-IV (QLI) overall score was the instrument used to measure QOL and the scores from each subscale to assess four unique domains of QOL. The QLI incorporates a common set of items across the general population as well as illness-specific versions with specific items developed for various disorders, including the version used in this study developed for use in patients with cardiovascular illnesses. The QLI is a self-administered two-part questionnaire, written at the fourth-grade reading level, takes approximately 10 minutes to administer and includes four subscales: (a) Health and Functioning, (b) Social and Economic, (c) Psychological and Spiritual, and (d) Family. The Health and Functioning subscale consists of 15 items that elicit respondents' subjective views of physical health status, ability to function in various activities and social roles, stress, concerns, worries, the ability to travel and participate in leisurely activities, and the potential for a happy old age/retirement. The Social and Economic subscale (seven items) is aligned with socioeconomic factors, including social support, friendships, neighborhood composition, education, employment status, one's standard of living, financial independence and includes concerns related to inflation, and economic stress. The seven items in the Psychological and Spiritual subscale are related to cognitive, emotional, and spiritual responses to various aspects of life, including peace of mind, faith in God, goal achievement, personal appearance, self-image, happiness, and general life satisfaction. Finally, the Family subscale includes five items related to family, including

satisfaction with a significant other, children, emotional support from family, and overall family health and happiness (Ferrans & Powers, 1985, 1992).

The QLI questionnaire includes two sections (Satisfaction and Importance) with 35 items per section (total of 70 responses required). The first section (Satisfaction) uses a 6-point Likert scale (1 = *very dissatisfied*, 2 = *moderately dissatisfied*, 3 = *slightly dissatisfied*, 4 = *slightly satisfied*, 5 = *moderately satisfied*, and 6 = *very satisfied*) and measures the respondents' satisfaction with various aspects of life. The second section (Importance) is a measure of the importance of the aspects of life assessed using a 6-point Likert scale (1 = *very unimportant*, 2 = *moderately unimportant*, 3 = *slightly unimportant*, 4 = *slightly important*, 5 = *moderately important*, and 6 = *very important*). Five scores are calculated from this questionnaire, one for each of the four QOL domains (subscales) and an overall QOL score combining each subscale.

The Importance items in the second QLI section are means to weigh the respondents' satisfaction of various aspects of life, resulting in scores reflecting respondents' satisfaction with the aspects of life they value most. Items rated as more important to a respondent have a greater impact on the QOL score than those deemed less important. Only the Satisfaction aspect of the QLI was reported in this study because it was the only aspect of the QLI measured in the parent study.

The QLI is a reliable and valid measure of QOL. Confirming its content validity occurred via a development process that included an extensive literature review of QOL, firsthand patient reports regarding the quality of their lives, and the Content Validity Index. Assessing construct validity entailed using the contrasted groups approach and factor analysis, resulting in the four factors underlying the QLI explaining 91% of the total variance. Supporting convergent validity were strong correlations between the QLI and Campbell et al.'s (1976) measure of life

satisfaction. The QLI has shown internal consistency with a Cronbach's alpha ranging from 0.73 to 0.99 across 48 studies (Ferrans, 1990; Ferrans & Powers, 1985; Ferrans and Powers Quality of Life Index; Ferrans and Powers Quality of Life Index, n.d.). Temporal reliability was assessed for the overall scale, and test-retest reliability was supported by correlations between 0.79 to 0.87 measured at 2-week and 1-month intervals (Dougherty et al., 1998; Ferrans & Powers, 1985) and of 0.78 at 3- to 4-week intervals (Rustoen et al., 1999). Each of the four subscales also showed temporal reliability by test-retest correlations at 2-week intervals: health and functioning ($r = .72$), social and economic ($r = .68$), psychological/spiritual ($r = .76$), and family ($r = .69$; Dougherty et al., 1998). Cronbach's alphas for the four subscales ranging from 0.63 to 0.96 have appeared in 24 studies. Finally, the QLI indicated sensitivity to change in 27 intervention studies (Ferrans, 1990; Ferrans & Powers, 1985; Ferrans and Powers Quality of Life Index, n.d.).

The QLI has received extensive use to measure QOL in chronic illness populations (Carroll et al., 1999; Delgado, 2007; DeSouza & Subrahmanya Nairy, 2003), chronically ill Black individuals (Greene, 2005; Johnson et al., 1998; Welch & Austin, 1999), and patients with cardiovascular illnesses (Flemme et al., 2001; Ghasemi et al., 2014; Grady et al., 1999), including those with HF (Koukouvou et al., 2004; Scott, 2000; Scott et al., 2004) and Black individuals with HF (de Leon et al., 2009).

Statistical Analysis

Data analysis entailed using SPSS 27 (IBM Corp., 2020). Calculating a score for each variable/instrument for each participant preceded descriptive statistics used to summarize the baseline sociodemographic characteristics, clinical characteristics, psychological characteristics, and QOL of the Black and the White samples. The Shapiro-Wilk test was the test used to assess for normality of the data (Ghasemi & Zahediasl, 2012). The level of significance was 0.05, and

the analysis required a minimum of 50 participants per group to have 80% power to detect differences between Blacks and Whites (Cohen, 1992; Serdar et al., 2021).

Aim 1

To describe the differences and examine the relationships between sociodemographic (age, gender, marital status, education, and employment status), clinical (disease severity, comorbidities, and functional status), and psychological (depression and anxiety) characteristics and QOL between Blacks and Whites with HF—Depending on the distribution of the data, a mean or median score was calculated for each sociodemographic, clinical, and psychological characteristic and QOL dichotomized by race (Black/White) and compared. Then, *t* tests (for continuous variables) and chi-square tests (for categorical variables) for independent samples were the statistics used to assess the mean or median for each measure.

To examine the relationships between or among sociodemographic (age, gender, marital status, education, and employment status), clinical (disease severity, comorbidities, and functional status), and psychological (depression and anxiety) characteristics, and QOL in Blacks and Whites with HF—To examine the relationships between or among these variables, Pearson’s correlation (for normally distributed data) or Spearman’s correlation (for data that do not meet parametric test assumptions) were used to determine significant relationships.

Aim 2

To examine the influence of sociodemographic (race, age, and gender), clinical (disease severity, comorbidities, and functional status), and psychological (depression and anxiety) characteristics on QOL in Blacks and Whites with HF—To examine the influence of the independent variables (psychological, sociodemographic, and clinical characteristics) that influence the dependent variable (QOL) in Blacks and Whites with HF, the scores of

psychological (depression and anxiety), and certain sociodemographic (age, race, and gender), clinical (disease severity, comorbidities, and functional status) characteristics were added to a regression model and a stepwise model-building approach used to determine the most significant predictors of QOL in Blacks for comparison with White patients with HF while trimming the model based on nonsignificant variables.

Limitations and Delimitations

With this study conducted in an outpatient clinical setting of one academic medical center and a small sample size, the findings might not be generalizable to the general HF patient population. Second, the variables examined were limited to those available in the dataset, although other variables might have been appropriate to investigate in a prospective study. Third, the parent study included patients who did not self-identify as Black or White race; however, they were excluded from this study, which aimed to compare characteristics of only Black and White racial groups. A fourth potential limitation was that researchers in the parent study included only participants with NYHA Class II to IV, thus excluding those with a NYHA Class of I, which could have influenced findings. Thus, comparisons to patients with NYHA Class I was not possible.

Another limitation was that the Satisfaction aspect of the QLI was the only subscale collected and reported in the parent study, and the Importance aspect was not available for analysis. The availability of this Importance data might have influenced findings because QOL is culturally influenced and Black individuals might weigh various aspects of their QOL differently than Whites (Blake & Darling, 2000; Matthews et al., 2012). Finally, the parent study occurred from 2011 to 2012 (Herr et al., 2015). Findings might vary from data collected today due to the elapsed time and changes in HF patient care and treatment that might influence racial disparities

among patients with HF. Despite these limitations, few researchers have investigated depression and anxiety in patients with HF and assessed for racial differences. This study provided foundational research contributing to the existing body of knowledge and informing future research directions regarding potential differences between Blacks and Whites with HF.

References

- American Heart Association. (2017a). *Classes of heart failure*. <https://www.heart.org/en/health-topics/heart-failure/what-is-heart-failure/classes-of-heart-failure>
- American Heart Association. (2017b). *Ejection fraction heart failure measurement*. <https://www.heart.org/en/health-topics/heart-failure/diagnosing-heart-failure/ejection-fraction-heart-failure-measurement>
- Aydemir, O., Ozdemir, C., & Koroglu, E. (2005). The impact of co-morbid conditions on the SF-36: A primary-care-based study among hypertensives. *Archives of Medical Research*, 36(2), 136–141. <https://doi.org/10.1016/j.arcmed.2004.12.009>
- Bakitas, M., Ejem, D., Dionne-Odom, J., Wells, R., Azuero, A., Stockdill, M., Keebler, K., Sockwell, E., Tims, S., Engler, S., Kvale, E., Tucker, R., Burgio, K., Tallaj, J., Swetz, K., Steinhauser, K., Durant, R., & Pamboukian, S. (2020). Educate, nurture, advise, before life ends: Comprehensive heart care for patients and families (ENABLE CHF-PC): An early palliative care telehealth randomized trial for patients with advanced heart failure (103A). *Journal of Pain and Symptom Management*, 60(1), 192. <https://doi.org/10.1016/j.jpainsymman.2020.04.033>
- Bener, A., Al-Hamaq, A. O. A. A., & Dafeeah, E. E. (2011). High prevalence of depression, anxiety and stress symptoms among diabetes mellitus patients. *The Open Psychiatry Journal*, 5(1), 5–12.
- Bjorner, J. B., Wolden, M. L., Gundgaard, J., & Miller, K. A. (2013). Benchmarks for interpretation of score differences on the SF-36 health survey for patients with diabetes. *Value in Health*, 16(6), 993–1000. <https://doi.org/10.1016/j.jval.2013.06.022>

- Blake, W. M., & Darling, C. A. (2000). Quality of life: Perceptions of African Americans. *Journal of Black Studies*, 30(3), 411–427. <https://doi.org/10.1177/002193470003000307>
- Braithwaite, D., Tammemagi, C. M., Moore, D. H., Ozanne, E. M., Hiatt, R. A., Belkora, J., West, D. W., Satariano, W. A., Liebman, M., & Esserman, L. (2009). Hypertension is an independent predictor of survival disparity between African-American and White breast cancer patients. *International Journal of Cancer*, 124(5), 1213–1219. <https://doi.org/10.1002/ijc.24054>
- Cameron, I. M., Crawford, J. R., Lawton, K., & Reid, I. C. (2008). Psychometric comparison of PHQ-9 and HADS for measuring depression severity in primary care. *British Journal of General Practice*, 58(546), 32–36. <https://doi.org/10.3399/bjgp08X263794>
- Cubbon, Gale, C. P., Kearney, L. C., Schechter, C. B., Brooksby, W. P., Nolan, J., Fox, K. A. A., Rajwani, A., Baig, W., Groves, D., Barlow, P., Fisher, A. C., Batin, P. D., Kahn, M. B., Zaman, A. G., Shah, A. M., Byrne, J. A., Lindsay, S. J., Sapsford, R. J., ... Kearney, M. T. (2011). Changing Characteristics and Mode of Death Associated With Chronic Heart Failure Caused by Left Ventricular Systolic Dysfunction A Study Across Therapeutic Eras. *Circulation. Heart Failure*, 4(4), 396–403. <https://doi.org/10.1161/CIRCHEARTFAILURE.110.959882>
- Blishen. (1977). *The Quality of American Life: Perceptions, Evaluations and Satisfactions* Angus Campbell, Philip E. Converse and Willard L. Rodgers New York: Russell Sage Foundation, 1976, pp. xi, 583. *Canadian Journal of Political Science*, 10(4), 875–877. <https://doi.org/10.1017/S0008423900050976>

- Carroll, D. L., Hamilton, G. A., & McGovern, B. A. (1999). Changes in health status and quality of life and the impact of uncertainty in patients who survive life-threatening arrhythmias. *Heart & Lung*, 28(4), 251–260. [https://doi.org/10.1016/S0147-9563\(99\)70071-3](https://doi.org/10.1016/S0147-9563(99)70071-3)
- Charlson, M. E., Pompei, P., Ales, K. L., & MacKenzie, C. R. (1987). A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *Journal of Chronic Diseases*, 40(5), 373–383. [https://doi.org/10.1016/0021-9681\(87\)90171-8](https://doi.org/10.1016/0021-9681(87)90171-8)
- Cohen. (1992). A Power Primer. *Psychological Bulletin*, 112(1), 155–159. <https://doi.org/10.1037/0033-2909.112.1.155>
- de Leon, C. F., Grady, K. L., Eaton, C., Rucker-Whitaker, C., Janssen, I., Calvin, J., & Powell, L. H. (2009). Quality of life in a diverse population of patients with heart failure: Baseline findings from the Heart Failure Adherence and Retention Trial (HART). *Journal of Cardiopulmonary Rehabilitation and Prevention*, 29(3), 171–178. <https://doi.org/10.1097/HCR.0b013e31819a0266>
- Del Core, M. A., Ahn, J., Wukich, D. K., Liu, G. T., Lalli, T., VanPelt, M. D., & Raspovic, K. M. (2018). Gender differences on SF-36 patient-reported outcomes of diabetic foot disease. *International Journal of Lower Extremity Wounds*, 17(2), 87–93. <https://doi.org/10.1177/1534734618774664>
- Delgado, C. (2007). Sense of coherence, spirituality, stress and quality of life in chronic illness. *Journal of Nursing Scholarship*, 39(3), 229–234. <https://doi.org/10.1111/j.1547-5069.2007.00173.x>

- DeSouza, M. S., & Subrahmanya Nairy, K. (2003). Nursing intervention for the quality of life of diabetic adults. *Clinical Effectiveness in Nursing*, 7(2), 63–72.
[https://doi.org/10.1016/S1361-9004\(03\)00044-X](https://doi.org/10.1016/S1361-9004(03)00044-X)
- Dickson, Buck, H., & Riegel, B. (2011). A Qualitative Meta-Analysis of Heart Failure Self-Care Practices Among Individuals With Multiple Comorbid Conditions. *Journal of Cardiac Failure*, 17(5), 413–419. <https://doi.org/10.1016/j.cardfail.2010.11.011>
- Dougherty, C. M., Dewhurst, T., Nichol, W. P., & Spertus, J. (1998). Comparison of three quality of life instruments in stable angina pectoris: Seattle Angina Questionnaire, Short Form Health Survey (SF-36), and Quality of Life Index-Cardiac Version III. *Journal of Clinical Epidemiology*, 51(7), 569–575. [https://doi.org/10.1016/S0895-4356\(98\)00028-6](https://doi.org/10.1016/S0895-4356(98)00028-6)
- Dunn, S. L., Arslanian-Engoren, C., DeKoekkoek, T., Jadack, R., & Scott, L. D. (2015). Secondary data analysis as an efficient and effective approach to nursing research. *Western Journal of Nursing Research*, 37(10), 1295–1307.
<https://doi.org/10.1177/0193945915570042>
- Edelmann, F., Stahrenberg, R., Gelbrich, G., Durstewitz, K., Angermann, C. E., Düngen, H.-D., Scheffold, T., Zugck, C., Maisch, B., Regitz-Zagrosek, V., Hasenfuß, G., Pieske, B. M., & Wachter, R. (2011). Contribution of comorbidities to functional impairment is higher in heart failure with preserved than with reduced ejection fraction. *Clinical Research in Cardiology*, 100(9), 755–764. <https://doi.org/10.1007/s00392-011-0305-4>
- Ferrans, C. E. (1990). Development of a quality of life index for patients with cancer. *Oncology Nursing Forum*, 17(3 Suppl), 15–19.

- Ferrans, C. E., & Powers, M. J. (1985). Quality of Life Index: Development and psychometric properties. *Advances in Nursing Science*, 8(1), 15–24. <https://doi.org/10.1097/00012272-198510000-00005>
- Ferrans, C. E., & Powers, M. J. (1992). Psychometric assessment of the Quality of Life Index. *Research in Nursing & Health*, 15(1), 29–38. <https://doi.org/10.1002/nur.4770150106>
- Flemme, I., Bolse, K., Ivarsson, A., Jinhage, B.-M., Sandstedt, B., Edvardsson, N., & Fridlund, B. (2001). Life situation of patients with an implantable cardioverter defibrillator: A descriptive longitudinal study. *Journal of Clinical Nursing*, 10(4), 563–572. <https://doi.org/10.1046/j.1365-2702.2001.00494.x>
- Formiga, F., Moreno-Gonzalez, R., Chivite, D., Franco, J., Montero, A., & Corbella, X. (2018). High comorbidity, measured by the Charlson Comorbidity Index, associates with higher 1-year mortality risks in elderly patients experiencing a first acute heart failure hospitalization. *Aging Clinical and Experimental Research*, 30(8), 927–933. <https://doi.org/10.1007/s40520-017-0853-1>
- Fukuhara, S., Yamazaki, S., Marumo, F., Akiba, T., Akizawa, T., Fujimi, S., Haruki, S., Kawaguchi, Y., Nihei, H., Shoji, T., & Maeda, T. (2006). Health-related quality of life of predialysis patients with chronic renal failure. *Nephron Clinical Practice*, 105(1), c1–c8. <https://doi.org/10.1159/000096802>
- Gaston-Johansson, F., Haisfield-Wolfe, M. E., Reddick, B., Goldstein, N., & Lawal, T. A. (2013, March). The relationships among coping strategies, religious coping, and spirituality in African American women with breast cancer receiving chemotherapy. *Oncology Nursing Forum*, 40(2).

- Ghasemi, E., Aliha, J. M., Bastani, F., Haghani, H., & Samiei, N. (2014). Quality of life in women with coronary artery disease. *Iranian Red Crescent Medical Journal*, *16*(7), Article e10188. <https://doi.org/10.5812/ircmj.10188>
- Ghasemi, A., & Zahediasl, S. (2012). Normality tests for statistical analysis: A guide for non-statisticians. *International Journal of Endocrinology and Metabolism*, *10*(2), 486–489. <https://doi.org/10.5812/ijem.3505>
- Gottlieb, S. S., Khatta, M., Friedmann, E., Einbinder, L., Katzen, S., Baker, B., Marshall, J., Minshall, S., Robinson, S., Fisher, M. L., Potenza, M., Sigler, B., Baldwin, C., & Thomas, S. A. (2004). The influence of age, gender, and race on the prevalence of depression in heart failure patients. *Journal of the American College of Cardiology*, *43*(9), 1542–1549. <https://doi.org/10.1016/j.jacc.2003.10.064>
- Grady, K. L., Jalowiec, A., & White-Williams, C. (1999). Predictors of quality of life in patients at one year after heart transplantation. *The Journal of Heart and Lung Transplantation*, *18*(3), 202–210. [https://doi.org/10.1016/S1053-2498\(98\)00048-5](https://doi.org/10.1016/S1053-2498(98)00048-5)
- Greene, R. A. (2005). Using the Ferrans and Powers Quality of Life Index of dialysis: A comparison of quality of life in older and younger African Americans receiving hemodialysis. *Topics in Geriatric Rehabilitation*, *21*(3), 230–232. <https://doi.org/10.1097/00013614-200507000-00009>
- Hays, R., Sherbourne, C., & Mazel, R. (1995). *User's manual for the Medical Outcomes Study (MOS) core measures of health-related quality of life*. Rand Corporaion.
- Herr, J. K., Salyer, J., Flattery, M., Goodloe, L., Lyon, D. E., Kabban, C. S., & Clement, D. G. (2015). Heart failure symptom clusters and functional status: A cross-sectional study. *Journal of Advanced Nursing*, *71*(6), 1274–1287. <https://doi.org/10.1111/jan.12596>

- Hill-Briggs, F., Gary, T. L., Hill, M. N., Bone, L. R., & Brancati, F. L. (2002). Health-related quality of life in urban African Americans with type 2 diabetes. *Journal of General Internal Medicine*, *17*(6), 412–419. <https://doi.org/10.1046/j.1525-1497.2002.11002.x>
- IBM Corp. (2020). IBM SPSS Statistics for Windows, Version 27.0.
- Jayadevappa, R., Johnson, J. C., Bloom, B. S., Nidich, S., Desai, S., Chhatre, S., Raziano, D. B., & Schneider, R. (2007). Effectiveness of transcendental meditation on functional capacity and quality of life of African Americans with congestive heart failure: A randomized control study. *Ethnicity and Disease*, *17*(1), 72–77. <https://doi.org/10.13016/v02r-skv3>
- Johnson, C. D., Wicks, M. N., Milstead, J., Hartwig, M., & Hathaway, D. K. (1998). Racial and gender differences in quality of life following kidney transplantation. *Image: The Journal of Nursing Scholarship*, *30*(2), 125–130. <https://doi.org/10.1111/j.1547-5069.1998.tb01266.x>
- Johnston, M. P. (2017). Secondary data analysis: A method of which the time has come. In *Qualitative and Quantitative Methods in Libraries*, *3*, 619–626.
- Konerman, M., Weeks, K. R., Shands, J. R., Tilburt, J. C., Dy, S., Bone, L. R., Levine, D. M., & Young, J. H. (2011). Short Form (SF-36) health survey measures are associated with decreased adherence among urban African Americans with severe, poorly controlled hypertension. *The Journal of Clinical Hypertension*, *13*(5), 385–390. <https://doi.org/10.1111/j.1751-7176.2010.00402.x>

- Koukouvou, G., Kouidi, E., Iacovides, A., Konstantinidou, E., Kaprinis, G., & Deligiannis, A. (2004). Quality of life, psychological and physiological changes following exercise training in patients with chronic heart failure. *Journal of Rehabilitation Medicine*, 36(1), 36–41. <https://doi.org/10.1080/11026480310015549>
- Kusek, J. W., Greene, P., Wang, S.-R., Beck, G., West, D., Jamerson, K., Agodoa, L. Y., Faulkner, M., & Level, B. (2002). Cross-sectional study of health-related quality of life in African Americans with chronic renal insufficiency: The African American Study of Kidney Disease and Hypertension Trial. *American Journal of Kidney Diseases*, 39(3), 513–524. <https://doi.org/10.1053/ajkd.2002.31401>
- Loosman, W. L., Siegert, C. E. H., Korzec, A., & Honig, A. (2010). Validity of the Hospital Anxiety and Depression Scale and the Beck Depression Inventory for use in end-stage renal disease patients. *British Journal of Clinical Psychology*, 49(4), 507–516. <https://doi.org/10.1348/014466509X477827>
- Marinus, J., Leentjens, A. F. G., Visser, M., Stiggelbout, A. M., & van Hilten, J. J. (2002). Evaluation of the hospital anxiety and depression scale in patients with Parkinson's disease. *Clinical Neuropharmacology*, 25(6), 318–324. <https://doi.org/10.1097/00002826-200211000-00008>
- Matsuda, R., Kohno, T., Kohsaka, S., Fukuoka, R., Maekawa, Y., Sano, M., Takatsuki, S., & Fukuda, K. (2016). The prevalence of poor sleep quality and its association with depression and anxiety scores in patients admitted for cardiovascular disease: A cross-sectional designed study. *International Journal of Cardiology*, 228, 977–982. <https://doi.org/10.1016/j.ijcard.2016.11.091>

- Matthews, A. K., Tejada, S., Johnson, T. P., Berbaum, M. L., & Manfredi, C. (2012). Correlates of quality of life among African American and White cancer survivors. *Cancer Nursing*, 35(5), 355–364. <https://doi.org/10.1097/NCC.0b013e31824131d9>
- Montague, E., & Perchonok, J. (2012). Health and wellness technology use by historically underserved health consumers: Systematic review. *Journal of Medical Internet Research*, 14(3), Article e2095. <https://doi.org/10.2196/jmir.2095>
- Nelson, C. J., Balk, E. M., & Roth, A. J. (2010). Distress, anxiety, depression, and emotional well-being in African-American men with prostate cancer. *Psycho-Oncology*, 19(10), 1052–1060. <https://doi.org/10.1002/pon.1659>
- O'Connor, S. (2020). Secondary data analysis in nursing research: A contemporary discussion. *Clinical Nursing Research*, 29(5), 279–284. <https://doi.org/10.1177/1054773820927144>
- Padilla, J., Krasnoff, J., DaSilva, M., Hsu, C.-Y., Frassetto, L., Johansen, K. L., & Painter, P. (2008). Physical functioning in patients with chronic kidney disease. *Journal of Nephrology*, 21(4), 550–559.
- Porter, A., Fischer, M. J., Wang, X., Brooks, D., Bruce, M., Charleston, J., Cleveland, W. H., Dowie, D., Faulkner, M., Gassman, J., Hiremath, L., Kendrick, C., Kusek, J. W., Norris, K. C., Thornley-Brown, D., Greene, T., & Lash, J. P. (2014). Quality of life and outcomes in African Americans with CKD. *Journal of the American Society of Nephrology*, 25(8), 1849–1855. <https://doi.org/10.1681/ASN.2013080835>
- Ferrans and Powers Quality of Life Index. (n.d.). *Reliability and validity*. <http://qli.org.uic.edu/reliability/reliabilityhome.htm>

- Rattanasompattikul, M., Feroze, U., Molnar, M. Z., Dukkipati, R., Kovesdy, C. P., Nissenson, A. R., Norris, K. C., Kopple, J. D., & Kalantar-Zadeh, K. (2011). Charlson Comorbidity Score is a strong predictor of mortality in hemodialysis patients. *International Urology and Nephrology*, 44(6), 1813–1823. <https://doi.org/10.1007/s11255-011-0085-9>
- Roberts, S. B., Bonnici, D. M., Mackinnon, A. J., & Worcester, M. C. (2001). Psychometric evaluation of the Hospital Anxiety and Depression Scale (HADS) among female cardiac patients. *British Journal of Health Psychology*, 6(4), 373–383. <https://doi.org/10.1348/135910701169278>
- Rustoen, T., Wiklund, I., Hanestad, B. R., & Burckhardt, C. S. (1999). Validity and reliability of the Norwegian version of the Ferrans and Powers Quality of Life Index. *Scandinavian Journal of Caring Sciences*, 13(2), 96–101. <https://doi.org/10.1080/02839319950162633>
- Salyer, J., Flattery, M., & Lyon, D. E. (2019). Heart failure symptom clusters and quality of life. *Heart & Lung*, 48(5), 366–372. <https://doi.org/10.1016/j.hrtlng.2019.05.016>
- Schoenthaler, A., Ogedegbe, G., & Allegrante, J. P. (2009). Self-efficacy mediates the relationship between depressive symptoms and medication adherence among hypertensive African Americans. *Health Education & Behavior*, 36(1), 127–137. <https://doi.org/10.1177/1090198107309459>
- Scott, L. D. (2000). Caregiving and care receiving among a technologically dependent heart failure population. *Advances in Nursing Science*, 23(2), 82–97. <https://doi.org/10.1097/00012272-200012000-00008>
- Scott, L. D., Setter-Kline, K., & Britton, A. S. (2004). The effects of nursing interventions to enhance mental health and quality of life among individuals with heart failure. *Applied Nursing Research*, 17(4), 248–256. [https://doi.org/10.1016/S0897-1897\(04\)00074-6](https://doi.org/10.1016/S0897-1897(04)00074-6)

- Serdar, C. C., Cihan, M., Yücel, D., & Serdar, M. A. (2021). Sample size, power and effect size revisited: Simplified and practical approaches in pre-clinical, clinical and laboratory studies. *Biochemia Medica*, 31(1), 27–53. <https://doi.org/10.11613/BM.2021.010502>
- Shuvy, M., Zwas, D. R., Keren, A., & Gotsman, I. (2020). The age-adjusted Charlson Comorbidity Index: A significant predictor of clinical outcome in patients with heart failure. *European Journal of Internal Medicine*, 73, 103–104. <https://doi.org/10.1016/j.ejim.2019.12.030>
- Siabani, S., Driscoll, T., Davidson, P. M., Najafi, F., Jenkins, M. C., & Leeder, S. R. (2016). Self-care and its predictors in patients with chronic heart failure in Western Iran. *The Journal of Cardiovascular Nursing*, 31(1), 22–30. <https://doi.org/10.1097/JCN.0000000000000211>
- Snaith, R. P. (2003). The Hospital Anxiety and Depression Scale. *Health and Quality of Life Outcomes*, 1(1), Article 29. <https://doi.org/10.1186/1477-7525-1-29>
- Alur-Gupta, S., Lee, I., Chemerinski, A., Liu, C., Lipson, J., Allison, K., Gallop, R., & Dokras, A. (2021). Racial differences in anxiety, depression, and quality of life in women with polycystic ovary syndrome. *F&S Reports*, 2(2), 230–237. <https://doi.org/10.1016/j.xfre.2021.03.003>
- Stafford, L., Berk, M., & Jackson, H. J. (2007). Validity of the Hospital Anxiety and Depression Scale and Patient Health Questionnaire-9 to screen for depression in patients with coronary artery disease. *General Hospital Psychiatry*, 29(5), 417–424. <https://doi.org/10.1016/j.genhosppsy.2007.06.005>

- Stefanick, M. L., Brunner, R. L., Leng, X., Limacher, M. C., Bird, C. E., Garcia, D. O., Hogan, P. E., LaMonte, M. J., Mackey, R. H., Johnson, K. C., LaCroix, A., Robinson, J. G., Seguin, R. A., Tindle, H. A., & Wassertheil-Smoller, S. (2016). The relationship of cardiovascular disease to physical functioning in women surviving to age 80 and above in the Women's Health Initiative. *The Journals of Gerontology: Series A*, *71*(3), S42–S53. <https://doi.org/10.1093/gerona/glv087>
- Supino, P. G., Borer, J. S., Franciosa, J. A., Preibisz, J. J., Hochreiter, C., Isom, O. W., Krieger, K. H., Girardi, L. N., Bouraad, D., & Forur, L. (2009). Acceptability and psychometric properties of the Minnesota Living with Heart Failure Questionnaire among patients undergoing heart valve surgery: Validation and comparison with SF-36. *Journal of Cardiac Failure*, *15*(3), 267–277. <https://doi.org/10.1016/j.cardfail.2008.10.003>
- Tamaki, S., Yamada, T., Morita, T., Furukawa, Y., Iwasaki, Y., Kawasaki, M., Kikuchi, A., Kawai, T., Seo, M., Ikeda, I., Fukuhara, E., Abe, M., Nakamura, J., Kayama, K., Tanabe, K., Kawahira, M., & Fukunami, M. (2018). Prognostic value of Charlson Comorbidity Index in patients admitted for acute decompensated heart failure with reduced or preserved left ventricular ejection fraction: A prospective study. *Journal of the American College of Cardiology*, *71*(11), Article A829. [https://doi.org/10.1016/S0735-1097\(18\)31370-6](https://doi.org/10.1016/S0735-1097(18)31370-6)
- Weiss, T., Weinberger, M. I., Holland, J., Nelson, C., & Moadel, A. (2011). Falling through the cracks: A review of psychological distress and psychosocial service needs in older Black and Hispanic patients with cancer. *Journal of Geriatric Oncology*, *3*(2), 163–173. <https://doi.org/10.1016/j.jgo.2011.12.001>

- Welch, J. L., & Austin, J. K. (1999). Quality of life in Black hemodialysis patients. *Advances in Renal Replacement Therapy*, 6(4), 351–357. [https://doi.org/10.1016/S1073-4449\(99\)70047-6](https://doi.org/10.1016/S1073-4449(99)70047-6)
- Yarlas, A. S., White, M. K., Yang, M., Saris-Baglana, R. N., Bech, P. G., & Christensen, T. (2011). Measuring the health status burden in hemodialysis patients using the SF-36 health survey. *Quality of Life Research*, 20(3), 383–389. <https://doi.org/10.1007/s11136-010-9764-8>
- Yuen, H. K., Holthaus, K., Kamen, D. L., Sword, D. O., & Breland, H. L. (2011). Using Wii Fit to reduce fatigue among African American women with systemic lupus erythematosus: A pilot study. *Lupus*, 20(12), 1293–1299. <https://doi.org/10.1177/0961203311412098>
- Zigmond, A. S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67(6), 361–370. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x>

CHAPTER 4: RESULTS

Introduction

This study addressed the differences and relationships between and the influence of sociodemographic, clinical, psychological characteristics, and the quality of life (QOL) on Black and White persons with heart failure (HF). This chapter presents the statistical and data analysis. The chapter also presents the univariate, bivariate, and multivariate analysis findings.

Data Analysis

Approach and Phases of Data Analysis

SPSS Version 27.0 (IBM Corp., 2020) was the software used to analyze the data in three phases. The first phase was descriptive analysis. All study variables were presented with descriptive statistics of means, standard deviations, minimum/maximum values for continuous variables (interval/ratio level), and frequencies and percentages for categorical variables (nominal/ratio level). The second phase was bivariate analysis, in which bivariate tests produced inferential statistics for the explanatory variables related to the dependent QOL variable at a statistically significant level ($p < .05$). Pearson's r zero-correlation indicated if the continuous explanatory variables significantly correlated to the continuous dependent variables. Independent-samples t tests were the means of measuring the differences between the continuous variables for Blacks and Whites. Finally, one-way ANOVA indicated if categorical explanatory variables with three or more categories significantly impacted QOL. The third phase, multivariate analysis, included the explanatory variables that correlated to the dependent variable at a statistically significant level ($p < .05$). Race was a design variable; therefore, it remained in each step of the regression model-building.

Data Preparation and Management

The original dataset included 114 participants (59 Black and 55 White). One participant did not self-identify a race and was excluded from the analysis. Due to a small amount of missing data, there was a plan to treat the missing data based on the missingness amount and patterns (Aguinis et al., 2013). The mean scale score was imputed for other missing data cases in the scales. The final data set did not include seven participants who did not provide sufficient data for the Quality of Life Index (QLI), QOL, and SF-36 functional status variables (at least 80% of items completed). Thus, the analysis did not include 5.4% ($n = 8$) participants due to missing data. The final sample included 104 patients with HF (51 Blacks and 53 Whites).

Creation of New Variables

The employment status variables in the parent study had eight categories. Due to small cell size by race, the employment status variable was collapsed into four categories: employed full-time, not employed full-time, receiving disability benefits, and retired. Initially, the marital status variable had three categories: never married, previously married, and currently married. However, this variable was collapsed into two categories, not currently married and currently married, to address associations with and the influence of married compared to unmarried status.

Data Distribution

The data distribution underwent examination to ensure each variable aligned with the parametric test assumptions of normality, linearity, homoskedasticity, multicollinearity, and outlier undue influence. The results showed an approximately normal score distribution due to skewness and kurtosis lower than three times the respective standard error (SE) of each value. Other than those indicated below, all data aligned with the parametric test assumptions and presented no issues. The Charlson Comorbidity Index had one outlier score (high = 11). The

HADS-D had one high outlier score of 12. The QLI had one outlier score (low = 48). The QOL Social/Economic Subscale had one outlier (low = 17). The Family Subscale had five outliers (low = 4, 4, 8, 9, 9).

Reliability

All the normed scales in this study showed sufficient internal consistency and reliability (Cronbach's $\alpha = \geq .70$; Bujang et al., 2018), including the QOL total scores (Cronbach's $\alpha = .96$), physical functioning scores (Cronbach's $\alpha = .90$), anxiety scores (Cronbach's $\alpha = .85$), and depression scores (Cronbach's $\alpha = .83$). The final multivariate analysis included the physical functioning scores to show the functional status construct. Cronbach's α also underwent examination for the QLI subscales and showed adequate internal consistency and reliability: Health/Functioning Subscale (Cronbach's $\alpha = .90$), Social and Economic Subscale (Cronbach's $\alpha = .88$), Psychological/Spiritual Subscale (Cronbach's $\alpha = .92$), and Family Subscale (Cronbach's $\alpha = .75$).

Statistical Power Analysis

Regarding statistical power, the G*Power software (Faul, Erdfelder, Buchner, & Lang, 2009; Faul, Erdfelder, Lang, & Buchner, 2007) indicated that a multiple linear regression model with seven explanatory variables, a medium effect size (Cohen's $f = .15$) between the explanatory and dependent variables, statistical power of $t .80$, and an alpha of $.05$ required a sample of 103 participants. Thus, the sample of 104 participants provided sufficient statistical power for analysis.

Descriptive Analysis

This section presents the sociodemographic, clinical, and psychological characteristics and QOL variables. The section also presents the characteristics of the overall sample and the

characteristics by race. This section addresses the mean comparisons by race and the bivariate and multivariate analysis results. The results showed an approximately normal score distribution due to skewness and kurtosis lower than approximately three times the respective standard error of each value.

Sociodemographic Characteristics

Table 4 presents a descriptive analysis of the categorical study variables. Table 5 presents the descriptive analysis of the continuous study variables. Table 4 includes the comparisons of the means of categorical variables by race. The overall mean participant age was 56.29 ($SD = 12.42$, min/max = 21.00–84.00). The sample had almost even division by race, with 51% ($n = 53$) White and 49.0% ($n = 51$) Black participants. The Black participants' ages ranged from 21 to 78, whereas the White participants' ages ranged from 40 to 84. There was a statistically significant difference between the Black and White participants' mean ages, $t(88.90) = 3.79$, $p < 0.001$). Men accounted for two-thirds of the overall sample ($n = 69$; 66.3%), with 42 White and 27 Black men. The female sample consisted of 11 White and 24 Black women. Race significantly related to gender, $\chi^2 = 8.05$, $p < .01$, as there was a higher percentage of Black ($n = 24$, 47.1%) than White women ($n = 11$, 20.8%; see Table 7). Over half the sample was married ($n = 59$, 56.7%). Race significantly associated with marital status, ($\chi^2 = 5.52$, $p < .05$), as more Whites were married ($n = 36$, 67.9%) than Blacks ($n = 23$, 45.1%; see Table 7).

Race significantly related to education level, $\chi^2 = 10.89$, $p < .01$, as more Black participants completed high school ($n = 34$, 66.7%) than White participants ($n = 19$, 35.8%). However, more White participants completed college ($n = 25$, 47.5%) than Black participants ($n = 10$, 19.6%). Regarding employment status, almost half the sample received disability ($n = 43$, 41.3%). Race did not significantly correlate with employment, $\chi^2 = 3.16$, $p = .37$.

Clinical Characteristics

The clinical characteristics examined by race included disease severity, comorbidity, and functional status; however, none showed statistical significance (see Table 6). The ejection fraction (EF) and New York Heart Association Classification (NYHA Class) were the means of measuring disease severity. The sample's mean EF ($n = 103$) was 24% ($SD = 9\%$, min/max = 10–40%). The mean difference in EF by race lacked statistical significance ($t = 1.21, p = .23$). The sample included 41 participants in NYHA Class II (21 Blacks and 20 Whites), 49 in NYHA Class III (25 Blacks and 24 Whites), and 14 in NYHA Class III (five Blacks and nine Whites). There were no statistically significant differences in mean NYHA Class ($t = .79, p = .43$) for comorbidities ($t = -0.02, p = .99$) or functional status by race ($t = -0.33, p = .74$).

Psychological Characteristics

The psychological characteristics of depression and anxiety were the variables measured with two subscales of the Hospital Anxiety and Depression Scale (HADS). Depression and anxiety highly correlated ($r = 0.651, p = <0.001$) in this sample. There were no statistically significant differences by race in the mean depression ($t = .25, p = .80$) or anxiety scores ($t = -1.15, p = .25$; Table 6).

When assessing the relationships between sociodemographic and psychological characteristics, the results showed higher depression levels ($r = -.211, p = .03$) and anxiety ($r = -.262, p = .007$) significantly correlated with lower education. Depression did not correlate with the remaining sociodemographic characteristics (see Table 8). Higher anxiety levels significantly correlated with age, as the younger participants reported more anxiety ($r = -.308, p = .001$).

Some clinical characteristics significantly correlated: NYHA Class and EF ($r = -.26, p = .007$), total comorbidity and functional status ($r = .34, p = <.001$), and NYHA Class and functional status ($r = .30, p = .002$). When assessing the relationships between clinical and psychological characteristics, higher NYHA Class (greater disease severity) correlated with higher levels of depression ($r = .251, p = .01$), suggesting that participants with more severe illnesses experienced more depression. Higher depression also significantly associated with lower functional status ($r = .582, p = <0.001$). However, depression did not significantly correlate with EF ($r = .008, p = 0.934$) or comorbidities ($r = .14, p = .078$). Anxiety significantly correlated with NYHA Class ($r = .262, p = .007$), suggesting that those with severe illnesses feel more anxious. Anxiety also significantly correlated with lower functional status ($r = .421, p = <0.001$) but not EF ($r = -.068, p = .493$) and comorbidities ($r = .102, p = .305$).

Quality of Life

There were no racial differences in overall QOL scores (see Table 5), even when assessing the four subscale scores for differences by race (see Table 6). The Health and Functioning Subscale score significantly correlated to age ($r = -.22, p = .03$), as the older participants had lower levels of health and functioning. Health and functioning did not significantly associate with the remaining sociodemographic characteristics of race ($r = .12, p = .22$); gender ($r = .07, p = .48$), marital status ($r = .48, p = .63$), educational status ($r = .10, p = .33$), and employment status ($r = -.12, p = .22$). The Health and Functioning Subscale did not correlate to the clinical characteristics of NYHA Class ($r = .14, p = .16$), EF ($r = -.10, p = .33$); comorbidities ($r = -.10, p = .31$), and functional status ($r = .03, p = .73$). The subscale also did not correlate with the psychological characteristics of depression ($r = .02, p = .88$) or anxiety ($r = .07, p = .49$).

The Social and Economic Subscale did not significantly correlate with the sociodemographic characteristics of race ($r = -0.2, p = .84$), age ($r = -.07, p = .49$), gender ($r = .09, p = .37$), marital status ($r = .05, p = .63$), education level ($r = .01, p = .91$); and employment status ($r = -.02, p = .84$). Additionally, the subscale did not correlate with the clinical characteristics of NYHA Class ($r = .24, p = .81$), EF ($r = -.03, p = .75$), comorbidities ($r = -.16, p = .12$), or functional status ($r = .10, p = .32$) or the psychological characteristics of depression ($r = .02, p = .82$) or anxiety ($r = -.16, p = .12$).

The Psychological/Spiritual Subscale did not significantly relate to the sociodemographic characteristics of race ($r = .83, p = .40$), age ($r = -.11, p = .27$), gender ($r = .02, p = .82$), marital status ($r = .10, p = .31$), educational status ($r = .10, p = .33$), and employment status ($r = -.07, p = .46$). The subscale did not correlate with the clinical characteristics of NYHA Class ($r = .07, p = .46$), EF ($r = -.06, p = .52$), comorbidities ($r = -.06, p = .52$), or functional status ($r = -.05, p = .63$). Additionally, the subscale did not correlate with the psychological characteristics of depression ($r = .01, p = .95$) or anxiety ($r = .05, p = .60$).

The Family Subscale did not significantly correlate with the sociodemographic characteristics of race ($r = -.03, p = .75$), age ($r = -.06, p = .53$), gender ($r = -.05, p = .60$), marital status ($r = .10, p = .31$), educational status ($r = .00, p = .33$), and employment status ($r = -.07, p = .46$). The subscale did not correlate with the clinical characteristics of NYHA Class ($r = -.07, p = .49$), EF ($r = -.02, p = .83$), comorbidities ($r = -.18, p = .06$), or functional status ($r = -.07, p = .47$). Additionally, the subscale did not correlate with the psychological characteristics of depression ($r = -.07, p = .46$) or anxiety ($r = -.12, p = .21$).

Table 1*Descriptive Analysis of Categorical Study Variables*

Variable	<i>n</i>	%
Gender		
Male	69	66.3
Female	35	33.7
Race		
White	53	51.0
Black	51	49.0
Education level		
Less than high school	16	15.4
Completed high school	30	28.8
Started college	23	22.1
Completed college	21	20.2
Education level (collapsed for analysis)		
Less than high school	16	15.4
High school	53	51.0
Completed college	35	33.7
Marital status		
Never married	20	19.2
Previously married	25	24.0
Currently married	59	56.7
Employment status		
Unemployed (but looking for work)	6	5.8
Employed part time	2	1.9
Employed full time	17	16.3
Homemaker	1	1.0
Receiving disability	43	41.3
Retired	32	30.8
Student	1	1.0
Volunteer	2	1.9
Employment status (collapsed for analysis)		
Employed full time	17	16.3
Not employed full time	12	11.5
Receiving disability	43	41.3
Retired	32	30.8

Note. *N* = 104

Table 2*Descriptive Analysis of Continuous Study Variables*

Variable	<i>M (SD)</i>	Minimum/maximum
Age	56.29 (12.30)	21.00–84.00
QOL (Total)	150.57 (33.34)	48.00–209.00
QOL-Health/Functioning	51.90 (16.37)	14.00–83.00
QOL-Social/Economic	43.15 (9.32)	17.00–64.00
QOL-Psych/Spiritual	35.76 (10.27)	10.00–48.00
QOL-Family	19.66 (4.38)	4.00–24.00
Physical Functioning	30.58 (25.39)	0.00–95.00
Ejection Fraction*	24% (9%)	10–40%
NYHA Class	2.74 (.68)	2.00–4.00
Comorbidity	3.15 (1.87)	0.00–8.00
Anxiety (HADS-A)	8.17 (4.66)	0.00–19.00
Depression (HADS-D)	7.54 (4.34)	0.00–19.00

Note. *N* = 104. **N* = 103

Table 3*Independent Samples t-Test Analysis Examining Racial Differences in Sociodemographic and Clinical Characteristics and Quality of Life*

Variable	n	<i>M (SD)</i>	<i>t (df)</i>	<i>p</i>
Age			3.82 (102)	<.001*
White	53	60.54 (9.37)		
Black	51	51.88 (13.47)		
QOL Total Scores			.13 (102)	.90
White	53	150.88 (28.73)		
Black	51	150.06 (37.84)		
QOL-Health/Functioning			-.57 (102)	.57
White	53	51.00 (14.84)		
Black	51	53.75 (15.50)		
QOL-Social/Economic			1.37 (102)	.17
White	53	44.38 (7.70)		
Black	51	41.88 (10.68)		

Variable	n	M (SD)	t (df)	p
QOL-Psych/Spiritual			-.48 (102)	.63
White	53	35.28 (9.23)		
Black	51	36.25 (11.33)		
QOL-Family			1.32 (102)	.19
White	53	20.22 (3.84)		
Black	51	19.08 (4.85)		
Physical Functioning			-.43 (102)	.67
White	53	29.53 (24.71)		
Black	51	31.68 (26.27)		
Ejection Fraction			1.21 (101)	.23
White	53	25% (9%)		
Black	51	23% (8%)		
NYHA Class			.79 (102)	.43
White	53	2.79 (.72)		
Black	51	2.69 (.65)		
Comorbidity			.131 (203)	.89
White	53	3.21 (2.24)		
Black	51	3.16 (1.67)		
Anxiety (HADS-A)			-1.15 (102)	.25
White	53	7.66 (4.72)		
Black	51	8.71 (4.58)		
Depression (HADS-D)			-.25 (102)	.80
White	53	7.43 (4.20)		
Black	51	7.64 (4.52)		

* $p < .05$, ** $p < .01$.

Table 4 presents the chi-square analyses of categorical sociodemographic characteristics by race.

Table 4*Chi-Square Analysis of Sociodemographic Characteristics by Race*

Gender						
		Male <i>n</i> (%)	Female <i>n</i> (%)	X^2 (<i>df</i>)	<i>p</i>	
Race				8.05 (1)	.005*	
White		42 (79.2)	11 (20.8)			
Black		27 (52.9)	24 (47.1)			
Marital status						
		Previously or never married <i>n</i> (%)	Currently married <i>n</i> (%)	<i>v</i>	<i>p</i>	
Race				5.52 (1)	.02*	
White		17 (32.1)	36 (67.9)			
Black		28 (54.9)	23 (45.1)			
Education level						
	< High school <i>n</i> (%)	High school <i>n</i> (%)	Completed college <i>n</i> (%)	X^2 (<i>df</i>)	<i>p</i>	
Race				10.89 (2)	.004*	
White	9 (17.0)	19 (35.8)	25 (47.2)			
Black	7 (13.7)	34 (66.7)	10 (19.6)			
Employment status						
	Employed full time <i>n</i> (%)	Not employed full time <i>n</i> (%)	Disability <i>n</i> (%)	Retired <i>n</i> (%)	X^2 (<i>df</i>)	<i>p</i>
Race					3.16 (3)	.37
White	10 (18.9)	4 (7.5)	20 (37.7)	19 (35.8)		
Black	7 (13.7)	8 (15.7)	23 (45.1)	13 (25.5)		

Note. *N* = 104. **N* = 103

Analysis for Aim 3

Table 5 presents Pearson's correlation of the relationship between QOL and the continuous explanatory study variables (sociodemographic, clinical, and psychological characteristics).

Sociodemographic Characteristics

The two-tailed correlation analysis showed that older participants had better QOL than younger participants, $r(102) = .26, p < .01$. The QOL varied significantly by marital status, $t(73.07) = -2.24, p < .05$, with the participants identifying as currently married ($M = 4.45, SD = .77$) having better QOL than those unmarried (previously/never married category; $M = 4.01, SD = 1.14$). QOL did not significantly correlate with race, $t(102) = .01, p = .99$; gender, $t(84.85) = -.12, p = .90$; education, $F(2, 101) = .13, p = .88$; or employment, $F(3, 100) = 2.50, p = .06$.

Clinical Characteristics

Worse physical functioning, $r(102) = -.44, p < .01$, and lower disease severity measured by the NYHA Class, $r(102) = -.32, p < .01$, were significantly associated with QOL. However, QOL did not significantly correlate with EF, $r(102) = .04, p = .71$, or comorbidity, $r(102) = -.13, p = .19$.

Psychological Characteristics

Better QOL significantly correlated with lower anxiety, $r(102) = -.59, p < .01$, and lower depression, $r(102) = -.73, p < .01$, scores.

Table 5

Pearson's r Correlation Analysis Examining the Relationship Between Sociodemographic Characteristics and Psychological Characteristics

	1	2	3	4	5	6	7	8
1. Race	–	-.353**	.278**	-.230*	-.180	-.040	.113	.025
2. Age		–	-.117	.347**	.896	.410**	-.308*	-.131
3. Gender			–	-.035	-.132	-.045	.123	.057
4. Marital status				–	.179	.150	-.127	-.179
5. Education					–	-.186	-.262**	-.211*
6. Employment status						–	-.044	
7. Anxiety (HADS-A)							–	.651**
8. Depression (HADS-D)								–

* $p < .05$, ** $p < .01$.

Note. $N = 104$.

Table 6

Pearson's r Correlation Analysis Examining the Relationship Between Sociodemographic Characteristics and Psychological Characteristics

	1	2	3	4	5	6	7	8
1. QOL	–	.268**	-.425**	.036	-.299**	-.128	-.586**	-.696**
2. Age		–	-.069	.202*	-.092	.208*	-.308**	-.131
3. Physical Functioning			–	.015	.297**	.342**	.421**	.582**
4. Ejection Function				–	-.264**	.166	-.068	.008
5. NYHA Class					–	.161	.262**	.251*
6. Comorbidity						–	.102	.174
7. Anxiety (HADS-A)							–	.651**
8. Depression (HADS-D)								–

* $p < .05$, ** $p < .01$.

Note. $N = 104$.

Table 7 presents the independent-samples t test and one-way ANOVA of the relationship between QOL total scores with the categorical explanatory study variables (sociodemographic, clinical, and psychological characteristics).

Table 7

Independent Samples t-Test and One-Way ANOVA Analysis Examining the Differences Between QOL and Categorical Sociodemographic and Clinical Characteristics

Variable	<i>n</i>	<i>M (SD)</i>	<i>t (df)</i>	<i>p</i>
Gender			-.12 (84.85)	.90
Male	69	4.25 (1.04)		
Female	35	4.28 (.81)		
Race			.01 (102)	.99
White	53	4.26 (.87)		
Black	51	4.26 (1.06)		
Education level			.13 (2, 101)	.88
Less than high school	16	4.15 (.87)		
High school	53	4.29 (1.00)		
Completed college	35	4.27 (.97)		
Marital status			-2.24 (73.07)	.03*
Previously/never married	45	4.01 (1.14)		
Currently married	59	4.45 (.77)		
Employment status			2.50 (3, 100)	.06
Employed full-time	17	4.60 (.80)		
Not employed full-time	12	4.08 (.97)		
Receiving disability	43	4.01 (1.00)		
Retired	32	4.49 (.93)		

* $p < .05$.

Multivariate Analysis

Except for race, the variables that significantly correlated with QOL in the bivariate analysis were entered into the hierarchical multiple linear regression model for QOL. This study focused on racial differences; therefore, race was the design variable entered and retained in the regression model, although it did not significantly correlate with QOL in the bivariate analysis. Forward and backward elimination approaches occurred for significant QOL predictors (see Table 11). Race was not a significant QOL predictor when assessed independently of other variables ($B = -.01$, $SE = .19$, $\beta = -.01$, $p = .95$). The forward elimination approach showed the

statistical significance of the full model, which included race, age, marital status, physical functioning, NYHA Class, anxiety, and depression, in Step 3, $F(98) = 21.32, p < .001$, and that depression accounted for 62% of the variance in QOL ($R^2 = .62$, Adjusted $R^2 = .59$). Regarding individual predictors, the sociodemographic characteristic of age and the psychological characteristic of depression significantly correlated with QOL in the full model. Younger age ($B = -.01, SE = .01, \beta = -.17, p = .03$) and greater depression ($B = -.14, SE = .02, \beta = -.62, p < .001$) significantly associated with lower QOL. Physical functioning and anxiety were significant predictors of QOL in Step 2. However, the independent variables (race, marital status, physical functioning, NYHA Class, and anxiety) did not significantly correlate with QOL in Step 3.

Table 8*Hierarchical Multiple Linear Regression Examining Quality of Life (Forward Elimination)*

Variable	B (SE)	β	<i>p</i>
Step 1			
Race	-.01 (.19)	-.01	.95
Step 2			
Race	.14 (.17)	.07	.39
Age	.01 (.01)	.13	.13
Marital status	-.06 (.16)	-.03	.70
Physical functioning (SF-36)	-.01 (.00)	.19	.04*
NYHA Class	-.20 (.12)	-.14	.10
Anxiety (HADS-A)	-.71 (.14)	-.47	<.001**
Step 3			
Race	.13 (.01)	.06	.36
Age	.01 (.01)	.17	.03*
Marital status	.18 (.14)	.09	.21
Physical functioning (SF-36)	-.00 (.00)	-.03	.72
NYHA Class	-.17 (.10)	-.11	.10
Anxiety (HADS-A)	-.23 (.14)	-.15	.10
Depression (HADS-D)	-.14 (.02)	-.62	<.001**

Step 1: $F(103) = .004, p = .95, R^2 = -.00, \text{adjusted } R^2 = -.01.$

Step 2: $F(98) = 12.52, p <.001, R^2 = .45, \text{adjusted } R^2 = .41.$

Step 3: $F(98) = 21.32, p <.001, R^2 = .6, \text{adjusted } R^2 = .59.$

* $p <.05, **p <.01.$

Note. $N = 104.$

A backward elimination approach showed the model's strength. Older age and greater depression remained significant QOL predictors in both the full (older age, $B = -.01, SE = .01, \beta = -.17, p = .03,$ and greater depression, $B = -.14, SE = .02, \beta = -.62, p <.001$) and trimmed (older age, $B = .02, SE = .01, \beta = .21, p = .00,$ and greater depression, $B = -.16, SE = .02, \beta = -.70, p <.001$), models (see Table 12).

Table 9*Hierarchical Multiple Linear Regression (Backward Elimination) Examining Quality of Life*

Variable	B (SE)	β	<i>p</i>
Step 1			
Race	.13 (.14)	.06	.36
Age	.01 (.01)	.17	.03*
Marital status	-.13 (.13)	-.07	.33
Physical functioning (SF-36)	-.00 (.00)	-.03	.72
NYHA Class	-.17 (.10)	-.11	.10
Anxiety (HADS-A)	-.23 (.14)	-.15	.10
Depression (HADS-D)	-.14 (.02)	-.62	<.001**
Step 2			
Race	.13 (.14)	.06	.37
Age	.01 (.00)	.17	.02*
Marital status	-.13 (.13)	-.07	.30
NYHA Class	-.16 (.09)	-.11	.11
Anxiety (HADS-A)	-.22 (.14)	-.15	.10
Depression (HADS-D)	-.13 (.02)	-.61	<.001**
Step 3			
Race	.15 (.14)	.08	.27
Age	.01 (.01)	.17	.03*
NYHA Class	-.17 (.09)	-.12	.08
Anxiety (HADS-A)	-.21 (.14)	-.14	.12
Depression (HADS-D)	-.13 (.02)	-.58	<.001**
Step 4			
Race	.17 (.14)	.09	.23
Age	.02 (.01)	.21	.00*
Depression (HADS-D)	-.16 (.02)	-.70	<.001**

Step 1: $F(98) = 21.32, p < .001, R^2 = .62, \text{adjusted } R^2 = .59.$

Step 2: $F(98) = 12.52, p < .001, R^2 = .62, \text{adjusted } R^2 = .60.$

Step 3: $F(101) = 28.13, p < .001, R^2 = .59, \text{adjusted } R^2 = .57.$

Step 4: $F(101) = 43.15, p < .001, R^2 = .57, \text{adjusted } R^2 = .56.$

* $p < .05, **p < .01.$

Note. $N = 104.$

References

- Aguinis, H., Gottfredson, R. K., & Joo, H. (2013). Best-practice recommendations for defining, identifying, and handling outliers. *Organizational Research Methods, 16*(2), 270–301. <https://doi.org/10.1177/1094428112470848>
- Bujang, M. A., Omar, E. D., & Baharum, N. A. (2018). A review on sample size determination for Cronbach's alpha test: A simple guide for researchers. *The Malaysian Journal of Medical Sciences, 25*(6), 85–99. <https://doi.org/10.21315/mjms2018.25.6.9>
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using GPower 3.1: Tests for correlation and regression analyses. *Behavior Research Methods, 41*(4), 1149–1160. <https://doi.org/10.3758/BRM.41.4.1149>
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). GPower 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods, 39*(2), 175–191. <https://doi.org/10.3758/BF03193146>
- IBM Corp. (2020). *IBM SPSS Statistics for Windows, Version 27.0*. IBM Corp.

CHAPTER 5: DISCUSSION

Introduction

The purpose of this descriptive, comparative study was to examine the relationships between sociodemographic, clinical, and psychological characteristics and quality of life (QOL) in patients with heart failure (HF) to determine racial differences between Blacks and Whites. The research question was, What are the differences between Blacks and Whites with heart failure with respect to sociodemographic and clinical characteristics and quality of life? The study focused on the differences, relationships, and influences of sociodemographic, clinical, and psychological characteristics on QOL in Blacks and Whites with HF. The HF racial disparities reduction framework (see Figures 2 and 3) guided the study to address racial disparities in the HF population. A research framework for racial disparities in gynecological cancer was appropriate to address potential racial differences to contribute to the design and testing of innovative approaches and interventions for effectiveness and efficacy (Doll, 2018). With a foundation of the HF racial disparities reduction framework (see Figures 2 and 3), the study provided a better understanding of the relationships between sociodemographic, clinical, and psychological characteristics and their influence on QOL and the racial differences contributing to persistent disparities among Blacks with HF.

This chapter presents the discussion and contextualization of the study's results. First addressed are the results related to the study's goals and limitations. Second, the chapter presents the study's conclusions and practice implications. Third, there are recommendations for future research on Black persons with HF and racial disparities.

Discussion

This section presents the results based on the study's goals, the differences by race, and the relationships between the sociodemographic, clinical, and psychological characteristics.

There are discussions of the sociodemographic, clinical, and psychological characteristics that correlated with and impacted QOL.

Aim 1: Racial Differences and Associations Between Sociodemographic, Clinical, and Psychological Characteristics

The results showed differences by race for the following sociodemographic characteristics: age, gender, and marital status. Education level and employment status did not significantly differ by race. Among the clinical characteristics, only ejection fraction (EF) differed by race, with Blacks having lower (worse) EF than Whites. This finding could have resulted from several factors, including time since diagnosis. Disease severity, as measured with the New York Heart Association Classification (NYHA Class), comorbidities, and functional status did not differ based on race. The psychological characteristics of depression and anxiety were significantly correlated, a finding consistent with prevalent clinical presentations and studies (Celano et al., 2018; Dekker et al., 2014; Evangelista, Ter-Galstanyan, Moughrabi, Moser, et al., 2009). However, the results indicated no differences in depression or anxiety between the Black and White participants.

Sociodemographic Characteristics. Race. Neither depression nor anxiety significantly correlated with race in this sample. This finding aligned with prior researchers who found no significant racial differences in depression between Black and White individuals with HF (Freedland et al., 2003a; Gottlieb et al., 2004; Griffin et al., 2007; Piamjariyakul et al., 2013; Rohyans & Pressler, 2009; Xiong et al., 2015). However, other scholars have identified racial

differences and concluded that Black individuals with HF had less depression than Whites (Ahmed et al., 2006; Koenig, 1998; Stamp et al., 2014; Vaccarino et al., 2001; Wierenga et al., 2017) or more depression than Whites (Dekker et al., 2014; Evangelista, Ter-Galstanyan, Moughrabi, & Moser, 2009; Freedland, Carney, et al., 2021). Sampling or measurement approaches could have contributed to these differences across studies.

Age. The Black participants in this sample were younger than the White participants. This finding aligned with HF epidemiology in Blacks, as Blacks with HF tend to be younger, on average, than Whites (Benjamin et al., 2017; Bibbins-Domingo et al., 2009; Go et al., 2013; Tandon et al., 2020; Vaccarino et al., 2002; Virani et al., 2020, 2021). This finding also aligned with other studies that found differences in age by race among Black and White patients with HF (Afzal et al., 1999; Glynn et al., 2019; Koenig, 1998; Lewis et al., 2011; Piamjariyakul et al., 2018; Stamp et al., 2014; Thomas et al., 2011; Wierenga et al., 2017; Ziaecian et al., 2017). Thus, younger Blacks remain disproportionately affected by HF (Tsao et al., 2022; Virani et al., 2020, 2021).

In this study, age did not correlate with depression. This finding differed from prior studies showing younger patients with HF had more depression (Athilingam et al., 2017; Chung et al., 2009; Freedland et al., 2003a; Gottlieb et al., 2004; Koenig, 1998; Mentz et al., 2015; Rohyans & Pressler, 2009; Stamp et al., 2014; Xiong et al., 2015). Conversely, Dickson et al. (2013) found that older patients with HF had more depression. Unlike other research, this study did not include participants in NYHA Class I, who often have less severe disease symptoms and more recent HF diagnoses.

Blacks are more likely to be diagnosed with HF at a younger age (Tsao et al., 2022; Virani et al., 2020, 2021), and younger age correlates with more depression with HF (Athilingam

et al., 2017; Chung et al., 2009; Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Koenig, 1998; Mentz et al., 2015; Rohyans & Pressler, 2009; Stamp et al., 2014; Xiong et al., 2015) and anxiety (Dekker et al., 2014). If this study's sample had included younger, more recently diagnosed participants in NYHA Class I, the results might have aligned with prior research showing increased depression and anxiety.

Age correlated with anxiety in this study, as the younger participants reported higher anxiety. This finding contributed to the limited research on anxiety in patients with HF. Little research has focused on anxiety in patients with HF, but Dekker et al. (2014) also found that anxiety correlated with younger age. A better understanding of the relationship between anxiety and HF could provide insight into younger patients with HF, of which Blacks comprise the majority (Tsao et al., 2022; Virani et al., 2020, 2021).

Gender. The sample included almost equal numbers of Black men and women ($n = 27$ and $n = 24$, respectively). However, White women were an underrepresented population in this sample ($n = 11$, with $n = 42$ White men), a finding that aligned with prior studies showing an underrepresentation of women in HF research (Bozkurt & Khalaf, 2017; De Bellis et al., 2019; DeFilippis et al., 2022; Habal et al., 2019; Heiat et al., 2002; Hsich & Pina, 2009; Reza et al., 2022). Although this study had a small White female sample, there was an adequate Black female sample. Gender and its intersection with race remain important, as prior studies have found gender differences (Dekker et al., 2014; Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Vaccarino et al., 2001; Xiong et al., 2015) in depression (Ahmed et al., 2006), anxiety (Dekker et al., 2014); QOL (Jackson & Emery, 2011), and race. For example, Black men are at the greatest risk of hospital readmissions (Chen et al., 2013), and Black women report more depression than Whites (Gottlieb et al., 2004).

Marital Status. The Black participants in this sample were less likely to be married. This was consistent with Wierenga et al.'s (2017) findings that Blacks were less likely to be married. Although this study did not include living arrangements, the finding suggests that Black individuals might have less access to social and financial support due to the absence of live-in marital partners. Patients with HF who are not married might need additional support to avoid poor clinical outcomes (Chung et al., 2009).

Education. In this study, more Blacks completed high school, and more Whites completed college. This finding aligned with prior research indicating that Black individuals with HF are less educated than their White counterparts (Lewis et al., 2011; Piamjariyakul et al., 2018). Also, education level correlates with health literacy in patients with HF (Cajita et al., 2016; Peterson et al., 2011). Blacks generally have less health literacy than Whites (Chaudhry et al., 2011; Morrow et al., 2006; Murray et al., 2009; Westlake et al., 2013). Less-educated Black patients with HF might have lower health literacy and struggle to understand HF's complex pathophysiology, severity, and self-management requirements (Cajita et al., 2016; Deek et al., 2020; Dennison et al., 2011; Rockwell & Riegel, 2001; Zou et al., 2017). In Allen et al.'s (2008) study, Black patients with HF in one sample without a college education overestimated their life expectancy, which suggests an incomplete understanding of their prognosis despite verbalizing their understanding of healthcare professionals' education.

Education also has implications for SES and health outcomes. SES correlates with depression in patients with HF (Hertzog et al., 2010; Koenig, 1998; Retrum et al., 2012; Vaccarino et al., 2001). Individuals with less education often earn less income (Eng & Feeny, 2007). Although there is no consistent relationship between higher education and increased income in Black individuals compared to Whites (Assari et al., 2018), there could be a

relationship between financial distress and depression in Blacks (Assari, 2019). Lower SES also negatively impacts access to health care, including among patients with chronic illnesses (Canedo et al., 2017; Georgopoulou et al., 2011; Sewell & Velayos, 2013). Thus, SES can impact QOL (Bielderma et al., 2015; Keyvanara et al., 2015; Tuzun et al., 2015), including for patients with HF (Verma et al., 2016). Patients with HF and less education could have fewer financial resources to adequately self-manage their health by adhering to dietary restrictions and medication regimens (Dickens et al., 2019; Macabasco-O'Connell et al., 2008).

Employment Status. This study found no racial differences in employment status. This finding was unsurprising, as many patients with HF lack employment due to HF's physiological and psychosocial impacts (Bennett & Sauv e, 2003; Ohlsson et al., 2021; Phillips et al., 2005; Pressler, 2008; R rth, Fosb l, et al., 2018; R rth, Wong, et al., 2016). This study's findings aligned with research indicating that Blacks with HF are likelier to be younger (Benjamin et al., 2017; Go et al., 2013; Lewis et al., 2011; Piamjariyakul et al., 2018; Roger et al., 2012; Stamp et al., 2014; Tsao et al., 2022; Virani et al., 2020, 2021; Wierenga et al., 2017). Most patients with HF are ineligible for retirement benefits, instead relying on disability benefits to meet their financial needs (Seo et al., 2021). However, disability benefits do not often cover living costs and might not enable living above the poverty level, resulting in increased financial strain and depression (Chung et al., 2009; Cirelli et al. (2018).

Clinical Characteristics. Ejection Fraction. Both measures of disease severity (NYHA Class and EF) correlated in this study's sample. The Black participants had lower EF than White participants, a finding that contributed to the literature because prior scholars had not stratified findings related to EF by race. The Black participants with HF were significantly younger than Whites in this sample, which aligned with the general HF patient population (Tandon et al.,

2020; Virani et al., 2020, 2021). However, the Black participants had lower EF, potentially indicating greater disease severity at a younger age. Greater disease severity could correlate with a greater risk of depression and anxiety and worse QOL (AbuRuz, 2018; Aggelopoulou et al., 2017; Carson et al., 2009; de Leon et al., 2009; Freedland, Rich, et al., 2021; Lee et al., 2005). Therefore, addressing this pattern of worse heart functioning in Blacks is critical to improving racial disparities.

NYHA Class, Comorbidities, and Functional Status. This study showed no statistically significant differences in mean NYHA Class, co-morbidities, and functional status by race. Testing associations between these clinical variables for racial differences contributed to the research on racial differences between Black and White persons with HF. The results indicate the need for further research on the associations between these variables and the factors contributing to negative outcomes among Blacks with HF.

Psychological Characteristics. Depression, Anxiety, and Comorbid Depression and Anxiety. Finding no differences by race in depression aligned with prior studies on racial differences in depression (Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Griffin et al., 2007; Piamjariyakul et al., 2013; Rohyans & Pressler, 2009; Xiong et al., 2015). However, other research has found differences in depression (Ahmed et al., 2006; Bean et al., 2009; Evangelista, Ter-Galstanyan, Moughrabi, Moser, et al., 2009; Gottlieb et al., 2004) and anxiety by race. A potential contributor to this study's findings was using the Hospital Anxiety and Depression Scale (HADS) to measure depression and anxiety. The HADS could have had measurement issues resulting in false positives and negatives (Y. Wu, Levis, Sun, He, et al., 2021; Y. Wu, Levis, Sun, Krishnan, et al., 2020). However, other scholars used the Minnesota Living with HF Questionnaire (Gottlieb et al., 2004), the Center for Epidemiologic Studies (Piamjariyakul,

Thompson, et al., 2018) Patient Health Questionnaire (Rohyans & Pressler, 2009), and the Diagnostic Interview Schedule (Freedland, Rich, Skala, et al., 2003) and found no significant differences in depression among Black and White participants.

The Black race did not correlate with anxiety in this sample. This finding contributed to the limited research on patients with HF and their anxiety and racial differences, which had little anxiety-related findings by race. Scholars have rarely studied anxiety among the overall HF patient population (Celano et al., 2018; Easton et al., 2016; Sokoreli et al., 2016), particularly Black individuals with HF. Although not reported by race, some research has suggested that patients with HF and anxiety have greater healthcare utilization and mortality (De Jong et al., 2011). Undiagnosed and untreated anxiety could contribute to disparate healthcare utilization and mortality rates among Blacks. Therefore, there is a need for further anxiety-focused research on potential racial differences.

Depression and anxiety were highly correlated in this sample, consistent with prior studies (Celano et al., 2018; Dekker et al., 2014; Evangelista, Ter-Galstanyan, et al., 2009; Freedland, Carney, et al., 2021). Evangelista, Ter-Galstanyan, et al. (2009) found that Blacks were likelier to have comorbid depression and anxiety than Whites. Comorbid depression and anxiety commonly occur in patients with HF, and their clinical presentation overlaps. Isolating each disorder's symptoms is challenging (Celano et al., 2018).

Relationships Between Psychological and Sociodemographic Characteristics.

Gender. Gender did not significantly correlate with depression or anxiety in this study's sample. The lack of a relationship between gender and depression aligned with studies that showed no significant differences in depression based on gender (Celik et al., 2016; Griffin et al., 2007; Rohyans & Pressler, 2009; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009). However, other

scholars found that women with HF reported significantly more depression (Ahmed et al., 2006; Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Jiang et al., 2001; Vaccarino et al., 2001; Xiong et al., 2015) and anxiety (Dekker et al., 2014) than men. However, this study's sample had a significant difference in participant gender, with men better represented, particularly among Whites. This sampling issue could have impacted the findings.

Marital Status. Depression and anxiety did not correlate in this sample, consistent with four studies that found no significant differences in depression among patients married or living alone and those living with family or friends (Dekker et al., 2014; Gottlieb et al., 2004; Rohyans & Pressler, 2009; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009). However, relationships between depression, anxiety, and marital status require further exploration because Chung et al. (2009) found that patients and their spouses felt depressed and anxious. There could be social support considerations and mental health implications for spouses of patients with HF.

Education Level. Lower education levels correlated with depression and anxiety in this sample. Although studies have had differing findings, a lower educational level is associated with greater depression (Celik et al., 2016; Chung et al., 2009; Vaccarino et al., 2001). Two other studies found correlations between higher levels of depression and anxiety (Aggelopoulou et al., 2017; Dekker et al., 2014). However, Brouwers et al. (2014) found that higher educational level correlated with greater depression in patients with HF. Thus, the research has had inconsistent findings. These three studies indicated above occurred in the United States, Turkey, and Greece. Cultural and societal norms and sampling, including of Blacks and other minorities in the United States compared to racially homogenous samples in other countries, could have impacted these studies' findings.

Employment Status. Employment status did not relate to depression or anxiety in this sample. Little research has focused on employment status and emotional well-being among patients with HF. However, Freedland, Rich, Skala, et al. (2003) found that unemployment due to disability significantly correlated with depression. Similarly, Ohlsson et al. (2021) identified a relationship between unemployment and increased mortality among patients with HF and adjusting for age and gender did not affect this finding.

Relationships Between Psychological and Sociodemographic Characteristics. The EF of this study's sample did not significantly differ by race. Interestingly, the results showed no significant differences by race in disease severity, as measured with EF and NYHA Class when comparing Blacks and Whites. This finding suggests that Black and White individuals with similar clinical status might have different outcomes, with Blacks tending to have negative or worse outcomes (Tsao et al., 2022; Virani et al., 2020, 2021). Hence, there is a need for further research on the factors contributing to different clinical outcomes.

Disease Severity: EF and NYHA Class. NYHA Class and EF were significantly related in this study's sample. This study's findings differed from Freedland, Rich, Skala, et al. (2003), who found depression more prevalent among individuals ($n = 682$) with lower EF within a sample that comprised 278 Blacks (41%). However, Freedland, Rich, Skala, et al. did not stratify the findings by race. This study did not find a significant association between depression and anxiety and EF. However, a higher NYHA Class was significantly associated with depression and anxiety. Thus, depression and anxiety significantly correlated with NYHA Class. This finding aligned with research that found associations with greater depression (Bean et al., 2009; Chung et al., 2009; De Jong et al., 2011; de Leon et al., 2009; Dekker et al., 2014; Dickson, McCarthy, & Katz, 2013; Freedland, Rich, Skala, et al., 2003; Gottlieb et al., 2004; Rohyans &

Pressler, 2009; Sharma et al., 2009; Trivedi et al., 2009) and NYHA Class. However, although EF and NYHA Class correlated, neither were significantly associated with depression and anxiety in this study.

Comorbidities. Neither depression nor anxiety was significantly associated with comorbidities in this study. This finding differed from prior studies that found comorbidities associated with depression (Patel et al., 2018) and anxiety (Dekker et al., 2014) in patients with HF. Lesman-Leegte et al. (2009) found an increased unadjusted odds ratio of 2.48 for depression in patients with HF without additional chronic conditions and 3.79 in those with additional chronic conditions. The risk for depression in patients with HF and participants with more chronic conditions was more than three times higher than in patients without chronic conditions (Lesman-Leegte et al., 2009). This finding aligned with Ahmed et al. (2006) and Sharma et al. (2009), who found no relationship between greater depression and comorbidities.

Functional Status. This study's results showed a correlation between lower functional status and depression and anxiety, aligning with prior studies on depression and, to a lesser extent, anxiety (Clarke et al., 2000; Dickson et al., 2013; Herr, Sayler, Flattery, et al., 2015; Shen et al., 2010; Vaccarino et al., 2001). Functional status limitations commonly occur among patients with HF, as HF symptoms frequently affect patients' ability to participate in activities of daily living (ADLs). Additionally, many patients with HF have fatigue and pronounced mobility challenges (Blinderman et al., 2008; Falk et al., 2008; Herr, Sayler, Lyon, et al., 2014; Herr, Sayler, Flattery, et al., 2015; Moser, Lee, et al., 2014). Patients unable to perform ADLs are likelier to be depressed and have worse functional status (de Leon et al., 2009; Freedland, Rich, Skala, et al., 2003). Improving functional status could be a way to reduce depression (X. Zhang et al., 2020).

Aim 2: Sociodemographic, Clinical, and Psychological Predictors of Quality of Life

This study showed that the sociodemographic characteristics of race, gender, education level, and employment status did not impact QOL. Of the clinical characteristics, EF and comorbidities did not impact QOL. Age, marital status, functional status, NYHA Class, and anxiety associated with QOL in the bivariate analysis were significant predictors of QOL in Steps 1 and 2 of the regression model building. However, depression was the most significant contributor to QOL.

Sociodemographic Characteristics and QOL. Race. An interesting finding was that race was not significantly associated with QOL in this sample, perhaps because the study did not include patients classified as NYHA Class I. Higher NYHA Class correlates with lower QOL (Coelho et al., 2005; Freedland, Rich, et al., 2021; Lee et al., 2005). However, most research has not focused on patients in NYHA Class I, likely because they are often asymptomatic. This omission could have resulted in limited interventions for this subpopulation, which might cause significant emotional distress. Relationships between race and QOL require further exploration, as does the influence of other sociodemographic, clinical, and psychological characteristics on patients with HF.

Age. In the bivariate analysis, age was significantly associated with QOL, as the older patients had better QOL. In the multivariate analysis, older age was a significant predictor of better QOL. Some studies found that younger patients have worse QOL. This study's results do not align with other research indicating worse QOL among older patients with HF (Luttik et al., 2006).

Gender. The bivariate analysis showed that gender was not significantly associated with QOL in this study's sample. This result aligned with other studies that found that QOL did not

vary by gender. However, some studies found that women with HF had worse QOL than men, even after adjusting for age and comorbidities (Lesman-Leegte et al., 2009; Riedinger et al., 2001).

Marital Status. Marital status was significantly associated with QOL in this study's sample, which aligns with prior research indicating that married individuals have higher QOL (Luttik et al., 2006). Similarly, Chung, Lennie, et al. (2009) found that patients with spouses had longer event-free survival. Marriage could be a protective factor and means of social support for patients with HF (Chung, Lennie, et al., 2009; Rohrbaugh et al., 2006). Many patients with HF are homebound and experience social isolation. Therefore, a marital partner who assists with ADLs, shares in financial responsibilities, assists with HF self-care, and provides emotional support could positively influence QOL. Chung, Moser, et al. (2013) also found that depression was a predictor of social support, and that social support was a QOL predictor QOL. Therefore, there are relationships between social support, lower depression, and better QOL.

Education and Employment Status. Education level and employment status did not significantly associate with QOL in this sample. In contrast, prior studies found that participants with less education had lower QOL (Barbareschi et al., 2011; Lee et al., 2005; Liu et al., 2021) and that patients with HF with employment had better QOL (AbuRuz, 2018). However, these studies occurred in other countries, and sociocultural differences could have resulted in differing findings. Unemployment correlates with greater mortality in patients with HF (Rørth, Fosbøl, et al., 2018). There is a need for further research on the relationships between employment status and QOL.

Clinical Characteristics and QOL. Disease Severity. Another interesting finding was that although the Black participants had significantly worse disease severity than Whites as

measured by the EF, they did not have significantly lower QOL than the White patients. This finding differed from the general HF patient population, as prior studies found an association between higher disease severity and lower QOL (AbuRuz, 2018; Carson et al., 2009; Freedland, Rich, et al., 2021). Additionally, like prior research, this study showed that a worse NYHA Class was associated with worse QOL (Coelho et al., 2005; Freedland, Rich, et al., 2021; Lee et al., 2005).

Comorbidities. In this study, comorbidities did not significantly associate with QOL. However, prior studies found that comorbidities negatively impacted QOL in patients with HF (Carson et al., 2009; de Leon et al., 2009; Lesman-Leegte et al., 2009; Reddy et al., 2020). Because Black individuals are likelier to have multiple chronic illnesses (Cuyjet & Akinboboye, 2014), they have the greatest risk for lower QOL. There is a need for further research on the relationships between comorbidities and QOL in patients with HF, as comorbidities with overlapping symptoms could impact a study's findings (de Groot et al., 2003). Managing multiple chronic illnesses is taxing and can negatively affect QOL. Future scholars could investigate comorbidity measures, as the long-standing Charlson Comorbidity Index (Charlson et al., 2022) might not be the best instrument for measuring phenomena with implications for clinical outcomes in large patient populations and reducing disparities among Blacks with HF.

Functional Status. Functional status is an element of QOL (Ferrans et al., 2005) measured in the QLI-Health and Functioning Subscale (Ferrans, 1990; Ferrans & Powers, 1985, 1992; Ghasemi et al., 2014; Warnecke et al., 1996). Based on prior research, one of this study's assumptions was that there would be a relationship with QOL (Wu et al., 2016). However, functional status does not consistently correlate with QOL in patients with HF (Carels, 2004; de

Leon et al., 2009). Different findings could have resulted from functional status measurements across studies.

Psychological Characteristics and QOL. *Depression.* Depression was the most significant predictor of QOL in this sample, accounting for 60% of QOL variance. The multivariate analysis results aligned with prior studies that found that depression negatively impacted QOL in patients with HF (AbuRuz, 2018; Aggelopoulou et al., 2017; Bekelman, Havranek, et al., 2007; Chung, Lennie, et al., 2009; Chung, Moser, et al., 2013; Cully et al., 2010; Gottlieb et al., 2004; Mentz et al., 2015; Salyer et al., 2019; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009). Despite limited evidence that QOL in Black individuals differs from the general population of patients with HF (de Leon et al., 2009; Warnecke et al., 1996), this study did not find racial differences in the sample. Instead of using a valid and reliable instrument, de Leon et al. (2009) measured QOL based on patient perceptions but excluded Black patients ($n = 209$) due to missing data. Black patients were also likelier to have lower QOL scores. Therefore, sampling and measurement issues could have contributed to the observed racial differences in this sample. Other studies had differing findings. For example, despite finding that participants with more depression and anxiety reported lower QOL, Bean et al. (2009) identified no statistically significant racial differences ($p > .05$).

Racial disparities, racial QOL differences, and the influence of depression on QOL require further exploration. Baseline and worsening depression correlate with poor QOL and all-cause mortality (Mentz et al., 2015), and Black patients have persistently worse clinical outcomes. Depression can negatively impact a patient's ability or willingness to engage in physical activity, regardless of ability (McCarthy et al., 2015). Thus, there is a need to discern whether patients have clinically significant depression (Bean et al., 2009; Blumenthal et al.,

2012), as it can negatively impact patients with HF, especially those who are Black (Mentz et al., 2015).

This study did not find significant differences in depression, as Black and White participants had similarly high depression rates. However, behavior caused by depression could be more detrimental to Blacks with HF and contribute to disparate clinical outcomes. Black individuals with HF with depression might be less likely than individuals of other races to adhere to diet, exercise, and medication regimens and report symptoms. Rather than seeking care for symptoms such as shortness of breath, fatigue, and fluid retention, Black patients with HF and depression might choose passive self-care, such as resting in bed or waiting for symptoms to subside. Some patients have attributed a lack of self-care to depression, sadness, or a lack of energy (Bean et al., 2009; de Leon et al., 2009; Dickson et al., 2013; Gottlieb et al., 2004; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009). Coping with depression can cause behavioral or mental disengagement, sometimes resulting in poor self-management practices, such as ignoring symptoms and medical recommendations (Trivedi et al., 2009). Individuals with HF and depression might recognize their health status is changing and their HF-related symptoms are worsening but delay contacting health care providers and seeking care (Evangelista et al., 2002).

Anxiety. Anxiety was a significant QOL predictor in Step 2 of the regression model—building process, but its significance decreased in the full model (Step 3). Although insignificant in the final model, this finding aligned with prior research that anxiety negatively impacts QOL in patients with HF (AbuRuz, 2018; Aggelopoulou et al., 2017; Chung, Moser, et al., 2009; Polikandrioti et al., 2019; Salyer et al., 2019).

Multivariate Analysis. Several variables were significant QOL predictors when entered into the stepwise regression model. In the final model, depression was the only significant

predictor of negative QOL and accounted for 60% of QOL variance. This study showed the negative impact of depression on the QOL of patients with HF. This finding aligned with prior studies indicating that depression negatively impacts QOL in patients with HF (Gottlieb et al., 2004).

Blacks are likelier to have severe, disabling, and untreated major depressive disorder (Williams et al., 2007). Additionally, Blacks with chronic illnesses have the highest depression rates, poorest health-related QOL, and worst functional status (Jackson-Triche et al., 2000). Black individuals are not screened, diagnosed, and treated at the same rates as Whites (Barnes & Bates, 2017), and depression and anxiety fluctuate over time (Hamatani et al., 2021; Hill-Joseph, 2019). Research might have had varied findings based on measurement timing. Contradictory evidence might have also resulted from cultural differences, cultural norms, and stigma in Blacks' perceptions of depression and anxiety and reluctance to identify with perceived negative mental health diagnoses (Das, Olfson, et al., 2006; Das, Roy, et al., 2019).

Among patients with HF, psychological distress can result in poor adherence to prescribed regimens (Eisele et al., 2020). In the presence of depression, patients with HF might struggle to engage in the self-care necessary to avoid hospital readmission and other adverse clinical outcomes (Lockhart et al., 2014). Some patients with HF prescribed antidepressant therapy remain depressed (Blumenthal et al., 2012; Chung, Moser, et al., 2013; Dekker et al., 2014). Depression was the most significant QOL predictor in patients with HF in this sample. Thus, depression in patients with HF requires attention and intervention by clinicians and researchers to prevent negative clinical outcomes.

Black patients have the highest rate of depression, the poorest health-related QOL, and the worst functional status (Jackson-Triche et al., 2000). Williams et al. (2007) found that Blacks

were likelier to have severe, disabling, and untreated major depressive disorder. Providers can address depression's significant influence on the QOL of patients with HF and disparities among Blacks (Tsao et al., 2022; Virani et al., 2020, 2021) by proactively screening Black patients with HF for depression. However, due to stigma (Das, Olfson, et al., 2006) and other cultural norms, some Blacks might not admit their feelings to healthcare providers (Dickson et al., 2013; Gottlieb et al., 2004; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009) or identify with or admit to feeling depressed. Instead, Black patients might use other terms for depression, such as sadness, overwhelm, low energy, or feeling-down energy (Dickson et al., 2013).

Limitations

This study has several methodological limitations. First, the secondary data analysis focused on extant data via study methods designed to address the study's goals. Secondary data analysis is a way to improve feasibility. However, remaining limited to the methods used in a previous study with different goals did not enable the full exploration of all potential variables.

Second, the study included data collected in 2014. Older data could have differed from more recent data, which would have impacted this study's findings. Third, the relatively small sample from one setting may not have represented the general population of Black and White patients with HF despite methodological rigor. The sample aligned with the criterion of 50 participants per group to find differences (Cohen, 1992; Serdar et al., 2021), and there was adequate power based on the G*Power analysis (Faul, Erdfelder, Buchner, & Lang, 2009; Faul, Erdfelder, Lang, & Buchner 2007). The data had normal distribution; the instruments showed adequate reliability (Cronbach's alphas ranging from .83 to .96), and addressing the missing data with well-established standards. However, despite using proven strategies for handling missing

data, the missing data could have impacted the findings, as 10 cases were excluded from the already small sample (see Figure 4).

Fourth, the parent study included participants in NYHA Classes II–IV and omitted those in Class I. Therefore, the study lacked findings regarding the sociodemographic, clinical, and psychological characteristics and QOL of patients in NYHA Class I. Also, the parent study occurred with only the satisfaction aspect of the Quality of Life Index (QLI). The inability to assess the importance of various QOL aspects could have resulted in limited findings, particularly regarding race. Fifth, this study did not include data on whether patients took prescribed antidepressants and anxiolytics and how their medications affected their observed depression and anxiety scores. Patients with HF prescribed antidepressants and anxiolytics often remain depressed (Chung et al., 2013; Freedl et al., 2016; Jiang et al., 2001) and anxious.

Conclusion

Like prior research (AbuRuz, 2018; Aggelopoulou et al., 2017; Bekelman, Havranek, et al., 2007; Chung, Lennie, et al., 2009; Chung, Moser, et al., 2013; Cully et al., 2010; de Leon et al., 2009; Gottlieb et al., 2004; Mentz et al., 2015; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009), this study found that greater depression and younger age were the most significant predictors of worse QOL among patients with HF. Despite the small sample and limitations, the results showed the prevalence of and need to screen for and treat depression in patients with HF because of its negative effects on QOL and clinical outcomes (Freedl et al., 2016; Sokoreli et al., 2016). Depression remains understudied in patients with HF (Ahmed et al., 2006; Bhatt et al., 2016; Holzapfel et al., 2008).

This study's findings also indicate the importance of considering how various sociodemographic, clinical, and psychosocial characteristics impact patients with HF. Although

the results showed no racial differences in depression, anxiety, or QOL, racial disparities persist among Black patients with HF. For example, Black individuals have the highest healthcare utilization rates (Pahuja et al., 2022; Tsao et al., 2022). Therefore, there is a need for more studies with racial comparisons to gain insight into the drivers of disparate clinical outcomes.

HF's global economic burden of over \$108 billion (Cook et al., 2013) and projected impact (Heidenreich, Fonarow, et al., 2022; Heidenreich, Trogdon, et al., 2011; Lesyuk et al., 2018; Moradi et al., 2021; Tsao et al., 2022; Urbich et al., 2020) indicate the need to research and design interventions for all patients with HF, especially those disparately affected, such as Blacks (Tsao et al., 2022; Virani et al., 2020, 2021). Future research could focus on physiological and psychosocial contributors to depression and QOL in patients with HF and the factors contributing to racial disparities and disparate clinical outcomes among Blacks with HF.

Recommendations and Future Research

Racial disparities persist among Black individuals with HF (Tsao et al., 2022; Virani et al., 2020, 2021). Success in Phase III of the HF Disparities Reduction Framework (see Figures 2 and 3) and improving QOL require increasing knowledge and developing innovative approaches to improve depression and anxiety diagnosis and treatment in patients with HF (Doll, 2018). Scholars and practitioners could design, develop, and deploy novel treatment strategies, culturally competent care, and culturally responsive interventions to address racial differences and disparate clinical outcomes among Black patients. Scholars could use several approaches and foci to advance the research.

First, future research could occur with different research designs and methodologies. Scholars could conduct quantitative studies with larger samples from multiple study sites, qualitative studies of patients' perceptions of depression and QOL, and mixed methods studies to

triangulate rich data about multiple facets of depression, anxiety, or QOL in HF patients. Considering patients' diverse experiences could provide a better understanding of their thoughts, feelings, perceptions, behaviors, beliefs, and values when developing culturally appropriate interventions to positively impact care (Riegel et al., 2016) and reduce healthcare disparities (Davidson et al., 2007; Watson, 1995).

HF is cyclical, and depression and anxiety intensity vary over time (Hamatani et al., 2021; Hill-Joseph, 2019; Mentz et al., 2015; Regan et al., 2019). Black individuals experience a more significant fluctuation in depressive symptoms (Hill-Joseph, 2019). Scholars could conduct longitudinal studies to assess depression and anxiety patterns and their influence on QOL at various points with repeated measures (Hill-Joseph, 2019; Konstam et al., 2005) and assessments for racial differences. Researchers could observe variations as patients move across the healthcare continuum and through various healthcare delivery settings, especially with stressors such as readmission to inpatient facilities or HF exacerbations requiring outpatient provider visits.

Researchers could conduct secondary analyses to identify means of interrupting racial disparities among patients with HF. To determine racial differences, investigators could conduct secondary data analyses using data sets that include sociodemographic, clinical, and psychological characteristics, QOL, and representative Black samples. Practitioners could use such findings in making depression and anxiety screening tools a standard of care in primary and acute care settings. Screening would help to identify patients who could benefit from proactive interventions to prevent depression and anxiety from negatively impacting QOL and contributing to other negative clinical outcomes.

Black individuals comprise the majority of the patients with HF (Benjamin, Blaha, et al., 2017; Benjamin, Munter, et al., 2019; Benjamin, Virani, et al., 2018; Go et al., 2013; Go et al., 2014; Mozaffarian et al., 2016; Mozaffarian et al., 2015; Roger et al., 2012; Tsao et al., 2022; Virani et al., 2020, 2021). Despite being disproportionately affected by HF (Tsao et al., 2022; Virani et al., 2020, 2021), Black persons are underrepresented in HF research (Evangelista, Ter-Galstanyan, Moughrabi, Moser, et al., 2009; Gaffey et al., 2022). Race remains largely unexamined as a variable of potential influence. Although Black participants are available, investigators might not recruit them. The scholars whose samples have included Black individuals with HF and had adequate power to detect racial differences have seldom stratified findings by race. Decreasing racial disparities among Blacks with HF requires further investigation into the associated drivers and contributors. Intentionally including Blacks and identifying racial differences could be a way to advance science in this area and inform the development of innovative interventions for racial disparities among Blacks with HF. Future scholars could include Black participants for more representative samples to compare phenomena among patients with HF by race and use race as a variable with predictive value.

Researchers conducting studies with representative Black samples could isolate race as a variable that could impact the outcomes under study. Rather than reporting race descriptively, scholars could assess race's influence on and predictive value for various clinical outcomes. Grouping minorities in analyses prevents the opportunity to detect racial differences, minimizing and disregarding race's potential influence on health outcomes. Race can impact health and health outcomes and is a social determinant of health (Bhopal et al., 2021; Culley, 2006). Researchers who consider sociodemographic factors and cultural differences in focused and

unbiased inquiries into patients' experiences could better understand racially, ethnically, and culturally influenced phenomena.

Black individuals' long-standing mistrust of the healthcare system and health-related research could be barriers to conducting health disparities research (Corbie-Smith, Thomas, St. George, 2002; Corbie-Smith, Thomas, Williams, & Moody-Ayers, 1999; Oh et al., 2015; Scharff et al., 2010; Simmonds, 2008). Therefore, setting specific recruitment goals (Durant et al., 2007), understanding barriers to (Branson et al., 2007; Hughes et al., 2017; Luebbert & Perez, 2016; Pugh et al., 2022; Rollins et al., 2018) and reasons for nonparticipation (Nooruddin et al., 2020), building trust, and gaining cultural awareness and competence (Rucker-Whitaker et al., 2006; Trantham et al., 2015) are essential when researching health disparities among Blacks (Byrd et al., 2011; Scharff et al., 2010). Innovative approaches to foster trust and connection, such as community engagement and involvement, could enable Black individuals to understand how participating in research might improve the human condition of other Blacks with HF and Black communities (Byrd et al., 2011; Lang et al., 2013; Spruill, 2010).

Scholars could also provide incentives, train staff in culturally congruent recruitment strategies, and diversify research teams to obtain more Black participants' agreement and commitment to participate in HF research (Billingsley, 2014; Branson et al., 2007; Browne et al., 2010; Ford et al., 2013; Fuqua et al., 2005; Graham et al., 2018; Hayes & Sharma, 2021; Huang & Coker, 2010; Hughes et al., 2017; Isler & Corbie-Smith, 2012; Lang et al., 2013; Luebbert & Perez, 2016; Mody et al., 2008). Further, investigators should not assume that Blacks will not participate in research, as Millon Underwood et al. (2013) found that 88.7% of their all-Black sample ($n = 212$) reported that they would have taken part in studies had anyone asked them.

Using this study's model (see Figure 3) to test the relationships between sociodemographic, clinical, and psychological characteristics, QOL, and racial differences could contribute to the HF racial disparities research. Adding new sociodemographic, clinical, and psychosocial variables to the model could provide a better understanding of the factors contributing to persistent racial disparities, which might address the psychosocial needs of younger patients with HF, who are disproportionately Black (Gottlieb et al., 2004; Tandon et al., 2020; Virani et al., 2020, 2021). There is a need for more research on the intersection of race and age and its influence on depression, anxiety, and QOL. There is also a need to focus on preventative efforts and targeted interventions for younger Blacks with different age-based needs.

Women with HF have physiological and psychosocial differences (De Bellis et al., 2019; Hopper et al., 2016; Lee & Riegel, 2018; Sethares & Chin, 2021). Therefore, scholars could compare race and gender intersections with representative samples of women from all races to generate knowledge and design tailored interventions. The relationships between marital status, depression, anxiety, and QOL require further exploration, with an emphasis on racial differences. Scholars could explore marital quality in the general HF patient population and stratify the data by race to determine whether there are racial differences.

Patients with higher education levels tend to have less depression and anxiety (Bjelland et al., 2008). Thus, there is a need to understand the relationships and racial differences among education, depression, and anxiety. Researchers should explore the relationships between employment status, depression, and QOL because patients with HF who are unemployed are likelier to have poor adherence to prescribed medication regimens (Dickson et al., 2008) and experience negative clinical outcomes related to loss of livelihood (Rørth, Fosbøl, et al., 2018).

Future studies could also address SES, an important focus of health disparities research (Shavers, 2007; Winkleby et al., 1992). Scholars could explore the relationships among education, employment status, income, depression, anxiety, and QOL in patients with HF. SES might be an important indicator of racial differences among Black and White individuals with HF. Although Black persons with HF might receive disability benefits, unemployment, underemployment, and limited financial resources could cause financial stress and strain and contribute to negative clinical outcomes (Clark et al., 2003; Dewan et al., 2019; Hung et al., 2021; Lindenauer et al., 2013; Tansey, 2010; Tromp et al., 2020).

Future studies with patients in NYHA Class I could have different findings regarding depression, anxiety, and QOL, as this study included only patients in NYHA Classes II–IV. Data across the HF severity spectrum could provide better insight into depression, anxiety, and QOL. Additionally, Black individuals receive HF diagnoses at younger ages than Whites (Tsao et al., 2022; Virani et al., 2020, 2021). Younger Black patients newly diagnosed with HF might fall into NYHA Class I. Although patients in NYHA Class I do not have severe symptoms, they are at risk for depression, anxiety, and poor QOL, especially during the adjustment phase after diagnosis. Poor self-care during NYHA Class I could contribute to disease progression and more rapid and severe illness phases. Focusing on this seemingly lower-risk subpopulation is necessary because its members might be despondent about their diagnosis and long-term prognosis.

Future scholars could investigate correlations between EF and NYHA Class and their influence when measuring disease severity. Scholars could also research the contributors to HF disease severity among Black individuals. Future studies could also indicate differences in phenomena based on disease severity, as measured by reduced versus preserved EF.

Relationships between disease severity, depression, anxiety, and QOL require further exploration focusing on the best approaches for measuring disease severity in patients with HF.

This study did not find a correlation between comorbidities and depression, anxiety, or QOL. However, other researchers found adverse outcomes associated with comorbidities, including lower QOL in patients with HF (Carson et al., 2009; Freedland, Rich, et al., 2021; Heo et al., 2012; Reddy et al., 2020). Future researchers could further assess these relationships and test for racial differences. Because Black individuals with HF might experience greater functional impairment (Vaccarino et al., 2002), there is a need for more research on the relationships between functional status and QOL to address racial differences. Future scholars could assess the medications prescribed to patients with HF, such as beta blockers, that might correlate with depression (Silver, 2010) and prescriptions for other chronic illnesses that might lead to adverse outcomes in patients with HF (Amabile & Spencer, 2004). Researchers could also assess medication adherence and its potential relationship with sociodemographic, clinical, and psychosocial characteristics, as medication adherence contributes to optimal outcomes in patients with HF. However, multiple factors often impact medication adherence (Andrews et al., 2017; Molloy et al., 2012).

Contemporary scholars quantifying disease severity, progression, and functional status in patients with HF tend to use the Kansas City Cardiomyopathy Questionnaire (KCCQ) rather than the NYHA Class (Greene et al., 2021) or EF. The KCCQ more comprehensively addresses patient-reported health outcomes, enables patient monitoring over time, and includes diverse patient characteristics (Spertus et al., 2020). Therefore, the KCCQ could be a better measure of the clinical and psychosocial characteristics of patients with HF, including disease severity.

Inflammation is a possible physiological contributor to depression in patients with HF due to an activated immune response (Brouwers et al., 2013; Ghosh et al., 2016; Petersen & Felker, 2006; Xiong et al., 2015). Other clinical variables, such as disease severity, functional status, poor appetite, and mortality, correlate with inflammatory biomarkers (von Haehling et al., 2009). Studying the relationships between inflammatory biomarkers (e.g., tumor necrosis factor; Interleukin 1, 6, and 18; and C-reactive protein and depression) could provide insight into depression in patients with HF, including those who are Black, who remain underrepresented in the literature. Further insight might provide information on improving QOL in patients with HF by addressing depression's physiological basis with interventions for preventing and treating depression related to inflammation.

Future researchers could also explore the relationships between inflammatory biomarkers and anxiety. Depression is a significant predictor of QOL, and depression and anxiety correlate (Celano et al., 2018; Dekker et al., 2014; Dickson et al., 2013), as was also found in this study. Examining the relationships between comorbid depression and anxiety and inflammatory biomarkers could address HF's physiological and psychosocial aspects and contribute to the biobehavioral research on the interplay of these chronic disorders.

Symptom science research has provided insight into the effects of symptoms and symptom clusters on patients with HF (Herr, Salyer, Flattery, et al., 2015; Herr, Salyer, Lyon, et al., 2013; Jurgens et al., 2009; Lee & Riegel, 2018; Moser et al., 2014; Park et al., 2019; Salyer et al., 2019; Sethares & Chin, 2021; Son & Won, 2018). However, most studies have lacked a cultural lens. There is a need for further study of symptoms and symptom clusters with race stratification because HF's often-intense symptoms can impact depression, anxiety, and QOL (Heo, Doering, et al., 2008; Heo, Lennie, et al., 2009; Heo, Moser, Lennie, et al., 2007). Using a

disparities lens to examine how symptoms contribute to and occur with depression, anxiety, and poor QOL could provide insight into how to improve QOL. Thus, scholars could stratify findings by race to capture any racial differences. Exploring differences in symptom clusters based on other sociodemographic and clinical characteristics would also contribute to the literature.

Future researchers could address clinical characteristics by assessing equity in the treatment of Black patients with HF, as inequities in the medical management of HF among Black individuals (Cascino et al., 2022; Mitchell et al., 2011) could contribute to adverse clinical outcomes, including hospital readmission (Wispelwey et al., 2019). Identifying inequities might indicate how to tailor interventions to foster equity in HF care and treatment for all races (Mensah et al., 2019; Ornelas et al., 2021).

Further study on psychological characteristics could show how to decrease depression and anxiety and improve QOL in patients with HF. Emotional distress can negatively impact Black patients with HF and contribute to adverse clinical outcomes, including decompensation, rehospitalization, and mortality (Ghali et al., 1988; Mentz et al., 2015). Black patients might not be screened, diagnosed, and treated for depression at the same rates as Whites (Barnes & Bates, 2017; Lewis et al., 2011; Mentz et al., 2015; Williams et al., 2007), and even when treated, Black patients might remain depressed (Chung et al., 2013; Freedl et al., 2016; Jiang et al., 2001). Access to care challenges and health equity also require further investigation because the mental health care provided to Black individuals could be inadequate compared to Whites (Cook et al., 2014).

This study found no racial differences in depression and anxiety. There is a need to understand depression and anxiety in Black individuals with HF because they might perceive depression and anxiety differently than other races, despite similarly high levels of psychological

distress. Due to cultural norms and stigma regarding negative emotions (Das, Olfson, et al., 2006; Das, Roy, et al., 2019), Black patients might underreport depression and anxiety to healthcare providers (Dickson, McCarthy, & Katz, 2013; Gottlieb et al., 2004; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009). They might also communicate less directly regarding depression, considering themselves overwhelmed with sadness, low spirits, or a lack of energy (Dickson et al., 2013). Thus, Black individuals might choose a passive response, such as resting in bed or waiting for symptoms to subside (Bean et al., 2009; de Leon et al., 2009; Dickson et al., 2013; Gottlieb et al., 2004; Sharma, Zehtabchi, Rojas, & Birkhahn, 2009). The passive approach has a cyclical effect because depression worsens among patients unable to effectively perform HF self-care (Dickson et al., 2013; Freedland, Carney, & Rich, 2011). Black individuals with chronic illnesses have the highest rates of depression, the poorest health-related QOL, and the worst functional status (Jackson-Triche et al., 2000) when compared to other races. Blacks are likelier to rate their depression as severe, very severe, disabling, and causing significant interference with functioning (Williams et al., 2007), potentially resulting in disparate HF clinical outcomes, including poor self-care (Alegría et al., 2008). Thus, Black individuals with HF require more proactive, intentional screening for depression and anxiety.

Providers cannot rely on Black patients to self-report depression or depressive symptoms, request antidepressants, or participate in therapy because Blacks are less likely to access mental health services than Whites (Alegría et al., 2008; Cook et al., 2014; Das, Olfson, et al., 2006; Ojeda & McGuire, 2006). Despite screening attempts, miscommunication regarding depression and anxiety could also contribute to missed diagnoses. Providers' attitudes and communication styles could impact Black patients' willingness to disclose depression. Some physicians minimize emotional expression in Black compared to White patients, resulting in less

antidepressant prescriptions (Sleath et al., 1998). Thus, there is a need for congruent provider communication regarding psychological distress, culturally appropriate screening, and a focus on understanding the patient's perspective (Humphreys et al., 2008).

Future researchers should examine methods for standardizing depression and anxiety measurement and screening of patients with HF to ensure measure reliability and validity in diverse samples. There are various measures for depression and anxiety in patients with HF. However, measurement errors could produce false high or low results because not all anxiety measures are equivalent (Clover et al., 2020). Thus, providers could misclassify depression depending on measurement approaches (Y. Wu, Levis, Sun, Krishnan, et al., 2020). Scholars have debated the best cutoff score for properly identifying depression (Tesio et al., 2014) and criticized the well-established HADS as a measure of depression (Y. Wu, Levis, Sun, He, et al., 2021; Y. Wu, Levis, Sun, Krishnan, et al., 2020) and anxiety (Clover et al., 2020).

Contending that the HADS might not be the best depression and anxiety measure, some investigators have suggested alternative scoring approaches to avoid false negatives and positives for major depression (Y. Wu, Levis, Sun, He, et al., 2021). Sensitivity and specificity issues in detecting major depression have led scholars to question the HADS and other instruments as accurate measures of depression. The measures might overlook symptoms that do not align with major depression diagnostic criteria but negatively impact patients with HF (Alhurani et al., 2015; Celano et al., 2018; Konstam et al., 2005; Nair et al., 2012; Rustad et al., 2013). Despite criticism, the HADS-D has adequate specificity for indicating depression in patients (Cassiani-Miranda et al., 2022).

Researchers could measure depression and anxiety to investigate the best measures for affective symptoms in patients with HF. Further research could show extant measures' cultural

congruence and appropriateness for identifying depression and anxiety in Blacks with HF. Scholars could focus on understanding the effectiveness of the best depression and anxiety measures in the general population of patients with HF and Black patients with HF. Researchers might focus on using culturally appropriate measures with internal consistency and reliability for all-Black samples. Future researchers could also isolate anxiety from depression to avoid confounding the two (Celano et al., 2018).

Depression is the most significant predictor of poor QOL, and anxiety highly correlates with depression (Celano et al., 2018; Dekker et al., 2014). Researchers could assess etiology, prevention methods, and strategies to minimize the effects and effectiveness of traditional (i.e., antidepressants, anxiolytics, and psychotherapy) and nontraditional depression and anxiety treatments (i.e., relaxation, meditation, and guided imagery) in patients with HF (Celano et al., 2018). Future scholars could also assess the treatment of patients with HF with antidepressants, compliance with prescribed medication regimens, and the medications prescribed for physical conditions. Finally, further exploring comorbid depression and anxiety with larger samples could contribute to the limited research on comorbid depression and anxiety in the general patient population and Blacks with HF to test for racial differences.

Depression profoundly affects the QOL of patients with HF. Future researchers could investigate depression-contributing factors and mitigation efforts in patients with HF, especially Black patients. Due to emotional and psychological distress, individuals with chronic illnesses (Ahmedani et al., 2017; Erlangsen et al., 2015; Fässberg et al., 2016; Fiske et al., 2008; Kye & Park, 2016; Le Strat et al., 2015; Sampaio et al., 2019), such as cardiovascular diseases (Dunn et al., 2021; Petersen et al., 2020; Suarez et al., 2015) and HF (Celano & Huffman, 2018; Korkmaz et al., 2019; Liu et al., 2018; Lossnitzer et al., 2009; V. C.-C. Wu et al., 2018), are at risk for

suicide ideation, suicide attempts, and death by suicide, especially when faced with untreated and unrelenting depression.

This study found that the Black patients were younger than the White participants, aligning with prior research (Tsao et al., 2022; Virani et al., 2020, 2021). Younger patients are at the greatest risk of depression (Choi et al., 2019; Fiske et al., 2008; Hirsch et al., 2009; Juurlink et al., 2004; V. C.-C. Wu et al., 2018). Additionally, there might be relationships between depression, QOL, and suicidality based on age. Black individuals with HF might have a higher risk of suicide because they often have more comorbidities, and comorbidities increase suicide risk (Juurlink et al., 2004) among patients with chronic illnesses (Sharma et al., 2014). Younger persons who receive disability pensions also have a higher mortality risk from suicide (Jonsson et al., 2013). There is a need for more research on the relationships between depression and suicidality in patients with HF to inform the development and testing of interventions to prevent, identify, and treat depression, avoid suicide risk, and lower suicide prevalence as HF prevalence increases, especially among younger Blacks.

This study did not indicate the importance of the overall QOL and four QOL domains in the QLI to patients or the racial differences in perceived importance. Future researchers should explore whether Black and White patients with HF consider various QOL domains (health and functioning, social and economic, psychological/spiritual, and family) equally important and stratify the findings by race. Assessing the importance of specific QOL aspects could indicate how to develop tailored interventions to improve QOL in Blacks with HF. Black patients with chronic illnesses may perceive the importance of QOL aspects differently from their White counterparts based on their perceptions. Future scholars could also assess the relationships between the four QOL domains and sociodemographic, clinical, and psychological

characteristics to determine whether the relationships differ from assessing total QOL. In addition to examining the contributors to and detractors of QOL in patients with HF, scholars could investigate the psychosocial characteristics contributing to QOL and patients' abilities to adhere to treatment regimens, such as self-management, sexual health and functioning, social support, spirituality, health literacy, depression, anxiety, and QOL in caregivers of patients with HF.

Because of its complexity and chronic nature, HF requires patients to actively engage in effective self-care at home to yield positive clinical outcomes (Jaarsma et al., 2021; Riegel, Dickson, & Faulkner, 2016; Riegel, Moser, et al., 2009). Culture differs among various racial and ethnic groups and can impact HF self-care (Davidson et al., 2007; Dickson, McCarthy, Howe, et al., 2013; Dickson, McCarthy, & Katz, 2013; Jaarsma et al., 2021; Riegel et al., 2009). Cultural influences on self-care, including those related to depression and anxiety, could contribute to persistent racial disparities among Blacks with HF (Dickson, McCarthy, Howe, et al., 2013).

Self-management is an aspect of HF self-care. Effective HF self-management is complex, requiring active engagement and motivation to monitor and manage medications, symptoms, diet, and exercise (Granger, Sandelowski, Tahshjain, Swedberg, & Ekman, 2009; Riegel & Dickson, 2008; Riegel, Lee, & Dickson, 2011; Riegel, Moser, & Anker, et al., 2009). Patients who effectively engage in self-care have less than half the risk of all-cause mortality, hospitalization, or emergency room admission than those less engaged (Lee, Moser, Lennie, & Riegel, 2011). Minorities, individuals with lower SES, and those with lower health literacy are the most vulnerable and at the greatest risk for poor self-management and worse clinical outcomes (Barnason et al., 2012; Murray et al., 2009). Blacks fall into all three high-risk

categories. Although patients might know about the prescribed HF self-management regimen, they could struggle to avoid exacerbation, clinical status declines, and hospitalization when they are depressed and anxious (Granger, Sandelowski, Tahshjain, Swedberg, & Ekman, 2009).

High healthcare utilization and increased mortality among patients with HF, especially Black patients, might correlate with undiagnosed and untreated depression and anxiety. Patients could struggle to implement the prescribed complex HF self-management regimens when depressed and anxious (DiMatteo et al., 2000; Mead et al., 2010; Riegel, Moser, et al., 2009). Scholars could research relationships between self-management, depression, anxiety, and QOL with a focus on racial differences to decrease healthcare costs, enhance QOL, decrease racial disparities, and improve other clinical outcomes. Future investigators could also consider examining relationships between depression, anxiety, and QOL and other aspects of HF self-care, including symptom perception and self-care maintenance.

Sexual health is an important QOL aspect addressed in the QLI. Sexual health and functioning affect health, well-being, and QOL among patients with chronic illnesses, including HF (Baert et al., 2019; Westlake et al., 1999). However, sexual functioning has received minimal focus in HF care, management, and research. Sexual dysfunction correlates with depression in the general population (Andrews et al., 2017; Atlantis & Sullivan, 2012; Fabre & Smith, 2012; Yazdanpanahi et al., 2018) and patients with chronic illnesses (Bak et al., 2017; Basson et al., 2010; Cichocka et al., 2020; Corona et al., 2016; Esen et al., 2015; Heiden-Rootes et al., 2017; McCabe et al., 2016; Slack & Aziz, 2020; Yazici et al., 2009). Additionally, sexual dysfunction contributes to poor QOL in individuals with chronic illnesses (Collins et al., 2012; Fernandes et al., 2010; Schwarz et al., 2008; Senol, 2018; Tański et al., 2022; Tutoglu et al., 2014; Yazici et al., 2009). Some antidepressants contribute to sexual dysfunction (Baldwin, 2001; Baldwin &

Foong, 2013; Basson & Gilks, 2018; Heiden-Rootes et al., 2017; McCabe et al., 2016; Montejo et al., 2018; Slack & Aziz, 2020). Therefore, treating depression with antidepressants might significantly contribute to a cyclical pattern of worsening depression and sexual dysfunction.

HF's complex negative physiological and psychological effects cause many patients to experience sexual dysfunction (da Silva et al., 2022; Jaarsma, 2017; Schwarz et al., 2008; Sztajzel, 2015), which contributes to poor QOL (Baert et al., 2019; Freitas et al., 2006). Sexual health is important to patients with HF, and sexual dysfunction is concerning (Fischer & Bekelman, 2017; Schwarz et al., 2008). Sexual dysfunction in younger individuals and noncompliance to prescribed regimens for chronic illnesses indicate masked depression (Shetty et al., 2018). Black patients with HF are likelier to be younger than White patients (Tsao et al., 2022; Virani et al., 2020, 2021). Therefore, sexual dysfunction could have greater implications for Black persons with HF and could be a major contributor to depression, anxiety, and poor QOL.

Patients with HF want to discuss their sexual health (Baert et al., 2019; van Driel et al., 2014), but healthcare professionals struggle to integrate sexual health into chronic illness management (Fennell & Grant, 2019; Haesler et al., 2016; McGrath et al., 2021), including for those with HF (Hoekstra et al., 2012; Kolbe et al., 2016). Sexual health changes could contribute to depression and poor QOL in patients with HF. Healthcare professionals should provide opportunities for patients to discuss and learn about their sexual health while living with HF and its potential activity and physical limitations.

Future scholars could explore the relationships between sexual health, depression, anxiety, and QOL and assess them for racial differences. Investigators could also explore barriers and facilitators to effective communication and patient sexual education among healthcare

professionals. Researchers could also test interventions related to improving the sexual health of patients with HF to decrease depression and improve QOL.

Patients lacking adequate social support are likelier to experience adverse physical and mental health, including poor coping abilities, depression, and inadequate self-care (Bean et al., 2009; de Leon et al., 2009; Dickson, McCarthy, & Katz, 2013; Evangelista, Ter-Galstanyan, S. Moughrabi, & Moser, 2009; Rohyans & Pressler, 2009; Trivedi et al., 2009). Social support is a way to prevent and reduce depression (Graven & Grant, 2013; Riegel, Moser, et al., 2009; Trivedi et al., 2009), as loneliness correlates with depression (Brouwers et al., 2014). The supporter's relationship to the patient or living arrangements might not indicate support efficacy; however, human beings who act as confidants and discuss depression and other symptoms can increase patients' ability to improve their coping mechanisms and self-care (Bean et al., 2009; de Leon et al., 2009; Dickson, McCarthy, & Katz, 2013; Evangelista, Ter-Galstanyan, Moughrabi, & Moser, 2009). A lack of social support also correlates with poor QOL. Social support might not modify the effects of HF's physical limitations; however, it contributes to mitigating depression, improving QOL (de Leon et al., 2009) and patients' ability to effectively cope with symptoms (Bean et al., 2009; Chung, Moser, Lennie, & Frazier, 2013; Chung, Moser, Lennie, & Rayens, 2009; de Leon et al., 2009; Dekker et al., 2014; Dickson, McCarthy, & Katz, 2013; Evangelista, Ter-Galstanyan, Moughrabi, & Moser, 2009; Rohyans & Pressler, 2009; Trivedi et al., 2009).

There might be racial, ethnic, and cultural differences in the relationships between depression, anxiety, QOL, and social support. Dickson, McCarthy, and Katz (2013) found that Black patients with HF rarely relied on healthcare providers for support, instead relying on family and friends of the same culture. Future researchers could explore the influence of social

support on depression and anxiety and whether there are differences between Black and White patients with HF. Scholars could also explore how to use eHealth and mHealth interventions to improve social support for patients (Mohr et al., 2014), especially Blacks with HF. Around-the-clock, indefinite access to healthcare professionals or caregivers might not be feasible, but technology could enable social interactions among patients from various locales. Thus, technology could enable patients to engage with others who understand their condition and interact with healthcare providers. Black individuals could use technology to interact with people of the same race who could provide culturally driven social support (Dickson, McCarthy, & Katz, 2013) for decreased depression and anxiety (Cornelius et al., 2022).

Culturally responsive eHealth interventions (Latulippe et al., 2017), such as an online social networking platform, or mHealth modalities, such as text messaging and mobile applications, could be means of socially supporting Black patients with HF (Vorderstrasse et al., 2016), decreasing depression and anxiety, and improving QOL. However, Black patients remain underrepresented in eHealth and mHealth studies (James et al., 2017). Before designing, developing, and testing technological interventions, scholars could assess Black individuals' willingness, perceptions, preferences, and priorities for using technology for social support. Scholars could research the technological modalities patients are likelier to use and the sociodemographic (Riegel, Moser, et al., 2009), clinical, or psychosocial factors that could impact use and perceptions about technology (Guzman-Clark et al., 2013; Peeters et al., 2013). Understanding the perspectives of Black patients with HF about using technology to improve social support could indicate how to design and develop interventions (Peeters et al., 2013) to improve acceptability and adoption. Such interventions could be ways to decrease depression and anxiety, improve social support and QOL, and reduce disparities among Blacks with HF.

Caregivers of patients with HF often serve as support systems. However, there has been growing concern for the mental and emotional well-being of caregivers of patients with HF. More research should occur to determine the depression, anxiety, and QOL of caregivers of patients with HF. Living with a family member with HF is challenging and could negatively impact family and caregiver physical and psychosocial well-being (Bahrami et al., 2014; Dracup et al., 2004; Ghasemi et al., 2020; Lum et al., 2014; Whittingham et al., 2013). Patients with HF and their spousal caregivers might have similar depression and anxiety (43%–57%; Chung et al., 2009). The emotional well-being of patients with HF also impacts their caregivers' emotional well-being (Evangelista, Dracup, Doering, et al., 2002). Caregivers of patients with HF require support to adequately function in their caregiving roles (Cameron et al., 2016). However, scholars have not stratified these limited findings by race.

There are also differences in the effects of HF caregiving on caregivers based on age, race, employment status, financial position, family dynamics, social support, and other factors (Dracup et al., 2004; Grigorovich et al., 2017; Hu et al., 2016; Saunders, 2008; Usher & Cammarata, 2009). However, scholars have not stratified their findings by race, and there is little research on the experiences of Black caregivers of Black patients with HF. Future research could focus on the influence of depression, anxiety, and QOL on caregivers of patients with HF and address potential racial differences. Depression and anxiety might contribute to poor QOL, morbidity, and mortality among caregivers of patients with HF. These adverse impacts are especially problematic for Blacks, who already have high rates of chronic illness morbidity and mortality (Doshi et al., 2016; Ellis et al., 2020; Ward & Schiller, 2013). However, there are effective interventions for caregiver burden (Etemadifar et al., 2014). More research and

interventions could indicate how to interrupt this pattern among Black caregivers of Black patients with HF.

Spirituality also impacts QOL (Abu et al., 2018; Beery et al., 2002; Borges et al., 2021; Burlacu et al., 2019; Counted et al., 2018; Koenig, 2004; Melo et al., 2015; Panzini et al., 2017; Sawatzky et al., 2005) and is one of the QOL domains assessed in the QLI (Ferrans, 1990; Ferrans & Powers, 1985). Patients with chronic illnesses (Delgado, 2007; Roger & Hatala, 2018), particularly Black patients (Becker & Newsom, 2005; Chin et al., 2000; Coats, 2015; Conner & Eller, 2004; Gibson & Hendricks, 2006; Holt et al., 2009; Siler et al., 2021; Warren-Findlow & Issel, 2010), rely on spirituality to cope and adjust (Coats et al., 2017) and decrease the negative effects of depression and anxiety (Chin et al., 2000; Loeb, 2006).

Spirituality can also positively impact depression (Bekelman, Dy, et al., 2007; Clark & Hunter, 2019; Mills et al., 2015; Westlake & Dracup, 2001) and anxiety (Bean et al., 2009; Clark & Hunter, 2019; Westlake et al., 2002) in patients with HF. However, few studies have addressed the relationships between spirituality and anxiety in patients with HF. Spiritual well-being has a significant role in illness perceptions and coping with sadness and loss. Additionally, spiritual well-being closely correlates with psychological well-being and responses to depression and anxiety. Spiritual well-being could be of greater importance to the psychological health of Black individuals with HF (Bean et al., 2009; Dickson et al., 2013), as Black persons often rely on their faith to protect against and mitigate depression (Bean et al., 2009; Dickson, McCarthy, Howe, et al., 2013).

Elhag et al. (2022) found relationships between spirituality and medication adherence in patients with cardiovascular disease, including HF. Thus, spirituality could support self-management activities and might correlate to less depression and anxiety and adequate QOL.

Scholars have measured and conceptualized spirituality in patients with HF in various ways (Beery et al., 2002; Clark & Hunter, 2019). Interventions in which spirituality is a means of improving depression, anxiety, and QOL could be effective (Naghi et al., 2012), especially among Black patients. Such interventions could be a way to decrease disparities and improve clinical outcomes among Black individuals. However, there is a need for more research to understand the relationships and their influence.

Lower health literacy contributes to negative clinical outcomes (Cajita et al., 2016; Fabbri et al., 2020; Liu et al., 2019; Moser et al., 2015; Peterson et al., 2011), including lower QOL (Macabasco-O'Connell et al., 2011) and greater mortality (Fabbri et al., 2020; Liu et al., 2019; McNaughton et al., 2015; Moser et al., 2015) in patients with HF. Scholars could explore lower health literacy and racial differences in psychosocial characteristics. Lower health literacy correlates with poor adherence to prescribed HF regimens (Cajita et al., 2016; Deek et al., 2020), greater depression and anxiety, and low QOL. Therefore, patients with HF could benefit from clinicians regularly assessing health literacy as often as vital signs (Evangelista, Rasmusson, et al., 2010; Weiss et al., 2005). Rather than focusing only on educational status, providers could improve depression, anxiety, and QOL in patients with HF by assessing and improving health literacy, especially in Black patients, who often have lower health literacy (Chaudhry et al., 2011; Morrow et al., 2006; Westlake et al., 2013). Future scholars could develop and test interventions, including those with technology, to improve health literacy in patients with HF (Bas-Sarmiento et al., 2022; Deek et al., 2020; Friel, 2016; Westlake et al., 2013). Other social determinants of health, including economic stability, neighborhood and physical environment, community and social context, and the healthcare system, have implications for the QOL of

patients with HF (Javed et al., 2022). This study did not address these concepts at length, indicating the need for future research.

Alignment With National Institute for Nursing Research

This study's foci and related directions for future research aligned with much-needed HF health disparities research and the National Institute for Nursing Research's (NINR) Strategic Plan for the next 4 years. This body of work examines social determinants of health, such as SES; proposes a focus on population and community-level interventions to address critical health challenges among patients with HF, including those that would leverage technology; focuses on tertiary prevention strategies among patients with HF by reducing disease severity, symptoms, and progression by considering relationships between sociodemographic, clinical, and psychosocial characteristics, and has the potential to reduce and ultimately eliminate systemic and structural inequities among Black patients with HF (NINR, 2022).

Advancing Health Disparities in Heart Failure Research

Structural racism is a social determinant of health (Nardi et al., 2020). More research on racial disparities among Black and White persons with HF could indicate how to improve HF racial disparities and usher in Phase IV: Reduced Mortality of the HF racial disparities framework (see Figure 3). Phase IV focuses on the wide availability of prevention and treatment methods for all patients with HF, especially Black patients, to significantly reduce or eliminate racial disparities (Doll, 2018). Although structural (Churchwell et al., 2020) and cultural (Cogburn, 2019) racism, including implicit bias (Nayak et al., 2020), have contributed to racial inequities and health disparities among Blacks with HF, it is possible to overcome these negative outcomes (Sharma et al., 2014). Researchers who investigate phenomena, recruit sufficient Black samples, and make comparisons by race with analytic approaches have contributed to the limited

knowledge (Wei et al., 2022) of racial disparities among Blacks with HF. The increased research is a charge for the scientific community and has implications for society regarding current and projected HF care costs, morbidity, and mortality (Benjamin, Blaha, et al., 2017; Benjamin, Muntner, et al., 2019; Benjamin, Virani, et al., 2018; Cook et al., 2013; Go et al., 2013; Heidenreich, Albert, et al., 2013; Heidenreich, Fonarow, et al., 2022; Lesyuk et al., 2018; Mensah & Dunbar, 2006; Mensah et al., 2005; Mensah et al., 2019; Tsao et al., 2022; Virani et al., 2020, 2021).

References

- Abu, H. O., Ulbricht, C., Ding, E., Allison, J. J., Salmoirago-Blotcher, E., Goldberg, R. J., & Kiefe, C. I. (2018). Association of religiosity and spirituality with quality of life in patients with cardiovascular disease: A systematic review. *Quality of Life Research*, 27(11), 2777–2797. <https://doi.org/10.1007/s11136-018-1906-4>
- AbuRuz, M. E. (2018). Anxiety and depression predicted quality of life among patients with heart failure. *Journal of Multidisciplinary Healthcare*, 11, 367–373. <https://doi.org/10.2147/JMDH.S170327>
- Afzal, A., Ananthasubramaniam, K., Sharma, N., Ai-Malki, Q., Ali, A. S., Jacobsen, G., & Jafri, S. M. (1999). Racial differences in patients with heart failure. *Clinical Cardiology*, 22(12), 791–794. <https://doi.org/10.1002/clc.4960221207>
- Aggelopoulou, Z., Fotos, N. V., Chatziefstratiou, A. A., Giakoumidakis, K., Elefsiniotis, I., & Brokalaki, H. (2017). The level of anxiety, depression and quality of life among patients with heart failure in Greece. *Applied Nursing Research*, 34, 52–56. <https://doi.org/10.1016/j.apnr.2017.01.003>
- Ahmed, A., Ali, M., Lefante, C. M., Mullick, M. S., & Kinney, F. C. (2006). Geriatric heart failure, depression, and nursing home admission: An observational study using propensity score analysis. *The American Journal of Geriatric Psychiatry*, 14(10), 867–875. <https://doi.org/10.1097/01.jgp.0000209639.30899.72>

Ahmedani, B. K., Peterson, E. L., Hu, Y., Rossom, R. C., Lynch, F., Lu, C. Y., Waitzfelder, B. E., Owen-Smith, A. A., Hubley, S., Prabhakar, D., Williams, L. K., Zeld, N., Mutter, E., Beck, A., Tolsma, D., & Simon, G. E. (2017). Major physical health conditions and risk of suicide. *American Journal of Preventive Medicine*, *53*(3), 308–315.

<https://doi.org/10.1016/j.amepre.2017.04.001>

Alegría, M., Chatterji, P., Wells, K., Cao, Z., Chen, C.-N., Takeuchi, D., Jackson, J., & Meng, X.-L. (2008). Disparity in depression treatment among racial and ethnic minority populations in the United States. *Psychiatric Services*, *59*(11), 1264–1272.

<https://doi.org/10.1176/ps.2008.59.11.1264>

Alhurani, A. S., Dekker, R. L., Abed, M. A., Khalil, A., Al Zaghal, M. H., Lee, K. S., Mudd-Martin, G., Biddle, M. J., Lennie, T. A., & Moser, D. K. (2015). The association of comorbid symptoms of depression and anxiety with all-cause mortality and cardiac rehospitalization in patients with heart failure. *Psychosomatics*, *56*(4), 371–380.

<https://doi.org/10.1016/j.psych.2014.05.022>

Allen, L. A., Yager, J. E., Funk, M. J., Levy, W. C., Tulskey, J. A., Bowers, M. T., Dodson, G. C., O'Connor, C. M., & Felker, G. M. (2008). Discordance between patient-predicted and model-predicted life expectancy among ambulatory patients with heart failure. *JAMA*, *299*(21), 2533–2542.

<https://doi.org/10.1001/jama.299.21.2533>

Amabile, C. M., & Spencer, A. P. (2004). Keeping your patient with heart failure safe: A review of potentially dangerous medications. *Archives of Internal Medicine*, *164*(7), 709–720.

<https://doi.org/10.1001/archinte.164.7.709>

- Andrews, A. M., Russell, C. L., & Cheng, A.-L. (2017). Medication adherence interventions for older adults with heart failure: A systematic review. *Journal of Gerontological Nursing*, 43(10), 37–45. <https://doi.org/10.3928/00989134-20170523-01>
- Andrews, J., Ziegelmann, M., Alom, M., Hebert, K., Westerman, M. E., & Trost, L. (2017). MP84-09 symptoms of depression and sexual dysfunction. *The Journal of Urology*, 197(4), e1142–e1143. <https://doi.org/10.1016/j.juro.2017.02.2650>
- Assari, S. (2019). Race, depression, and financial distress in a nationally representative sample of American adults. *Brain Sciences*, 9(2), Article 29. <https://doi.org/10.3390/brainsci9020029>
- Assari, S., Preiser, B., & Kelly, M. (2018). Education and income predict future emotional well-being of Whites but not Blacks: A ten-year cohort. *Brain Sciences*, 8(7), Article 122. <https://doi.org/10.3390/brainsci8070122>
- Athilingam, P., Jenkins, B., Johansson, M., & Labrador, M. A. (2017). Mobile health intervention to improve self-care in patients with heart failure: Pilot randomized control trial. *JMIR Cardio*, 1(2), Article e3. <https://doi.org/10.2196/cardio.7848>
- Atlantis, E., & Sullivan, T. (2012). Bidirectional association between depression and sexual dysfunction: A systematic review and meta-analysis. *Journal of Sexual Medicine*, 9(6), 1497–1507. <https://doi.org/10.1111/j.1743-6109.2012.02709.x>
- Baert, A., Pardaens, S., De Smedt, D., Puddu, P. E., Ciancarelli, M. C., Dawodu, A., De Sutter, J., De Bacquer, D., & Clays, E. (2019). Sexual activity in heart failure patients: Information needs and association with health-related quality of life. *International Journal of Environmental Research and Public Health*, 16(9), Article 1570. <https://doi.org/10.3390/ijerph16091570>

- Bahrami, M., Etemadifar, S., Shahriari, M., & Farsani, A. K. (2014). Caregiver burden among Iranian heart failure family caregivers: A descriptive, exploratory, qualitative study. *Iranian Journal of Nursing and Midwifery Research*, *19*(1), 56–63.
- Bąk, E., Marcisz, C., Krzemińska, S., Dobrzym-Matusiak, D., Foltyn, A., & Drosdzol-Cop, A. (2017). Relationships of sexual dysfunction with depression and acceptance of illness in women and men with type 2 diabetes mellitus. *International Journal of Environmental Research and Public Health*, *14*(9), Article 1073. <https://doi.org/10.3390/ijerph14091073>
- Baldwin, D. S. (2001). Depression and sexual dysfunction. *British Medical Bulletin*, *57*(1), 81–99. <https://doi.org/10.1093/bmb/57.1.81>
- Baldwin, D. S., & Foong, T. (2013). Antidepressant drugs and sexual dysfunction. *British Journal of Psychiatry*, *202*(6), 396–397. <https://doi.org/10.1192/bjp.bp.112.110650>
- Barbareschi, G., Sanderman, R., Lesman-Leegte, I., Van Veldhuisen, D. J., & Jaarsma, T. (2011). Educational level and the quality of life of heart failure patients: A longitudinal study. *Journal of Cardiac Failure*, *17*(1), 47–53. <https://doi.org/10.1016/j.cardfail.2010.08.005>
- Barnason, S., Zimmerman, L., & Young, L. (2012). An integrative review of interventions promoting self-care of patients with heart failure. *Journal Clinical Nursing*, *21*(3–4), 448–475. <https://doi.org/10.1111/j.1365-2702.2011.03907.x>
- Barnes, D. M., & Bates, L. M. (2017). Do racial patterns in psychological distress shed light on the Black–White depression paradox? A systematic review. *Social Psychiatry and Psychiatric Epidemiology*, *52*(8), 913–928. <https://doi.org/10.1007/s00127-017-1394-9>

- Bas-Sarmiento, P., Fernández-Gutiérrez, M., Poza-Méndez, M., Marín-Paz, A. J., Paloma-Castro, O., & Romero-Sánchez, J. M. (2022). Development and effectiveness of a mobile health intervention in improving health literacy and self-management of patients with multimorbidity and heart failure: Protocol for a randomized controlled trial. *JMIR Research Protocols*, *11*(4), Article e35945. <https://doi.org/10.2196/35945>
- Basson, R., & Gilks, T. (2018). Women's sexual dysfunction associated with psychiatric disorders and their treatment. *Women's Health*, *14*, Article 1745506518762664. <https://doi.org/10.1177/1745506518762664>
- Basson, R., Rees, P., Wang, R., Montejo, A. L., & Incrocci, L. (2010). Sexual function in chronic illness. *Journal of Sexual Medicine*, *7*(1), 374–388. <https://doi.org/10.1111/j.1743-6109.2009.01621.x>
- Bean, M. K., Gibson, D., Flattery, M., Duncan, A., & Hess, M. (2009). Psychosocial factors, quality of life, and psychological distress: Ethnic differences in patients with heart failure. *Progress in Cardiovascular Nursing*, *24*(4), 131–140. <https://doi.org/10.1111/j.1751-7117.2009.00051.x>
- Becker, G., & Newsom, E. (2005). Resilience in the face of serious illness among chronically ill African Americans in later life. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, *60*(4), S214–S223. <https://doi.org/10.1093/geronb/60.4.S214>
- Beery, T. A., Baas, L. S., Fowler, C., & Allen, G. (2002). Spirituality in persons with heart failure. *Journal of Holistic Nursing*, *20*(1), 5–25. <https://doi.org/10.1177/089801010202000102>

Bekelman, D. B., Dy, S. M., Becker, D. M., Wittstein, I. S., Hendricks, D. E., Yamashita, T. E., & Gottlieb, S. H. (2007). Spiritual well-being and depression in patients with heart failure. *Journal of General Internal Medicine*, 22(4), 470–477.

<https://doi.org/10.1007/s11606-006-0044-9>

Bekelman, D. B., Havranek, E. P., Becker, D. M., Kutner, J. S., Peterson, P. N., Wittstein, I. S., Gottlieb, S. H., Yamashita, T. E., Fairclough, D. L., & Dy, S. M. (2007). Symptoms, depression, and quality of life in patients with heart failure. *Journal of Cardiac Failure*, 13(8), 643–648. <https://doi.org/10.1016/j.cardfail.2007.05.005>

Benjamin, E. J., Blaha, J. M., Chiuve, E. S., Cushman, R. M., Das, D. S., Deo, R. R., De Ferranti, C. S., Floyd, C. J., Fornage, E. M., Gillespie, H. C., Isasi, T. C., Jiménez, H. M., Jordan, E. L., Judd, W. S., Lackland, K. D., Lichtman, R. J., Lisabeth, J. L., Liu, J. S., Longenecker, A. C., ... Muntner, S. P. (2017). Heart disease and stroke statistics—2017 update: A report from the American Heart Association. *Circulation*, 135(10), e146–e603.

<https://doi.org/10.1161/CIR.0000000000000485>

Benjamin, E. J., Muntner, P., Alonso, A., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Chang, A. R., Cheng, S., Das, S. R., Delling, F. N., Djousse, L., Elkind, M. S. V., Ferguson, J. F., Fornage, M., Jordan, L. C., Khan, S. S., Kissela, B. M., Knutson, K. L., ... Virani, S. S. (2019). Heart disease and stroke statistics—2019 update: A report from the American Heart Association. *Circulation*, 139(10), e56–e66.

<https://doi.org/10.1161/CIR.0000000000000659>

Benjamin, E. J., Virani, S. S., Callaway, C. W., Chamberlain, A. M., Chang, A. R., Cheng, S., Chiuve, S. E., Cushman, M., Dellinger, F. N., Deo, R., de Ferranti, S. D., Ferguson, J. F., Fornage, M., Gillespie, C., Isasi, C. R., Jiménez, M. C., Jordan, L. C., Judd, S. E., Lackland, D., ... Muntner, P. (2018). Heart disease and stroke statistics—2018 update: A report from the American Heart Association. *Circulation*, *137*(12), e67–e492.

<https://doi.org/10.1161/CIR.0000000000000558>

Bennett, S. J., & Sauvé, M. J. (2003). Cognitive deficits in patients with heart failure: A review of the literature. *The Journal of Cardiovascular Nursing*, *18*(3), 219–242.

<https://doi.org/10.1097/00005082-200307000-00007>

Bhatt et al., 2016

Bhopal, R., Gruer, L., Agyemang, C., Davidovitch, N., de-Graft Aikins, A., Krasnik, A., Martinez-Donate, A. P., Miranda, J. J., Pottie, K., Segal, U., Zwi, A., & Kumar, B. (2021). The global society on migration, ethnicity, race and health: Why race can't be ignored even if it causes discomfort. *European Journal of Public Health*, *31*(1), 3–4.

<https://doi.org/10.1093/eurpub/ckaa191>

Bibbins-Domingo, K., Pletcher, M. J., Lin, F., Vittinghoff, E., Gardin, J. M., Arynchyn, A., Lewis, C. E., Williams, O. D., & Hulley, S. B. (2009). Racial differences in incident heart failure among young adults. *New England Journal of Medicine*, *360*(12), 1179–1190.

<https://doi.org/10.1056/NEJMoa0807265>

Bielderman, A., de Greef, M., Krijnen, W., & van der Schans, C. (2015). Relationship between socioeconomic status and quality of life in older adults: a path analysis. *Quality of Life Research*, *24*(7), 1697–1705. <https://doi.org/10.1007/s11136-014-0898-y>

- Billingsley, G. (2014). Black community involvement as a strategy for overcoming barriers to African-American participation in health research—What is community involvement? *Race, Gender & Class, 21*(3/4), 122–137.
- Bjelland, I., Krokstad, S., Mykletun, A., Dahl, A. A., Tell, G. S., & Tambs, K. (2008). Does a higher educational level protect against anxiety and depression? The HUNT study. *Social Science & Medicine, 66*(6), 1334–1345. <https://doi.org/10.1016/j.socscimed.2007.12.019>
- Blinderman et al., 2008
- Blumenthal, J. A., Babyak, M. A., O'Connor, C., Keteyian, S., Landzberg, J., Howlett, J., Kraus, W., Gottlieb, S., Blackburn, G., Swank, A., & Whellan, D. J. (2012). Effects of exercise training on depressive symptoms in patients with chronic heart failure. *JAMA, 308*(5), 565–474. <https://doi.org/10.1001/jama.2012.8720>
- Borges, C. C., dos Santos, P. R., Alves, P. M., Borges, R. C. M., Lucchetti, G., Barbosa, M. A., Porto, C. C., & Fernandes, M. R. (2021). Association between spirituality/religiousness and quality of life among healthy adults: A systematic review. *Health and Quality of Life Outcomes, 19*(1), Article 246. <https://doi.org/10.1186/s12955-021-01878-7>
- Bozkurt, B., & Khalaf, S. (2017). Heart failure in women. *Methodist DeBakey Cardiovascular Journal, 13*(4), 216–223. <https://doi.org/10.14797/mdcj-13-4-216>
- Branson, R. D., Davis, K., Jr., & Butler, K. L. (2007). African Americans' participation in clinical research: Importance, barriers, and solutions. *The American Journal of Surgery, 193*(1), 32–39. <https://doi.org/10.1016/j.amjsurg.2005.11.007>

- Brouwers, C., Kupper, N., Pelle, A. J., Szabó, B. M., Westerhuis, B. L., & Denollet, J. (2014). Depressive symptoms in outpatients with heart failure: Importance of inflammatory biomarkers, disease severity and personality. *Psychology & Health, 29*(5), 564–582. <https://doi.org/10.1080/08870446.2013.869813>
- Brouwers, C., Mommersteeg, P. M., Nyklicek, I., Pelle, A. J., Westerhuis, B. L., Szabo, B. M., & Denollet, J. (2013). Positive affect dimensions and their association with inflammatory biomarkers in patients with chronic heart failure. *Biological Psychology, 92*(2), 220–226. <https://doi.org/10.1016/j.biopsycho.2012.10.002>
- Browne, D. C., Gross, S. M., & Scott-Johnson, P. E. (2010). Web-based data collection: An effective strategy for increasing African Americans' participation in health-related research. *Ethnicity & Disease, 20*(1), 201–206.
- Burlacu, A., Artene, B., Nistor, I., Buju, S., Jugrin, D., Mavrichi, I., & Covic, A. (2019). Religiosity, spirituality and quality of life of dialysis patients: A systematic review. *International Urology and Nephrology, 51*(5), 839–850. <https://doi.org/10.1007/s11255-019-02129-x>
- Byrd, G. S., Edwards, C. L., Kelkar, V. A., Phillips, R. G., Byrd, J. R., Pirn-Pong, D. S., Starks, T. D., Taylor, A. L., Mckinley, R. E., Li, Y. J., & Pericak-Vance, M. (2011). Recruiting intergenerational African American males for biomedical research studies: A major research challenge. *Journal of the National Medical Association, 103*(6), 480–487. [https://doi.org/10.1016/S0027-9684\(15\)30361-8](https://doi.org/10.1016/S0027-9684(15)30361-8)
- Cajita, M. I., Cajita, T. R., & Han, H.-R. (2016). Health literacy and heart failure: A systematic review. *The Journal of Cardiovascular Nursing, 31*(2), 121–130. <https://doi.org/10.1097/JCN.0000000000000229>

Cameron, J., Rhodes, K. L., Ski, C. F., & Thompson, D. R. (2016). Carers' views on patient self-care in chronic heart failure. *Journal of Clinical Nursing*, 25(1–2), 144–152.

<https://doi.org/10.1111/jocn.13124>

Canedo, J. R., Miller, S. T., Schlundt, D., Fadden, M. K., & Sanderson, M. (2017). Racial/ethnic disparities in diabetes quality of care: The role of healthcare access and socioeconomic status. *Journal of Racial and Ethnic Health Disparities*, 5(1), 7–14.

<https://doi.org/10.1007/s40615-016-0335-8>

Carels, R. A. (2004). The association between disease severity, functional status, depression and daily quality of life in congestive heart failure patients. *Quality of Life Research*, 13(1), 63–72. <https://doi.org/10.1023/B:QURE.0000015301.58054.51>

Carson, P., Tam, S. W., Ghali, J. K., Archambault, W. T., Taylor, A., Cohn, J. N., Braman, V. M., Worcel, M., & Anand, I. S. (2009). Relationship of quality of life scores with baseline characteristics and outcomes in the African-American Heart Failure Trial. *Journal of Cardiac Failure*, 15(10), 835–842.

<https://doi.org/10.1016/j.cardfail.2009.05.016>

Cascino, T. M., Somanchi, S., Colvin, M., Chung, G. S., Brescia, A. A., Pienta, M., Thompson, M. P., Stewart, J. W., Sukul, D., Watkins, D. C., Pagani, F. D., Likosky, D. S., Aaronson, K. D., & McCullough, J. S. (2022). Racial and sex inequities in the use of and outcomes after left ventricular assist device implantation among Medicare beneficiaries. *JAMA Network Open*, 5(7), Article e2223080.

<https://doi.org/10.1001/jamanetworkopen.2022.23080>

- Cassiani-Miranda, C. A., Rueda, M., & Camacho, P. A. (2022). Diagnostic accuracy of scales for depression screening in patients with heart failure: Systematic review and meta-analysis. *Revista Ciencias Biomédicas*, *11*(2), 127–144. <https://doi.org/10.32997/rcb-2022-3934>
- Celano, C. M., & Huffman, J. C. (2018). Heart failure and suicide: The role of depression. *Journal of Cardiac Failure*, *24*(11), 801–802. <https://doi.org/10.1016/j.cardfail.2018.10.003>
- Celano, C. M., Villegas, A. C., Albanese, A. M., Gaggin, H. K., & Huffman, J. C. (2018). Depression and anxiety in heart failure: A review. *Harvard Review of Psychiatry*, *26*(4), 175–184. <https://doi.org/10.1097/HRP.000000000000162>
- Celik, E., Cay, S., Sensoy, B., Murat, S., Oksuz, F., Cankurt, T., & Mendi, M. A. (2016). Heart failure functional class associated with depression severity but not anxiety severity. *Acta Cardiologica Sinica*, *32*(1), 55–61. <https://doi.org/10.6515/ACS20150509A>
- Charlson, M. E., Carrozzino, D., Guidi, J., & Patierno, C. (2022). Charlson Comorbidity Index: A critical review of clinimetric properties. *Psychotherapy and Psychosomatics*, *91*(1), 8–35. <https://doi.org/10.1159/000521288>
- Chaudhry, S. I., Herrin, J., Phillips, C., Butler, J., Mukerjee, S., Murillo, J., Onwuanyi, A., Seto, T. B., Spertus, J., & Krumholz, H. M. (2011). Racial disparities in health literacy and access to care among patients with heart failure. *Journal of Cardiac Failure*, *17*(2), 122–127. <https://doi.org/10.1016/j.cardfail.2010.09.016>
- Chen, J., Dharmarajan, K., Wang, Y., & Krumholz, H. M. (2013). National trends in heart failure hospital stay rates, 2001 to 2009. *Journal of the American College of Cardiology*, *61*(10), 1078–1088. <https://doi.org/10.1016/j.jacc.2012.11.057>

- Chin, M. H., Polonsky, T. S., Thomas, V. D., & Nerney, M. P. (2000). Developing a conceptual framework for understanding illness and attitudes in older, urban African Americans with diabetes. *The Diabetes Educator*, 26(3), 439–449.
<https://doi.org/10.1177/014572170002600311>
- Choi, N. G., DiNitto, D. M., Marti, C. N., & Conwell, Y. (2019). Physical health problems as a late-life suicide precipitant: Examination of coroner/medical examiner and law enforcement reports. *The Gerontologist*, 59(2), 356–367.
<https://doi.org/10.1093/geront/gnx143>
- Chung, M. L., Lennie, T. A., Riegel, B., Wu, J.-R., Dekker, R. L., & Moser, D. K. (2009). Marital status as an independent predictor of event-free survival of patients with heart failure. *American Journal of Critical Care*, 18(6), 562–570.
<https://doi.org/10.4037/ajcc2009388>
- Chung, M. L., Moser, D. K., Lennie, T. A., & Frazier, S. K. (2013). Perceived social support predicted quality of life in patients with heart failure, but the effect is mediated by depressive symptoms. *Quality of Life Research*, 22(7), 1555–1563.
<https://doi.org/10.1007/s11136-012-0294-4>
- Chung, M. L., Moser, D. K., Lennie, T. A., & Rayens, M. K. (2009). The effects of depressive symptoms and anxiety on quality of life in patients with heart failure and their spouses: Testing dyadic dynamics using actor-partner interdependence model. *Journal of Psychosomatic Research*, 67(1), 29–35. <https://doi.org/10.1016/j.jpsychores.2009.01.009>

- Churchwell, K., Elkind, M. S., Benjamin, R. M., Carson, A. P., Chang, E. K., Lawrence, W., Mills, A., Odom, T. M., Rodriguez, C. J., & Rodriguez, F. (2020). Call to action: Structural racism as a fundamental driver of health disparities: A presidential advisory from the American Heart Association. *Circulation*, *142*(24), e454–e468. <https://doi.org/10.1161/CIR.0000000000000936>
- Cichocka, E., Jagusiewicz, M., & Gumprecht, J. (2020). Sexual dysfunction in young women with type 1 diabetes. *International Journal of Environmental Research and Public Health*, *17*(12), Article 4468. <https://doi.org/10.3390/ijerph17124468>
- Clark, C. C., & Hunter, J. (2019). Spirituality, spiritual well-being, and spiritual coping in advanced heart failure: Review of the literature. *Journal of Holistic Nursing*, *37*(1), 56–73. <https://doi.org/10.1177/0898010118761401>
- Clark, D. O., Tu, W., Weiner, M., & Murray, M. D. (2003). Correlates of health-related quality of life among lower-income, urban adults with congestive heart failure. *Heart & Lung*, *32*(6), 391–401. <https://doi.org/10.1016/j.hrtlng.2003.07.005>
- Clarke, S. P., Frasure-Smith, N., Lespérance, F., & Bourassa, M. G. (2000). Psychosocial factors as predictors of functional status at 1 year in patients with left ventricular dysfunction. *Research in Nursing & Health*, *23*(4), 290–300. [https://doi.org/10.1002/1098-240X\(200008\)23:4<290::AID-NUR5>3.0.CO;2-0](https://doi.org/10.1002/1098-240X(200008)23:4<290::AID-NUR5>3.0.CO;2-0)

- Clover, K., Lambert, S. D., Oldmeadow, C., Britton, B., King, M. T., Mitchell, A. J., & Carter, G. L. (2020). Apples to apples? Comparison of the measurement properties of Hospital Anxiety and Depression-Anxiety Subscale (HADS-A), Depression, Anxiety and Stress Scale – Anxiety subscale (DASS-A), and Generalised Anxiety Disorder (GAD-7) scale in an oncology setting using Rasch analysis and diagnostic accuracy statistics. *Current Psychology*, 41(7), 4592–4601. <https://doi.org/10.1007/s12144-020-00906-x>
- Coats, H., Crist, J. D., Berger, A., Sternberg, E., & Rosenfeld, A. G. (2017). African American elders' serious illness experiences: Narratives of “God did,” “God will,” and “Life is better.” *Qualitative Health Research*, 27(5), 634–648. <https://doi.org/10.1177/1049732315620153>
- Coats, H. L. (2015). *African American elders' psycho-social-spiritual healing across serious illness* [Doctoral dissertation, The University of Arizona]. UA Theses and Dissertations. <https://repository.arizona.edu/handle/10150/578887>
- Coelho, R., Ramos, S., Prata, J., Bettencourt, P., Ferreira, A., & Cerqueira-Gomes, M. (2005). Heart failure and health related quality of life. *Clinical Practice and Epidemiology in Mental Health*, 1(1), Article 19. <https://doi.org/10.1186/1745-0179-1-19>
- Cogburn, C. D. (2019). Culture, race, and health: Implications for racial inequities and population health. *The Milbank Quarterly*, 97(3), 736–761. <https://doi.org/10.1111/1468-0009.12411>
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112(1), 155–159. <https://doi.org/10.1037/0033-2909.112.1.155>

- Collins, E. G., Halabi, S., Langston, M., Schnell, T., Tobin, M. J., & Laghi, F. (2012). Sexual dysfunction in men with COPD: Impact on quality of life and survival. *Lung*, *190*(5), 545–556. <https://doi.org/10.1007/s00408-012-9398-4>
- Conner, N. E., & Eller, L. S. (2004). Spiritual perspectives, needs and nursing interventions of Christian African-Americans. *Journal of Advanced Nursing*, *46*(6), 624–632. <https://doi.org/10.1111/j.1365-2648.2004.03053.x>
- Cook, B. L., Zuvekas, S. H., Carson, N., Wayne, G. F., Vesper, A., & McGuire, T. G. (2014). Assessing racial/ethnic disparities in treatment across episodes of mental health care. *Health Services Research*, *49*(1), 206–229. <https://doi.org/10.1111/1475-6773.12095>
- Cook, C., Cole, G., Asaria, P., Jabbour, R., & Francis, D. P. (2013). The annual global economic burden of heart failure. *International Journal of Cardiology*, *171*(3), 368–376. <https://doi.org/10.1016/j.ijcard.2013.12.028>
- Corbie-Smith, G., Thomas, S. B., & St. George, D. M. M. (2002). Distrust, race, and research. *Archives of Internal Medicine*, *162*(21), 2458–2463. <https://doi.org/10.1001/archinte.162.21.2458>
- Corbie-Smith, G., Thomas, S. B., Williams, M. V., & Moody-Ayers, S. (1999). Attitudes and beliefs of African Americans toward participation in medical research. *Journal of General Internal Medicine*, *14*(9), 537–546. <https://doi.org/10.1046/j.1525-1497.1999.07048.x>
- Cornelius, J., Whitaker-Brown, C., Smoot, J., Hart, S., Lewis, Z., & Smith, O. (2022). A text messaging-enhanced intervention for African American patients with heart failure, depression, and anxiety (TXT COPE-HF): Protocol for a pilot feasibility study. *JMIR Research Protocols*, *11*(1), Article e32550. <https://doi.org/10.2196/32550>

- Corona, G., Giorda, C. B., Cucinotta, D., Guida, P., & Nada, E. (2016). Sexual dysfunction in type 2 diabetes at diagnosis: Progression over time and drug and non-drug correlated factors. *PLoS ONE*, *11*(10), Article e0157915. <https://doi.org/10.1371/journal.pone.0157915>
- Counted, V., Possamai, A., & Meade, T. (2018). Relational spirituality and quality of life 2007 to 2017: An integrative research review. *Health and Quality of Life Outcomes*, *16*, Article 75. <https://doi.org/10.1186/s12955-018-0895-x>
- Culley, L. (2006). Transcending transculturalism? Race, ethnicity and health-care. *Nursing Inquiry*, *13*(2), 144–153. <https://doi.org/10.1111/j.1440-1800.2006.00311.x>
- Cully, J. A., Phillips, L. L., Kunik, M. E., Stanley, M. A., & Deswal, A. (2010). Predicting quality of life in veterans with heart failure: The role of disease severity, depression, and comorbid anxiety. *Behavioral Medicine*, *36*(2), 70–76. <https://doi.org/10.1080/08964280903521297>
- Cuyjet, A. B., & Akinboboye, O. (2014). Acute heart failure (AHF) in the African American patient. *Journal of Cardiac Failure*, *20*(7), 533–540.
- da Silva, M. L., Costa, N. L., Jacomo, R. H., Alves, A. T., de Melo, R. A., & Cunha, K. D. C. (2022). The impact of heart failure on the sexual response cycle: A systematic review. *Heart & Lung*, *56*, 50–56. <https://doi.org/10.1016/j.hrtlng.2022.05.013>
- Das, A. K., Olfson, M., McCurtis, H. L., & Weissman, M. M. (2006). Depression in African Americans: Breaking barriers to detection and treatment. *Journal of Family Practice*, *55*(1), 30–39. <https://doi.org/10.1016/j.cardfail.2014.04.018>

Das, A. K., Roy, B., Schwarzer, G., Silverman, M. G., Ziegler, O., Bandyopadhyay, D., Philpotts, L. L., Sinha, S., Blumenthal, J. A., & Das, S. (2019). Comparison of treatment options for depression in heart failure: A network meta-analysis. *Journal of Psychiatric Research, 108*, 7–23. <https://doi.org/10.1016/j.jpsychires.2018.10.007>

Davidson et al., 2007

De Bellis, A., De Angelis, G., Fabris, E., Cannatà, A., Merlo, M., & Sinagra, G. (2019). Gender-related differences in heart failure: Beyond the “one-size-fits-all” paradigm. *Heart Failure Review, 25*(2), 245–255. <https://doi.org/10.1007/s10741-019-09824-y>

de Groot, V., Beckerman, H., Lankhorst, G. J., & Bouter, L. M. (2003). How to measure comorbidity: A critical review of available methods. *Journal of Clinical Epidemiology, 56*(3), 221–229. [https://doi.org/10.1016/S0895-4356\(02\)00585-1](https://doi.org/10.1016/S0895-4356(02)00585-1)

De Jong, M. J., Chung, M. L., Wu, J. R., Riegel, B., Rayens, M. K., & Moser, D. K. (2011). Linkages between anxiety and outcomes in heart failure. *Heart Lung, 40*(5), 393–404. <https://doi.org/10.1016/j.hrtlng.2011.02.002>

de Leon, C. F., Grady, K. L., Eaton, C., Rucker-Whitaker, C., Janssen, I., Calvin, J., & Powell, L. H. (2009). Quality of life in a diverse population of patients with heart failure: Baseline findings from the Heart Failure Adherence and Retention Trial (HART). *Journal of Cardiopulmonary Rehabilitation and Prevention, 29*(3), 171–178. <https://doi.org/10.1097/HCR.0b013e31819a0266>

Deek, H., Itani, L., & Davidson, P. M. (2020). Literacy critical to heart failure management: A scoping review. *Heart Failure Review, 26*(6), 1413–1419. <https://doi.org/10.1007/s10741-020-09964-6>

- DeFilippis, E. M., Beale, A., Martyn, T., Agarwal, A., Elkayam, U., Lam, C. S. P., & Hsich, E. (2022). Heart failure subtypes and cardiomyopathies in women. *Circulation Research*, *130*(4), 436–454. <https://doi.org/10.1161/CIRCRESAHA.121.319900>
- Dekker, R. L., Lennie, T. A., Doering, L. V., Chung, M. L., Wu, J. R., & Moser, D. K. (2014). Coexisting anxiety and depressive symptoms in patients with heart failure. *European Journal of Cardiovascular Nursing*, *13*(2), 168–176. <https://doi.org/10.1177/1474515113519520>
- Delgado, C. (2007). Sense of coherence, spirituality, stress and quality of life in chronic illness. *Journal of Nursing Scholarship*, *39*(3), 229–234. <https://doi.org/10.1111/j.1547-5069.2007.00173.x>
- Dennison, C. R., McEntee, M. L., Samuel, L., Johnson, B. J., Rotman, S., Kielty, A., & Russell, S. D. (2011). Adequate health literacy is associated with higher heart failure knowledge and self care confidence in hospitalized patients. *The Journal of Cardiovascular Nursing*, *26*(5), 359–367. <https://doi.org/10.1097/JCN.0b013e3181f16f88>
- Dewan, P., Rørth, R., Jhund, P. S., Ferreira, J. P., Zannad, F., Shen, L., Køber, L., Abraham, W. T., Desai, A. S., Dickstein, K., Packer, M., Rouleau, J. L., Solomon, S. D., Swedberg, K., Zile, M. R., & McMurray, J. J. V. (2019). Income inequality and outcomes in heart failure: A global between-country analysis. *JACC: Heart Failure*, *7*(4), 336–346. <https://doi.org/10.1016/j.jchf.2018.11.005>
- Dickens, C., Dickson, V. V., & Piano, M. R. (2019). Perceived stress among patients with heart failure who have low socioeconomic status: A mixed-methods study. *The Journal of Cardiovascular Nursing*, *34*(3), E1–E8. <https://doi.org/10.1097/JCN.0000000000000562>

- Dickson, V. V., McCarthy, M. M., Howe, A., Schipper, J., & Katz, S. M. (2013). Sociocultural influences on heart failure self-care among an ethnic minority Black population. *Journal of Cardiovascular Nursing*, 28(2), 111–118.
<https://doi.org/10.1097/JCN.0b013e31823db328>
- Dickson, V. V., McCarthy, M. M., & Katz, S. M. (2013). How do depressive symptoms influence self-care among an ethnic minority population with heart failure? *Ethnicity & Disease*, 23(1), 22–28.
- Dickson, V. V., McCauley, L. A., & Riegel, B. (2008). Work–heart balance: The influence of biobehavioral variables on self-care among employees with heart failure. *Workplace Health & Safety*, 56(2), 63–75. <https://doi.org/10.1177/216507990805600203>
- DiMatteo, M. R., Lepper, H. S., & Croghan, T. W. (2000). Depression is a risk factor for noncompliance with medical treatment: Meta-analysis of the effects of anxiety and depression on patient adherence. *Archives of Internal Medicine*, 160(14), 2101–2107.
<https://doi.org/10.1001/archinte.160.14.2101>
- Doll, K. M. (2018). Investigating Black-White disparities in gynecologic oncology: Theories, conceptual models, and applications. *Gynecologic Oncology*, 149(1), 78–83.
<https://doi.org/10.1016/j.ygyno.2017.10.002>
- Doshi, R. P., Aseltine, R. H., Sabina, A. B., & Graham, G. N. (2016). Racial and ethnic disparities in preventable hospitalizations for chronic disease: Prevalence and risk factors. *Journal of Racial and Ethnic Health Disparities*, 4(6), 1100–1106.
<https://doi.org/10.1007/s40615-016-0315-z>

Dracup, K., Evangelista, L. S., Doering, L., Tullman, D., Moser, D. K., & Hamilton, M. (2004).

Emotional well-being in spouses of patients with advanced heart failure. *Heart & Lung*, 33(6), 354–361. <https://doi.org/10.1016/j.hrtlng.2004.06.003>

Dunn, S. L., DeVon, H. A., Collins, E. G., Luong, A., Buursma, M. P., Gutierrez-Kapheim, M.,

& Bronas, U. G. (2021). Suicide risk management protocol for a randomized controlled trial of cardiac patients reporting hopelessness. *Nursing Research*, 70(1), 72–79.

<https://doi.org/10.1097/NNR.0000000000000474>

Durant, R. W., Davis, R. B., George, D. M. M. S., Williams, I. C., Blumenthal, C., & Corbie-

Smith, G. M. (2007). Participation in research studies: Factors associated with failing to meet minority recruitment goals. *Annals of Epidemiology*, 17(8), 634–642.

<https://doi.org/10.1016/j.annepidem.2007.02.003>

Easton, K., Coventry, P., Lovell, K., Carter, L.-A., & Deaton, C. (2016). Prevalence and

measurement of anxiety in samples of patients with heart failure: Meta-analysis. *Journal of Cardiovascular Nursing*, 31(4), 367–379.

<https://doi.org/10.1097/JCN.0000000000000265>

Eisele, M., Harder, M., Rakebrandt, A., Boczor, S., Marx, G., Blozik, E., Träder, J.-M., Störk, S.,

Herrmann-Lingen, C., & Scherer, M. (2020). Association of depression and anxiety with adherence in primary care patients with heart failure-cross-sectional results of the observational RECODE-HF cohort study. *Family Practice*, 37(5), 695–702.

<https://doi.org/10.1093/fampra/cmaa042>

- Elhag, M., Awaisu, A., Koenig, H. G., & Mohamed Ibrahim, M. I. (2022). the association between religiosity, spirituality, and medication adherence among patients with cardiovascular diseases: A systematic review of the literature. *Journal of Religion and Health*, 61(5), 3988–4027. <https://doi.org/10.1007/s10943-022-01525-5>
- Ellis, K. R., Hecht, H. K., Young, T. L., Oh, S., Thomas, S., Hoggard, L. S., Ali, Z., Olawale, R., Carthron, D., Corbie-Smith, G., & Eng, E. (2020). Chronic disease among African American families: A systematic scoping review. *Preventing Chronic Disease*, 17, Article E167. <https://doi.org/10.5888/pcd17.190431>
- Eng, K., & Feeny, D. (2007). Comparing the health of low income and less well educated groups in the United States and Canada. *Population Health Metrics*, 5(1), Article 10. <https://doi.org/10.1186/1478-7954-5-10>
- Erlangsen, A., Stenager, E., & Conwell, Y. (2015). Physical diseases as predictors of suicide in older adults: A nationwide, register-based cohort study. *Social Psychiatry and Psychiatric Epidemiology*, 50(9), 1427–1439. <https://doi.org/10.1007/s00127-015-1051-0>
- Esen, B., Kahvecioglu, S., Atay, A. E., Ozgen, G., Okumus, M. M., Seyahi, N., Sit, D., & Kadioglu, P. N. (2015). Evaluation of relationship between sexual functions, depression and quality of life in patients with chronic kidney disease at predialysis stage. *Renal Failure*, 37(2), 262–267. <https://doi.org/10.3109/0886022X.2014.990348>
- Etemadifar, S., Bahrami, M., Shahriari, M., & Farsani, A. K. (2014). The effectiveness of a supportive educative group intervention on family caregiver burden of patients with heart failure. *Iranian Journal of Nursing and Midwifery Research*, 19(3), 217–223.

Evangelista, L. S., Dracup, K., & Doering, L. V. (2002). Racial differences in treatment-seeking delays among heart failure patients. *Journal of Cardiac Failure*, 8(6), 381–386.

<https://doi.org/10.1054/jcaf.2002.129234>

Evangelista, L. S., Dracup, K., Doering, L., Westlake, C., Fonarow, G. C., & Hamilton, M. (2002). Emotional well-being of heart failure patients and their caregivers. *Journal of Cardiac Failure*, 8(5), 300–305. <https://doi.org/10.1054/jcaf.2002.128005>

Evangelista, L. S., Ter-Galstanyan, A., Moughrabi, S., & Moser, D. K. (2009). Anxiety and depression in ethnic minorities with chronic heart failure. *Journal of Cardiac Failure*, 15(7), 572–579. <https://doi.org/10.1016/j.cardfail.2009.03.005>

Evangelista, L. S., Ter-Galstanyan, A., Moughrabi, S., Moser, D. K., Evangelista, L. S., Ter-Galstanyan, A., Moughrabi, S., & Moser, D. K. (2009). Anxiety and depression in ethnic minorities with chronic heart failure. *Journal of Cardiac Failure*, 15(7), 572–579.

<https://doi.org/10.1016/j.cardfail.2009.03.005>

Evangelista, L. S., Rasmusson, K. D., Laramée, A. S., Barr, J., Ammon, S. E., Dunbar, S., Ziesche, S., Patterson, J. H., & Yancy, C. W. (2010). Health literacy and the patient with heart failure—Implications for patient care and research: A consensus statement of the Heart Failure Society of America. *Journal of Cardiac Failure*, 16(1), 9–16.

<https://doi.org/10.1016/j.cardfail.2009.10.026>

Evangelista, L. S., Ter-Galstanyan, A., Moughrabi, S., & Moser, D. K. (2009). Anxiety and depression in ethnic minorities with chronic heart failure. *Journal of Cardiac Failure*, 15(7), 572–579. <https://doi.org/10.1016/j.cardfail.2009.03.005>

- Fabbri, M., Murad, M. H., Wennberg, A. M., Turcano, P., Erwin, P. J., Alahdab, F., Berti, A., Manemann, S. M., Yost, K. J., Finney Rutten, L. J., & Roger, V. L. (2020). Health literacy and outcomes among patients with heart failure: A systematic review and meta-analysis. *JACC: Heart Failure*, 8(6), 451–460. <https://doi.org/10.1016/j.jchf.2019.11.007>
- Fabre, L. F., & Smith, L. C. (2012). The effect of major depression on sexual function in women. *Journal of Sexual Medicine*, 9(1), 231–239. <https://doi.org/10.1111/j.1743-6109.2011.02445.x>
- Falk, Doll, S., Gummert, J., Jacquet, L.-M., Glineur, D., Fisher, K., Shipkowitz, T., Crane, D., & Belalcazar, A. (2007). Pilot Study of Chronic Monitoring of Left Ventricular Pressure in Heart Failure Patients. *Journal of Cardiac Failure*, 13(6), S119–S119. <https://doi.org/10.1016/j.cardfail.2007.06.466>
- Fässberg, M. M., Cheung, G., Canetto, S. S., Erlangsen, A., Lapierre, S., Lindner, R., Draper, B., Gallo, J. J., Wong, C., Wu, J., Duberstein, P., & Wærn, M. (2016). A systematic review of physical illness, functional disability, and suicidal behaviour among older adults. *Aging & Mental Health*, 20(2), 166–194. <https://doi.org/10.1080/13607863.2015.1083945>
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using GPower 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*, 41(4), 1149–1160. <https://doi.org/10.3758/BRM.41.4.1149>
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). GPower 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191. <https://doi.org/10.3758/BF03193146>

- Fennell, R., & Grant, B. (2019). Discussing sexuality in health care: A systematic review. *Journal of Clinical Nursing*, 28(17–18), 3065–3076. <https://doi.org/10.1111/jocn.14900>
- Fernandes, G. V., Dos Santos, R. R., Soares, W., De Lima, L. G., De Macêdo, B. S., Da Fonte, J. E., De Carvalho, B. S. P., Coelho, S. N., & Calado, A. A. (2010). The impact of erectile dysfunction on the quality of life of men undergoing hemodialysis and its association with depression. *Journal of Sexual Medicine*, 7(12), 4003–4010. <https://doi.org/10.1111/j.1743-6109.2010.01993.x>
- Ferrans, C. E. (1990). Development of a quality of life index for patients with cancer. *Oncology Nursing Forum*, 17(3 Suppl), 15–19.
- Ferrans, C. E., & Powers, M. J. (1985). Quality of Life Index: Development and psychometric properties. *Advances in Nursing Science*, 8(1), 15–24. <https://doi.org/10.1097/00012272-198510000-00005>
- Ferrans, C. E., & Powers, M. J. (1992). Psychometric assessment of the Quality of Life Index. *Research in Nursing & Health*, 15(1), 29–38. <https://doi.org/10.1002/nur.4770150106>
- Ferrans, C. E., Zerwic, J. J., Wilbur, J. E., & Larson, J. L. (2005). Conceptual model of health-related quality of life. *Journal of Nursing Scholarship*, 37(4), 336–342. <https://doi.org/10.1111/j.1547-5069.2005.00058.x>
- Fischer, S., & Bekelman, D. (2017). Gender differences in sexual interest or activity among adults with symptomatic heart failure. *Journal of Palliative Medicine*, 20(8), 890–894. <https://doi.org/10.1089/jpm.2016.0476>
- Fiske, A., O’Riley, A. A., & Widoe, R. K. (2008). Physical health and suicide in late life: An evaluative review. *Clinical Gerontologist*, 31(4), 31–50. <https://doi.org/10.1080/07317110801947151>

- Ford, M. E., Siminoff, L. A., Pickelsimer, E., Mainous, A. G., Smith, D. W., Diaz, V. A., Soderstrom, L. H., Jefferson, M. S., & Tilley, B. C. (2013). Unequal burden of disease, unequal participation in clinical trials: Solutions from African American and Latino community members. *Health & Social Work, 38*(1), 29–38.
<https://doi.org/10.1093/hsw/hlt001>
- Freedland, K. E., Hesseler, M. J., Carney, R. M., Steinmeyer, B. C., Skala, J. A., Davila-Roman, V. G., & Rich, M. W. (2016). Major depression and long-term survival of patients with heart failure. *Psychosomatic Medicine, 78*(8), 896–903.
<https://doi.org/10.1097/PSY.0000000000000346>
- Freedland, K. E., Carney, R. M., & Rich, M. W. (2011). Effect of depression on prognosis in heart failure. *Heart Failure Clinics, 7*(1), 11–21.
<https://doi.org/10.1016/j.hfc.2010.08.003>
- Freedland, K. E., Carney, R. M., Steinmeyer, B. C., Skala, J. A., & Rich, M. W. (2021). Left ventricular dysfunction and depression in hospitalized patients with heart failure. *Psychosomatic Medicine, 83*(3), 274–282.
<https://doi.org/10.1097/PSY.0000000000000915>
- Freedland, K. E., Rich, M. W., & Carney, R. M. (2021). Improving quality of life in heart failure. *Current Cardiology Reports, 23*(11), Article 159. <https://doi.org/10.1007/s11886-021-01588-y>
- Freedland, K. E., Rich, M. W., Skala, J. A., Carney, R. M., Davila-Roman, V. G., & Jaffe, A. S. (2003). Prevalence of depression in hospitalized patients with congestive heart failure. *Psychosomatic Medicine, 65*(1), 119–128.
<https://doi.org/10.1097/01.PSY.0000038938.67401.85>

- Freitas, D., Athanazio, R., Almeida, D., Dantas, N., & Reis, F. (2006). Sildenafil improves quality of life in men with heart failure and erectile dysfunction. *International Journal of Impotence Research*, 18(2), 210–212. <https://doi.org/10.1038/sj.ijir.3901385>
- Friel, C. J. (2016). Improving health outcomes for low health literacy heart failure patients. *Home Healthcare Now*, 34(8), 434–439. <https://doi.org/10.1097/NHH.0000000000000433>
- Fuqua, S. R., Wyatt, S. B., Andrew, M. E., Sarpong, D. F., Henderson, F. R., Cunningham, M. F., & Taylor, H. A., Jr. (2005). Recruiting African-American research participation in the Jackson Heart Study: Methods, response rates, and sample description. *Ethnicity & Disease*, 15(4_Suppl 6), S6–18.
- Gaffey, A. E., Cavanagh, C. E., Rosman, L., Wang, K., Deng, Y., Sims, M., O'Brien, E. C., Chamberlain, A. M., Mentz, R. J., Glover, L. M., & Burg, M. M. (2022). Depressive symptoms and incident heart failure in the Jackson Heart Study: Differential risk among Black men and women. *Journal of the American Heart Association*, 11(5), Article e022514. <https://doi.org/10.1161/JAHA.121.022514>
- Georgopoulou, S., Wright, A. J., Weinman, J., Booth, H., Thornton, H., & White, P. (2011). The relationship between socioeconomic status, quality of life and healthcare access in COPD: A systematic review. *European Respiratory Journal*, 38(Suppl 55), Article 684.
- Ghali, J. K., Kadakia, S., Cooper, R., & Ferlinz, J. (1988). Precipitating factors leading to decompensation of heart failure: Traits among urban Blacks. *Archives of Internal Medicine*, 148(9), 2013–2016. <https://doi.org/10.1001/archinte.1988.00380090087021>

- Ghasemi, E., Aliha, J. M., Bastani, F., Haghani, H., & Samiei, N. (2014). Quality of life in women with coronary artery disease. *Iranian Red Crescent Medical Journal*, 16(7), Article e10188. <https://doi.org/10.5812/ircmj.10188>
- Ghasemi, M., Arab, M., & Mangolian Shahrababaki, P. (2020). Relationship between caregiver burden and family functioning in family caregivers of older adults with heart failure. *Journal of Gerontological Nursing*, 46(6), 25–33. <https://doi.org/10.3928/00989134-20200511-04>
- Ghosh, R. K., Ball, S., Prasad, V., & Gupta, A. (2016). Depression in heart failure: Intricate relationship, pathophysiology and most updated evidence of interventions from recent clinical studies. *International Journal of Cardiology*, 224, 170–177. <https://doi.org/10.1016/j.ijcard.2016.09.063>
- Gibson, L. M., & Hendricks, C. S. (2006). Integrative review of spirituality in African American breast cancer survivors. *ABNF Journal*, 17(2), 67–72.
- Glynn, P., Lloyd-Jones, D. M., Feinstein, M. J., Carnethon, M., & Khan, S. S. (2019). Disparities in cardiovascular mortality related to heart failure in the United States. *Journal of the American College of Cardiology*, 73(18), 2354–2355. <https://doi.org/10.1016/j.jacc.2019.02.042>
- Go, A. S., Mozaffarian, D., Roger, V. L., Benjamin, E. J., Berry, J. D., Blaha, M. J., Dai, S., Ford, E. S., Fox, C. S., Franco, S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Huffman, M. D., Judd, S. E., Kissela, B. M., Kittner, S. J., ... Turner, M. B. (2013). Heart disease and stroke statistics—2014 update: A report from the American Heart Association. *Circulation*, 129(3). <https://doi.org/10.1161/01.cir.0000441139.02102.80>

Go et al., 2014

Gottlieb, S. S., Khatta, M., Friedmann, E., Einbinder, L., Katzen, S., Baker, B., Marshall, J., Minshall, S., Robinson, S., Fisher, M. L., Potenza, M., Sigler, B., Baldwin, C., & Thomas, S. A. (2004). The influence of age, gender, and race on the prevalence of depression in heart failure patients. *Journal of the American College of Cardiology*, 43(9), 1542–1549. <https://doi.org/10.1016/j.jacc.2003.10.064>

Graham, L. A., Ngwa, J., Ntekim, O., Ogunlana, O., Wolday, S., Johnson, S., Johnson, M., Castor, C., Fungwe, T. V., & Obisesan, T. O. (2018). Best strategies to recruit and enroll elderly Blacks into clinical and biomedical research. *Clinical Interventions in Aging*, 13, 43–50. <https://doi.org/10.2147/CIA.S130112>

Granger, B. B., Sandelowski, M., Tahshjain, H., Swedberg, K., & Ekman, I. (2009). A qualitative descriptive study of the work of adherence to a chronic heart failure regimen: patient and physician perspectives. *Journal of Cardiovascular Nursing*, 24(4), 308–315. <https://doi.org/10.1097/JCN.0b013e3181a4be30>

Graven, L. J., & Grant, J. (2013). The impact of social support on depressive symptoms in individuals with heart failure: Update and review. *The Journal of Cardiovascular Nursing*, 28(5), 429–443. <https://doi.org/10.1097/JCN.0b013e3182578b9d>

Greene, S. J., Butler, J., Spertus, J. A., Hellkamp, A. S., Vaduganathan, M., DeVore, A. D., Albert, N. M., Duffy, C. I., Patterson, J. H., Thomas, L., Williams, F. B., Hernandez, A. F., & Fonarow, G. C. (2021). Comparison of New York Heart Association class and patient-reported outcomes for heart failure with reduced ejection fraction. *JAMA Cardiology*, 6(5), 522–531. <https://doi.org/10.1001/jamacardio.2021.0372>

- Griffin, M. T. Q., Lee, Y.-H., Salman, A., Seo, Y., Marin, P. A., Starling, R. C., & Fitzpatrick, J. J. (2007). Spirituality and well being among elders: Differences between elders with heart failure and those without heart failure. *Clinical Interventions in Aging*, 2(4), 669–675. <https://doi.org/10.2147/cia.s874>
- Grigorovich, A., Lee, A., Ross, H., Woodend, A. K., Forde, S., & Cameron, J. I. (2017). A longitudinal view of factors that influence the emotional well-being of family caregivers to individuals with heart failure. *Aging & Mental Health*, 21(8), 844–850. <https://doi.org/10.1080/13607863.2016.1168361>
- Guzman-Clark, J. R. S., van Servellen, G., Chang, B., Mentes, J., & Hahn, T. J. (2013). Predictors and outcomes of early adherence to the use of a home telehealth device by older veterans with heart failure. *Telemedicine and e-Health*, 19(3), 217–223. <https://doi.org/10.1089/tmj.2012.0096>
- Habal, M. V., Axsom, K., & Farr, M. (2019). Advanced therapies for advanced heart failure in women. *Heart Failure Clinics*, 15(1), 97–107. <https://doi.org/10.1016/j.hfc.2018.08.010>
- Haesler, E., Bauer, M., & Fetherstonhaugh, D. (2016). Sexuality, sexual health and older people: A systematic review of research on the knowledge and attitudes of health professionals. *Nurse Education Today*, 40, 57–71. <https://doi.org/10.1016/j.nedt.2016.02.012>
- Hamatani, Y., Iguchi, M., Ikeyama, Y., Kunugida, A., Ogawa, M., Yasuda, N., Fujimoto, K., Ichihara, H., Sakai, M., & Kinoshita, T. (2021). Prevalence, temporal change, and determinants of anxiety and depression in hospitalized patients with heart failure. *Journal of Cardiac Failure*, 28(2), 181–190. <https://doi.org/10.1016/j.cardfail.2021.07.024>

- Hayes, T., & Sharma, M. (2021). Applying the integrated marketing communication approach to recruit and retain African American women. *Health Promotion Perspectives, 11*(4), 460–466. <https://doi.org/10.34172/hpp.2021.58>
- Heiat, A., Gross, C. P., & Krumholz, H. M. (2002). Representation of the elderly, women, and minorities in heart failure clinical trials. *Archives of Internal Medicine, 162*(15), 1682–1688. <https://doi.org/10.1001/archinte.162.15.1682>
- Heiden-Rootes, K. M., Salas, J., Gebauer, S., Witthaus, M., Scherrer, J., McDaniel, K., & Carver, D. (2017). Sexual dysfunction in primary care: An exploratory descriptive analysis of medical record diagnoses. *Journal of Sexual Medicine, 14*(11), 1318–1326. <https://doi.org/10.1016/j.jsxm.2017.09.014>
- Heidenreich, P. A., Albert, N. M., Allen, L. A., Bluemke, D. A., Butler, J., Fonarow, G. C., Ikonomidis, J. S., Khavjou, O., Konstam, M. A., Maddox, T. M., Nichol, G., Pham, M., Piña, I. L., & Trogon, J. G. (2013). Forecasting the impact of heart failure in the United States: A policy statement from the American Heart Association. *Circulation: Heart Failure, 6*(3), 606–629. <https://doi.org/10.1161/HHF.0b013e318291329a>
- Heidenreich, P. A., Fonarow, G. C., Opsha, Y., Sandhu, A. T., Sweitzer, N. K., Warraich, H. J., Butler, J., Hsich, E., Pressler, S. B., Shah, K., Taylor, K., Sabe, M., & Ng, T. (2022). Economic issues in heart failure in the United States. *Journal of Cardiac Failure, 28*(3), 453–466. <https://doi.org/10.1016/j.cardfail.2021.12.017>

- Heidenreich, P. A., Trogon, J. G., Khavjou, O. A., Butler, J., Dracup, K., Ezekowitz, M. D., Finkelstein, E. A., Hong, Y., Johnston, S. C., Khera, A., Lloyd-Jones, D. M., Nelson, S. A., Nichol, G., Orenstein, D., Wilson, P. W., & Woo, Y. J. (2011). Forecasting the future of cardiovascular disease in the United States: A policy statement from the American Heart Association. *Circulation*, *123*(8), 933–944.
<https://doi.org/10.1161/CIR.0b013e31820a55f5>
- Heo, S., Doering, L. V., Widener, J., & Moser, D. K. (2008). Predictors and effect of physical symptom status on health-related quality of life in patients with heart failure. *American Journal of Critical Care*, *17*(2), 124–132. <https://doi.org/10.4037/ajcc2008.17.2.124>
- Heo, S., Lennie, T. A., Okoli, C., & Moser, D. K. (2009). Quality of life in patients with heart failure: Ask the patients. *Heart & Lung*, *38*(2), 100–108.
<https://doi.org/10.1016/j.hrtlng.2008.04.002>
- Heo, S., Moser, D. K., Chung, M. L., & Lennie, T. A. (2012). Social status, health-related quality of life, and event-free survival in patients with heart failure. *European Journal of Cardiovascular Nursing*, *11*(2), 141–149. <https://doi.org/10.1016/j.ejcnurse.2010.10.003>
- Heo, S., Moser, D. K., Lennie, T. A., Zambroski, C. H., & Chung, M. L. (2007). A comparison of health-related quality of life between older adults with heart failure and healthy older adults. *Heart & Lung*, *36*(1), 16–24. <https://doi.org/10.1016/j.hrtlng.2006.06.003>
- Herr, J. K., Salyer, J., Flattery, M., Goodloe, L., Lyon, D. E., Kabban, C. S., & Clement, D. G. (2015). Heart failure symptom clusters and functional status: A cross-sectional study. *Journal of Advanced Nursing*, *71*(6), 1274–1287. <https://doi.org/10.1111/jan.12596>

- Herr, J. K., Salyer, J., Lyon, D. E., Goodloe, L., Schubert, C., & Clement, D. G. (2014). Heart failure symptom relationships: A systematic review. *Journal of Cardiovascular Nursing*, 29(5), 416–422. <https://doi.org/10.1097/JCN.0b013e31829b675e>
- Hertzog, M. A., Pozehl, B., & Duncan, K. (2010). Cluster analysis of symptom occurrence to identify subgroups of heart failure patients: A pilot study. *The Journal of Cardiovascular Nursing*, 25(4), 273–283. <https://doi.org/10.1097/JCN.0b013e3181cfbb6c>
- Hill-Joseph, E. A. (2019). Coping while Black: Chronic illness, mastery, and the Black-White health paradox. *Journal of Racial and Ethnic Health Disparities*, 6(5), 935–943. <https://doi.org/10.1007/s40615-019-00594-9>
- Hirsch, J. K., Duberstein, P. R., & Unützer, J. (2009). Chronic medical problems and distressful thoughts of suicide in primary care patients: Mitigating role of happiness. *International Journal of Geriatric Psychiatry*, 24(7), 671–679. <https://doi.org/10.1002/gps.2174>
- Hoekstra, T., Lesman-Leegte, I., Couperus, M. F., Sanderman, R., & Jaarsma, T. (2012). What keeps nurses from the sexual counseling of patients with heart failure? *Heart & Lung*, 41(5), 492–499. <https://doi.org/10.1016/j.hrtlng.2012.04.009>
- Holt, C. L., Schulz, E., & Wynn, T. A. (2009). Perceptions of the religion-health connection among African Americans in the Southeastern United States: Sex, age, and urban/rural differences. *Health Education & Behavior*, 36(1), 62–80. <https://doi.org/10.1177/1090198107303314>
- Holzapfel et al., 2008**
- Hopper, I., Kotecha, D., Chin, K. L., Mentz, R. J., & von Lueder, T. G. (2016). Comorbidities in heart failure: Are there gender differences? *Current Heart Failure Reports*, 13(1), 1–12. <https://doi.org/10.1007/s11897-016-0280-1>

Hsich, E. M., & Pina, I. L. (2009). Heart failure in women: A need for prospective data. *Journal of the American College of Cardiology*, 54(6), 491–498.

<https://doi.org/10.1016/j.jacc.2009.02.066>

Hu, X., Dolansky, M. A., Hu, X., Zhang, F., & Qu, M. (2016). Factors associated with the caregiver burden among family caregivers of patients with heart failure in Southwest China. *Nursing & Health Sciences*, 18(1), 105–112. <https://doi.org/10.1111/nhs.12253>

Huang, H.-H., & Coker, A. D. (2010). Examining issues affecting African American participation in research studies. *Journal of Black Studies*, 40(4), 619–636.

<https://doi.org/10.1177/0021934708317749>

Hughes, T. B., Varma, V. R., Pettigrew, C., & Albert, M. S. (2017). African Americans and clinical research: Evidence concerning barriers and facilitators to participation and recruitment recommendations. *The Gerontologist*, 57(2), 348–358.

<https://doi.org/10.1093/geront/gnv118>

Humphreys et al., 2008

Hung, C.-L., Chao, T.-F., Su, C.-H., Liao, J.-N., Sung, K.-T., Yeh, H.-I., & Chiang, C.-E. (2021). Income level and outcomes in patients with heart failure with universal health coverage.

Heart, 107(3), 208–216. <https://doi.org/10.1136/heartjnl-2020-316793>

Isler, M. R., & Corbie-Smith, G. (2012). Practical steps to community engaged research: From inputs to outcomes. *The Journal of Law, Medicine & Ethics*, 40(4), 904–914.

<https://doi.org/10.1111/j.1748-720X.2012.00719.x>

Jaarsma, T. (2017). Sexual function of patients with heart failure: Facts and numbers. *ESC Heart Failure*, 4(1), 3–7. <https://doi.org/10.1002/ehf2.12108>

- Jaarsma, T., Hill, L., Bayes-Genis, A., La Rocca, H.-P. B., Castiello, T., Celutkiene, J., Marques-Sule, E., Plymen, C. M., Piper, S. E., Riegel, B., Rutten, F. H., Ben Gal, T., Bauersachs, J., Coats, A. J. S., Chioncel, O., Lopatin, Y., Lund, L. H., Lainscak, M., Moura, B., ... Stromberg, A. (2021). Self-care of heart failure patients: Practical management recommendations from the Heart Failure Association of the European Society of Cardiology. *European Journal of Heart Failure*, 23(1), 157–174.
<https://doi.org/10.1002/ejhf.2008>
- Jackson-Triche, M. E., Greer Sullivan, J., Wells, K. B., Rogers, W., Camp, P., & Mazel, R. (2000). Depression and health-related quality of life in ethnic minorities seeking care in general medical settings. *Journal of Affective Disorders*, 58(2), 89–97.
[https://doi.org/10.1016/S0165-0327\(99\)00069-5](https://doi.org/10.1016/S0165-0327(99)00069-5)
- James, D. C., Harville, C., Sears, C., Efunbumi, O., & Bondoc, I. (2017). Participation of African Americans in e-Health and m-Health studies: A systematic review. *Telemedicine and e-Health*, 23(5), 351–364. <https://doi.org/10.1089/tmj.2016.0067>
- Javed, Z., Haisum Maqsood, M., Yahya, T., Amin, Z., Acquah, I., Valero-Elizondo, J., Andrieni, J., Dubey, P., Jackson, R. K., Daffin, M. A., Cainzos-Achirica, M., Hyder, A. A., & Nasir, K. (2022). Race, racism, and cardiovascular health: Applying a social determinants of health framework to racial/ethnic disparities in cardiovascular disease. *Circulation Cardiovascular Quality and Outcomes*, 15(1), Article e007917.
<https://doi.org/10.1161/CIRCOUTCOMES.121.007917>

- Jiang, W., Alexander, J., Christopher, E., Kuchibhatla, M., Gaulden, L. H., Cuffe, M. S., Blazing, M. A., Davenport, C., Califf, R. M., Krishnan, R. R., & O'Connor, C. M. (2001). Relationship of depression to increased risk of mortality and rehospitalization in patients with congestive heart failure. *Archives of Internal Medicine*, *161*(15), 1849–1856. <https://doi.org/10.1001/archinte.161.15.1849>
- Jonsson, U., Alexanderson, K., Kjeldgård, L., Westerlund, H., & Mittendorfer-Rutz, E. (2013). Diagnosis-specific disability pension predicts suicidal behaviour and mortality in young adults: A nationwide prospective cohort study. *BMJ Open*, *3*(2), Article e002286. <https://doi.org/10.1136/bmjopen-2012-002286>
- Jurgens, C. Y., Moser, D. K., Armola, R., Carlson, B., Sethares, K., & Riegel, B. (2009). Symptom clusters of heart failure. *Research in Nursing & Health*, *32*(5), 551–560. <https://doi.org/10.1002/nur.20343>
- Juurink, D. N., Herrmann, N., Szalai, J. P., Kopp, A., & Redelmeier, D. A. (2004). Medical illness and the risk of suicide in the elderly. *Archives of Internal Medicine*, *164*(11), 1179–1184. <https://doi.org/10.1001/archinte.164.11.1179>
- Keyvanara, M. M., Khasti, B. Y. B. Y., Zadeh, M. R. M. R., & Modaber, F. F. (2015). Study of the relationship between quality of life and socioeconomic status in Isfahan at 2011. *Journal of Education and Health Promotion*, *4*, Article 92. <https://doi.org/10.4103/2277-9531.171806>
- Koenig, H. G. (1998). Depression in hospitalized older patients with congestive heart failure. *General Hospital Psychiatry*, *20*(1), 29–43. [https://doi.org/10.1016/S0163-8343\(98\)80001-7](https://doi.org/10.1016/S0163-8343(98)80001-7)

- Koenig, H. G. (2004). Spirituality, wellness, and quality of life. *Sexuality, Reproduction and Menopause*, 2(2), 76–82. <https://doi.org/10.1016/j.sram.2004.04.004>
- Kolbe, N., Kugler, C., Schnepf, W., & Jaarsma, T. (2016). Sexual counseling in patients with heart failure: A silent phenomenon: Results from a convergent parallel mixed method study. *The Journal of Cardiovascular Nursing*, 31(1), 53–61. <https://doi.org/10.1097/JCN.0000000000000215>
- Konstam, V., Moser, D. K., & De Jong, M. J. (2005). Depression and anxiety in heart failure. *Journal of Cardiac Failure*, 11(6), 455–463. <https://doi.org/10.1016/j.cardfail.2005.03.006>
- Korkmaz, H., Korkmaz, S., & Çakar, M. (2019). Suicide risk in chronic heart failure patients and its association with depression, hopelessness and self esteem. *Journal of Clinical Neuroscience*, 68, 51–54. <https://doi.org/10.1016/j.jocn.2019.07.062>
- Kye, S. Y., & Park, K. (2016). Suicidal ideation and suicidal attempts among adults with chronic diseases: A cross-sectional study. *Comprehensive Psychiatry*, 73, 160–167. <https://doi.org/10.1016/j.comppsy.2016.12.001>
- Lang, R., Kelkar, V. A., Byrd, J. R., Edwards, C. L., Pericak-Vance, M., & Byrd, G. S. (2013). African American participation in health-related research studies: Indicators for effective recruitment. *Journal of Public Health Management and Practice*, 19(2), 110–118. <https://doi.org/10.1097/PHH.0b013e31825717ef>
- Latulippe, K., Hamel, C., & Giroux, D. (2017). Social health inequalities and ehealth: A literature review with qualitative synthesis of theoretical and empirical studies. *Journal of Medical Internet Research*, 19(4), Article e136. <https://doi.org/10.2196/jmir.6731>

- Le Strat, Y., Le Foll, B., & Dubertret, C. (2015). Major depression and suicide attempts in patients with liver disease in the United States. *Liver International*, 35(7), 1910–1916.
<https://doi.org/10.1111/liv.12612>
- Lee, C. S., Moser, D. K., Lennie, T. A., & Riegel, B. (2011). Event-free survival in adults with heart failure who engage in self-care management. *Heart & Lung*, 40(1), 12–20.
- Lee, D. T. F., Yu, D. S. F., Woo, J., & Thompson, D. R. (2005). Health-related quality of life in patients with congestive heart failure. *European Journal of Heart Failure*, 7(3), 419–422.
<https://doi.org/10.1016/j.ejheart.2004.08.004>
- Lee, S., & Riegel, B. (2018). State of the science in heart failure symptom perception research: An integrative review. *Journal of Cardiovascular Nursing*, 33(3), 204–210.
<https://doi.org/10.1097/JCN.0000000000000445>
- Lesman-Leegte, I., Jaarsma, T., Coyne, J. C., Hillege, H. L., Van Veldhuisen, D. J., & Sanderman, R. (2009). Quality of life and depressive symptoms in the elderly: A comparison between patients with heart failure and age- and gender-matched community controls. *Journal of Cardiac Failure*, 15(1), 17–23.
<https://doi.org/10.1016/j.cardfail.2008.09.006>
- Lesyuk, W., Kriza, C., & Kolominsky-Rabas, P. (2018). Cost-of-illness studies in heart failure: A systematic review 2004–2016. *BMC Cardiovascular Disorders*, 18(1), Article 74.
<https://doi.org/10.1186/s12872-018-0815-3>

- Lewis, T. T., Guo, H., Lunos, S., Mendes de Leon, C. F., Skarupski, K. A., Evans, D. A., & Everson-Rose, S. A. (2011). Depressive symptoms and cardiovascular mortality in older African-American and White adults: Evidence for a differential association by race. *Circulation Cardiovascular Quality and Outcomes*, *4*(3), 293–299. <https://doi.org/10.1161/CIRCOUTCOMES.110.957548>
- Lindenauer, P. K., Lagu, T., Rothberg, M. B., Avrunin, J., Pekow, P. S., Wang, Y., & Krumholz, H. M. (2013). Income inequality and 30 day outcomes after acute myocardial infarction, heart failure, and pneumonia: Retrospective cohort study. *BMJ*, *346*, Article f521. <https://doi.org/10.1136/bmj.f521>
- Liu, C.-H., Wang, J.-H., Weng, S.-C., Cheng, Y.-H., Yeh, M.-K., Bai, M.-Y., & Chang, J.-C. (2018). Is heart failure associated with risk of suicide? *Journal of Cardiac Failure*, *24*(11), 795–800. <https://doi.org/10.1016/j.cardfail.2018.07.006>
- Liu, M.-H., Chiou, A.-F., Wang, C.-H., Yu, W.-P., & Lin, M.-H. (2021). Relationship of symptom stress, care needs, social support, and meaning in life to quality of life in patients with heart failure from the acute to chronic stages: A longitudinal study. *Health Quality of Life Outcomes*, *19*(1), Article 252. <https://doi.org/10.1186/s12955-021-01885-8>
- Liu, X. B., Ayatollahi, Y., Yamashita, T., Jaradat, M., Shen, J. J., Kim, S. J., Lee, Y.-J., Hwang, J., Yeom, H., & Upadhyay, S. (2019). Health literacy and mortality in patients with heart failure: A systematic review and meta-analysis. *Research in Gerontological Nursing*, *12*(2), 99–108. <https://doi.org/10.3928/19404921-20181018-01>

- Lockhart, E., Foreman, J., Mase, R., & Heisler, M. (2014). Heart failure patients' experiences of a self-management peer support program: A qualitative study. *Heart and Lung: Journal of Acute and Critical Care*, 43(4), 292–298. <https://doi.org/10.1016/j.hrtlng.2014.04.008>
- Loeb, S. J. (2006). African American older adults coping with chronic health conditions. *Journal of Transcultural Nursing*, 17(2), 139–147. <https://doi.org/10.1177/1043659605285415>
- Lossnitzer, N., Müller-Tasch, T., Löwe, B., Zugck, C., Nelles, M., Remppis, A., Haass, M., Rauch, B., Jünger, J., Herzog, W., & Wild, B. (2009). Exploring potential associations of suicidal ideation and ideas of self-harm in patients with congestive heart failure. *Depression & Anxiety*, 26(8), 764–768. <https://doi.org/10.1002/da.20587>
- Luebbert, R., & Perez, A. (2016). Barriers to clinical research participation among African Americans. *Journal of Transcultural Nursing*, 27(5), 456–463. <https://doi.org/10.1177/1043659615575578>
- Lum, H. D., Lo, D., Hooker, S., & Bekelman, D. B. (2014). Caregiving in heart failure: Relationship quality is associated with caregiver benefit finding and caregiver burden. *Heart & Lung*, 43(4), 306–310. <https://doi.org/10.1016/j.hrtlng.2014.05.002>
- Luttik, M. L., Jaarsma, T., Veeger, N., & van Veldhuisen, D. J. (2006). Marital status, quality of life, and clinical outcome in patients with heart failure. *Heart & Lung*, 35(1), 3–8. <https://doi.org/10.1016/j.hrtlng.2005.08.001>
- Macabasco-O'Connell, A., Crawford, M. H., Stotts, N., Stewart, A., & Froelicher, E. S. (2008). Self-care behaviors in indigent patients with heart failure. *The Journal of Cardiovascular Nursing*, 23(3), 223–230. <https://doi.org/10.1097/01.JCN.0000317427.21716.5f>

- Macabasco-O'Connell, A., DeWalt, D. A., Broucksou, K. A., Hawk, V., Baker, D. W., Schillinger, D., Ruo, B., Bibbins-Domingo, K., Holmes, G. M., & Erman, B. (2011). Relationship between literacy, knowledge, self-care behaviors, and heart failure-related quality of life among patients with heart failure. *Journal of General Internal Medicine*, 26(9), 979–986. <https://doi.org/10.1007/s11606-011-1668-y>
- McCabe, M. P., Sharlip, I. D., Lewis, R., Atalla, E., Balon, R., Fisher, A. D., Laumann, E., Lee, S. W., & Segraves, R. T. (2016). Risk factors for sexual dysfunction among women and men: A consensus statement from the Fourth International Consultation on Sexual Medicine 2015. *Journal of Sexual Medicine*, 13(2), 153–167. <https://doi.org/10.1016/j.jsxm.2015.12.015>
- McCarthy, M., Katz, S. D., Schipper, J., & Dickson, V. V. (2015). “I just can’t do it anymore”: Patterns of African Americans with heart failure: A mixed method study. *Healthcare*, 3(4), 973–986. <https://doi.org/10.3390/healthcare3040973>
- McGrath, M., Low, M. A., Power, E., McCluskey, A., & Lever, S. (2021). Addressing sexuality among people living with chronic disease and disability: A systematic mixed methods review of knowledge, attitudes, and practices of health care professionals. *Archives of Physical Medicine and Rehabilitation*, 102(5), 999–1010. <https://doi.org/10.1016/j.apmr.2020.09.379>
- McNaughton, C. D., Cawthon, C., Kripalani, S., Liu, D., Storrow, A. B., & Roumie, C. L. (2015). Health literacy and mortality: A cohort study of patients hospitalized for acute heart failure. *Journal of the American Heart Association*, 4(5), Article e001799. <https://doi.org/10.1161/JAHA.115.001799>

- Mead, H., Andres, E., Ramos, C., Siegel, B., & Regenstein, M. (2010). Barriers to effective self-management in cardiac patients: The patient's experience. *Patient Education and Counseling*, 79(1), 69–76. <https://doi.org/10.1016/j.pec.2009.08.003>
- Melo, C. D. F., Sampaio, I. S., Souza, D. L. D. A., & Pinto, N. D. S. (2015). Correlation between religiousness, spirituality and quality of life: A review of literature. *Estudos e Pesquisas em Psicologia*, 15(2), 447–464.
- Mensah, G. A., & Dunbar, S. B. (2006). A framework for addressing disparities in cardiovascular health. *The Journal of Cardiovascular Nursing*, 21(6), 451–456. <https://doi.org/10.1097/00005082-200611000-00007>
- Mensah, G. A., Mokdad, A. H., Ford, E. S., Greenlund, K. J., & Croft, J. B. (2005). State of disparities in cardiovascular health in the United States. *Circulation*, 111(10), 1233–1241. <https://doi.org/10.1161/01.CIR.0000158136.76824.04>
- Mensah, G. A., Roth, G. A., & Fuster, V. (2019). *The global burden of cardiovascular diseases and risk factors: 2020 and beyond*. American College of Cardiology Foundation.
- Mensah, G. A., Stoney, C. M., Freemer, M. M., Smith, S., Engelgau, M. M., Hoots, W. K., Kiley, J. P., & Goff, D. C. (2019). The National Heart, Lung, and Blood Institute strategic vision implementation for health equity research. *Ethnicity & Disease*, 29(Suppl 1), 57–64. <https://doi.org/10.18865/ed.29.S1.57>
- Mentz, R. J., Babyak, M. A., Bittner, V., Fleg, J. L., Keteyian, S. J., Swank, A. M., Pina, I. L., Kraus, W. E., Whellan, D. J., O'Connor, C. M., & Blumenthal, J. A. (2015). Prognostic significance of depression in Blacks with heart failure: Insights from Heart Failure: A Controlled Trial Investigating Outcomes of Exercise Training. *Circulation: Heart Failure*, 8(3), 497–503. <https://doi.org/10.1161/CIRCHEARTFAILURE.114.001995>

- Millon Underwood, S., Buseh, A. G., Kelber, S. T., Stevens, P. E., & Townsend, L. (2013). Enhancing the participation of African Americans in health-related genetic research: Findings of a collaborative academic and community-based research study. *Nursing Research and Practice*, 2013, Article 749563. <https://doi.org/10.1155/2013/749563>
- Mills, P. J., Wilson, K., Iqbal, N., Iqbal, F., Alvarez, M., Pung, M. A., Wachmann, K., Rutledge, T., Maglione, J., Zisook, S., Dimsdale, J. E., Lunde, O., Greenberg, B. H., Maisel, A., Raisinghani, A., Natarajan, L., Jain, S., Hufford, D. J., & Redwine, L. (2015). Depressive symptoms and spiritual wellbeing in asymptomatic heart failure patients. *Journal of Behavioral Medicine*, 38(3), 407–415. <https://doi.org/10.1007/s10865-014-9615-0>
- Mitchell, J. E., Ferdinand, K. C., Watson, K. E., Wenger, N. K., Watkins, L. O., Flack, J. M., Gavin, J. R., III, Reed, J. W., Saunders, E., & Wright, J. T., Jr. (2011). Treatment of heart failure in African Americans—A call to action. *Journal of the National Medical Association*, 103(2), 86–98. [https://doi.org/10.1016/S0027-9684\(15\)30257-1](https://doi.org/10.1016/S0027-9684(15)30257-1)
- Mody, L., Miller, D. K., McGloin, J. M., Freeman, M., Marcantonio, E. R., Magaziner, J., & Studenski, S. (2008). Recruitment and retention of older adults in aging research. *Journal of the American Geriatrics Society*, 56(12), 2340–2348. <https://doi.org/10.1111/j.1532-5415.2008.02015.x>
- Mohr, D. C., Schueller, S. M., Montague, E., Burns, M. N., & Rashidi, P. (2014). The behavioral intervention technology model: An integrated conceptual and technological framework for eHealth and mHealth interventions. *Journal of Medical Internet Research*, 16(6), Article e146. <https://doi.org/10.2196/jmir.3077>

- Molloy, G. J., O'Carroll, R. E., Witham, M. D., & McMurdo, M. E. T. (2012). Interventions to enhance adherence to medications in patients with heart failure: A systematic review. *Circulation: Heart Failure*, 5(1), 126–133.
<https://doi.org/10.1161/CIRCHEARTFAILURE.111.964569>
- Montejo, A. L., Montejo, L., & Baldwin, D. S. (2018). The impact of severe mental disorders and psychotropic medications on sexual health and its implications for clinical management. *World Psychiatry*, 17(1), 3–11. <https://doi.org/10.1002/wps.20509>
- Moradi, M., Doostkami, M., Behnamfar, N., Rafiemanesh, H., & Behzadmehr, R. (2021). Global prevalence of depression among heart failure patients: A systematic review and meta-analysis. *Current Problems in Cardiology*, Article 100848.
<https://doi.org/10.1016/j.cpcardiol.2021.100848>
- Morrow, D., Clark, D., Tu, W., Wu, J., Weiner, M., Steinley, D., & Murray, M. D. (2006). Correlates of health literacy in patients with chronic heart failure. *The Gerontologist*, 46(5), 669–676. <https://doi.org/10.1093/geront/46.5.669>
- Moser, D. K., Lee, K. S., Wu, J.-R., Mudd-Martin, G., Jaarsma, T., Huang, T.-Y., Fan, X.-Z., Strömberg, A., Lennie, T. A., & Riegel, B. (2014). Identification of symptom clusters among patients with heart failure: An international observational study. *International Journal of Nursing Studies*, 51(10), 1366–1372.
<https://doi.org/10.1016/j.ijnurstu.2014.02.004>
- Moser, D. K., Robinson, S., Biddle, M. J., Pelter, M. M., Nesbitt, T. S., Southard, J., Cooper, L., & Dracup, K. (2015). Health literacy predicts morbidity and mortality in rural patients with heart failure. *Journal of Cardiac Failure*, 21(8), 612–618.
<https://doi.org/10.1016/j.cardfail.2015.04.004>

Mozaffarian, D., Benjamin, E. J., Go, A. S., Arnett, D. K., Blaha, M. J., Cushman, M., Das, S. R., de Ferranti, S., Després, J.-P., Fullerton, H. J., Howard, V. J., Huffman, M. D., Isasi, C. R., Jiménez, M. C., Judd, S. E., Kissela, B. M., Lichtman, J. H., Lisabeth, L. D., Liu, S., ... Turner, M. B. (2016). Heart disease and stroke statistics—2016 update: A report from the American Heart Association. *Circulation*, *133*(4), e38–e360.

<https://doi.org/10.1161/CIR.0000000000000350>

Mozaffarian, D., Benjamin, E. J., Go, A. S., Arnett, D. K., Blaha, M. J., Cushman, M., de Ferranti, S., Després, J.-P., Fullerton, H. J., Howard, V. J., Huffman, M. D., Judd, S. E., Kissela, B. M., Lackland, D. T., Lichtman, J. H., Lisabeth, L. D., Liu, S., Mackey, R. H., Matchar, D. B., ... Turner, M. B. (2015). Heart disease and stroke statistics—2015 update: A report from the American Heart Association. *Circulation*, *131*(4), e29–e322.

<https://doi.org/10.1161/CIR.0000000000000152>

Murray, M., Tu, W., Wu, J., Morrow, D., Smith, F., & Brater, D. (2009). Factors associated with exacerbation of heart failure include treatment adherence and health literacy skills. *Clinical Pharmacology & Therapeutics*, *85*(6), 651–658.

Naghi, J. J., Philip, K. J., Phan, A., Cleenewerck, L., & Schwarz, E. R. (2012). The effects of spirituality and religion on outcomes in patients with chronic heart failure. *Journal of Religion and Health*, *51*(4), 1124–1136. <https://doi.org/10.1007/s10943-010-9419-7>

Nair, N., Farmer, C., Gongora, E., & Dehmer, G. J. (2012). Commonality between depression and heart failure. *American Journal of Cardiology*, *109*(5), 768–772.

<https://doi.org/10.1016/j.amjcard.2011.10.039>

- Nardi, D., Waite, R., Nowak, M., Hatcher, B., Hines-Martin, V., & Stacciarini, J. M. R. (2020). Achieving health equity through eradicating structural racism in the United States: A call to action for nursing leadership. *Journal of Nursing Scholarship*, 52(6), 696–704.
<https://doi.org/10.1111/jnu.12602>
- Nayak, A., Hicks, A. J., & Morris, A. A. (2020). Understanding the complexity of heart failure risk and treatment in Black patients. *Circulation: Heart Failure*, 13(8), Article e007264.
<https://doi.org/10.1161/CIRCHEARTFAILURE.120.007264>
- National Institute for Nursing Research. (2022). *NINR 2022–2026 Strategic Plan*. National Institute for Nursing Research.
https://www.ninr.nih.gov/sites/files/docs/Zenk_NINR_2022-2026_Strategic_Plan_508c.pdf
- Nooruddin, M., Scherr, C., Friedman, P., Subrahmanyam, R., Banagan, J., Moreno, D., Sathyanarayanan, M., Nutescu, E., Jeyaram, T., Harris, M., Zhang, H., Rodriguez, A., Shaazuddin, M., Perera, M., & Tuck, M. (2020). Why African Americans say “no”: A study of pharmacogenomic research participation. *Ethnicity & Disease*, 30(Suppl 1), 159–166. <https://doi.org/10.18865/ed.30.S1.159>
- Oh, S. S., Galanter, J., Thakur, N., Pino-Yanes, M., Barcelo, N. E., White, M. J., de Bruin, D. M., Greenblatt, R. M., Bibbins-Domingo, K., Wu, A. H. B., Borrell, L. N., Gunter, C., Powe, N. R., & Burchard, E. G. (2015). Diversity in clinical and biomedical research: A promise yet to be fulfilled. *PLoS Medicine*, 12(12), Article e1001918.
<https://doi.org/10.1101/034538>

- Ohlsson, A., Eckerdal, N., Lindahl, B., Hanning, M., & Westerling, R. (2021). Non-employment and low educational level as risk factors for inequitable treatment and mortality in heart failure: A population-based cohort study of register data. *BMC Public Health*, 21(1), Article 1040. <https://doi.org/10.1186/s12889-021-10919-1>
- Ojeda, & McGuire, T. G. (2006). Gender and racial/ethnic differences in use of outpatient mental health and substance use services by depressed adults. *Psychiatric Quarterly*, 77(3), 211–222. <https://doi.org/10.1007/s11126-006-9008-9>
- Ornelas, C., Fadadu, R. P., Richardson, M. A., Agboghidi, O. H., & Davis, J. D. (2021). Bridge the gap: Reducing inequity in hospital readmissions for African American patients with heart failure through quality improvement initiatives. *Health Equity*, 5(1), 3–34. <https://doi.org/10.1089/heq.2020.0082>
- Pahuja, M., Leifer, E., Clarke, J. R., & Pina, I. L. (2022). The impact of race and ethnicity on clinical outcomes in patients with heart failure (on behalf of the Guide-IT Steering Committee). *Journal of Cardiac Failure*, 28(5), Article S2. <https://doi.org/10.1016/j.cardfail.2022.03.007>
- Panzini, R. G., Mosqueiro, B. P., Zimpel, R. R., Bandeira, D. R., Rocha, N. S., & Fleck, M. P. (2017). Quality-of-life and spirituality. *International Review of Psychiatry*, 29(3), 263–282. <https://doi.org/10.1080/09540261.2017.1285553>
- Park, J., Moser, D. K., Griffith, K., Harring, J. R., & Johantgen, M. (2019). Exploring symptom clusters in people with heart failure. *Clinical Nursing and Research*, 28(2), 165–181. <https://doi.org/10.1177/1054773817729606>

- Patel, R. S., Shrestha, S., Saeed, H., Raveendranathan, S., Isidahome, E. E., Ravat, V., Fakorede, M. O., & Patel, V. (2018). Comorbidities and consequences in hospitalized heart failure patients with depression. *Cureus, 10*(8), Article e3193.
<https://doi.org/10.7759/cureus.3193>
- Peeters, J., Wieggers, T. A., & Friele, R. D. (2013). How technology in care at home affects patient self-care and self-management: A scoping review. *International Journal of Environmental Research and Public Health, 10*(11), 5541-5564.
<https://doi.org/10.3390/ijerph10115541>
- Petersen, B. D., Stenager, E., Mogensen, C. B., & Erlangsen, A. (2020). The association between heart diseases and suicide: A nationwide cohort study. *Journal of Internal Medicine, 287*(5), 558-568. <https://doi.org/10.1111/joim.13025>
- Petersen, J. W., & Felker, G. M. (2006). Inflammatory biomarkers in heart failure. *Congestive Heart Failure, 12*(6), 324-328. <https://doi.org/10.1111/j.1527-5299.2006.05595.x>
- Peterson, P. N., Shetterly, S. M., Clarke, C. L., Bekelman, D. B., Chan, P. S., Allen, L. A., Matlock, D. D., Magid, D. J., & Masoudi, F. A. (2011). Health literacy and outcomes among patients with heart failure. *JAMA, 305*(16), 1695–1701.
<https://doi.org/10.1001/jama.2011.512>
- Phillips, L., Harrison, T., & Houck, P. (2005). Return to work and the person with heart failure. *Heart & Lung, 34*(2), 79–88. <https://doi.org/10.1016/j.hrtlng.2004.06.001>
- Piamjariyakul, U., Smith, C. E., Wick, J., & Fitzgerald, S. (2013). Nurse-led group clinic intervention reduced depression among African American patients with heart failure. *Clinical and Translational Science, 6*(2), 128–132.
<https://doi.org/10.1016/j.hrtlng.2018.02.005>

- Piamjariyakul, U., Thompson, N. C., Russell, C., & Smith, C. E. (2018). The effect of nurse-led group discussions by race on depressive symptoms in patients with heart failure. *Heart & Lung, 47*(3), 211–215. <https://doi.org/10.1016/j.hrtlng.2018.02.005>
- Polikandrioti, M., Panoutsopoulos, G., Tsami, A., Gerogianni, G., Saroglou, S., Thomai, E., & Leventzonis, I. (2019). Assessment of quality of life and anxiety in heart failure outpatients. *Archives of Medical Sciences: Atherosclerotic Diseases, 4*, Article e38. <https://doi.org/10.5114/amsad.2019.84444>
- Pressler, S. J. (2008). Cognitive functioning and chronic heart failure: A review of the literature (2002–July 2007). *The Journal of Cardiovascular Nursing, 23*(3), 239–249. <https://doi.org/10.1097/01.JCN.0000305096.09710.ec>
- Pugh, E., Stewart, J., Carter, L., Calamia, M., Carmichael, O., & Newton, J. R. L. (2022). Beliefs, understanding, and barriers related to dementia research participation among older African Americans. *Alzheimer Disease and Associated Disorders, 36*(1), 52–57. <https://doi.org/10.1097/WAD.0000000000000476>
- Reddy, Y. N. V., Rikhi, A., Obokata, M., Shah, S. J., Lewis, G. D., AbouEzzedine, O. F., Dunlay, S., McNulty, S., Chakraborty, H., Stevenson, L. W., Redfield, M. M., & Borlaug, B. A. (2020). Quality of life in heart failure with preserved ejection fraction: Importance of obesity, functional capacity, and physical inactivity. *European Journal of Heart Failure, 22*(6), 1009–1018. <https://doi.org/10.1002/ejhf.1788>
- Regan, J. A., Kitzman, D. W., Leifer, E. S., Kraus, W. E., Fleg, J. L., Forman, D. E., Whellan, D. J., Wojdyla, D., Parikh, K., O'Connor, C. M., & Mentz, R. J. (2019). Impact of age on comorbidities and outcomes in heart failure with reduced ejection fraction. *JACC: Heart Failure, 7*(12), 1056–1065. <https://doi.org/10.1016/j.jchf.2019.09.004>

Retrum, J. H., Boggs, J., Hersh, A., Wright, L., Main, D. S., Magid, D. J., & Allen, L. A. (2012).

Patient-identified factors related to heart failure readmissions. *Circulation:*

Cardiovascular Quality and Outcomes, 5(3), Article A124.

https://doi.org/10.1161/circoutcomes.5.suppl_1.A124

Reza, N., Gruen, J., & Bozkurt, B. (2022). Representation of women in heart failure clinical

trials: Barriers to enrollment and strategies to close the gap. *American Heart Journal*

Plus: Cardiology Research and Practice, 13, Article 100093.

<https://doi.org/10.1016/j.ahjo.2022.100093>

Riedinger, M. S., Dracup, K. A., Brecht, M.-L., Padilla, G., Sarna, L., & Ganz, P. A. (2001).

Quality of life in patients with heart failure: Do gender differences exist? *Heart & Lung*,

30(2), 105–116. <https://doi.org/10.1067/mhl.2001.114140>

Riegel & Dickson, 2008

Riegel, B., Dickson, V. V., & Faulkner, K. M. (2016). The situation-specific theory of heart

failure self-care: Revised and updated. *The Journal of Cardiovascular Nursing*, 31(3),

226–235. <https://doi.org/10.1097/JCN.0000000000000244>

Riegel, B., Lee, C. S., & Dickson, V. V. (2011). Self care in patients with chronic heart failure.

Nature Reviews Cardiology, 8(11), 644–654. <https://doi.org/10.1038/nrcardio.2011.95>

Riegel, B., Moser, D. K., Anker, S. D., Appel, L. J., Dunbar, S. B., Grady, K. L., Gurvitz, M. Z.,

Havranek, E. P., Lee, C. S., & Lindenfeld, J. (2009). State of the science promoting self-

care in persons with heart failure: A scientific statement from the American Heart

Association. *Circulation*, 120(12), 1141–1163.

<https://doi.org/10.1161/CIRCULATIONAHA.109.192628>

- Rockwell, J. M., & Riegel, B. (2001). Predictors of self-care in persons with heart failure. *Heart & Lung, 30*(1), 18–25. <https://doi.org/10.1067/mhl.2001.112503>
- Roger, K. S., & Hatala, A. (2018). Religion, spirituality & chronic illness: A scoping review and implications for health care practitioners. *Journal of Religion & Spirituality in Social Work, 37*(1), 24–44. <https://doi.org/10.1080/15426432.2017.1386151>
- Roger, V. L., Go, A. S., Lloyd-Jones, D. M., Benjamin, E. J., Berry, J. D., Borden, W. B., Bravata, D. M., Dai, S., Ford, E. S., Fox, C. S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Kissela, B. M., Kittner, S. J., Lackland, D. T., Lichtman, J. H., ... Turner, M. B. (2012). Heart disease and stroke statistics—2012 update: A report from the American Heart Association. *Circulation, 125*(1), e2–e220. <https://doi.org/10.1161/CIR.0b013e31823ac046>
- Rohrbaugh, M. J., Shoham, V., & Coyne, J. C. (2006). Effect of marital quality on eight-year survival of patients with heart failure. *American Journal of Cardiology, 98*(8), 1069–1072. <https://doi.org/10.1016/j.amjcard.2006.05.034>
- Rohyans, L. M., & Pressler, S. J. (2009). Depressive symptoms and heart failure: Examining the sociodemographic variables. *Clinical Nurse Specialist, 23*(3), 138–144. <https://doi.org/10.1097/NUR.0b013e3181a443b4>
- Rollins, L., Sy, A., Crowell, N., Rivers, D., Miller, A., Cooper, P., Teague, D., Jackson, C., Henry Akintobi, T., & Ofili, E. (2018). Learning and action in community health: Using the health belief model to assess and educate African American community residents about participation in clinical research. *International Journal of Environmental Research and Public Health, 15*(9), Article 1862. <https://doi.org/10.3390/ijerph15091862>

- Rørth, R., Fosbøl, E. L., Mogensen, U. M., Kragholm, K., Numé, A. K., Gislason, G. H., Jhund, P. S., Petrie, M. C., McMurray, J. J. V., Torp-Pedersen, C., Køber, L., & Kristensen, S. L. (2018). Employment status at time of first hospitalization for heart failure is associated with a higher risk of death and rehospitalization for heart failure. *European Journal of Heart Failure*, 20(2), 240–247. <https://doi.org/10.1002/ejhf.1046>
- Rørth, R., Wong, C., Kragholm, K., Fosbøl, E. L., Mogensen, U. M., Lamberts, M., Petrie, M. C., Jhund, P. S., Gerds, T. A., Torp-Pedersen, C., Gislason, G. H., McMurray, J. J. V., Køber, L., & Kristensen, S. L. (2016). Return to the workforce after first hospitalization for heart failure: A Danish nationwide cohort study. *Circulation*, 134(14), 999–1009. <https://doi.org/10.1161/CIRCULATIONAHA.116.021859>
- Rucker-Whitaker, C., Flynn, K. J., Kravitz, G., Eaton, C., Calvin, J. E., & Powell, L. H. (2006). Understanding African-American participation in a behavioral intervention: Results from focus groups. *Contemporary Clinical Trials*, 27(3), 274–286. <https://doi.org/10.1016/j.cct.2005.11.006>
- Rustad, J. K., Stern, T. A., Hebert, K. A., & Musselman, D. L. (2013). Diagnosis and treatment of depression in patients with congestive heart failure: A review of the literature. *The Primary Care Companion for CNS Disorders*, 15(4), Article 26254. <https://doi.org/10.4088/PCC.13r01511>
- Salyer, J., Flattery, M., & Lyon, D. E. (2019). Heart failure symptom clusters and quality of life. *Heart & Lung*, 48(5), 366–372. <https://doi.org/10.1016/j.hrtlng.2019.05.016>

- Sampaio, M. S., Vieira, W. D. A., Bernardino, Í. D. M., Herval, Á. M., Flores-Mir, C., & Paranhos, L. R. (2019). Chronic obstructive pulmonary disease as a risk factor for suicide: A systematic review and meta-analysis. *Respiratory Medicine*, *151*, 11–18.
<https://doi.org/10.1016/j.rmed.2019.03.018>
- Saunders, M. M. (2008). Factors associated with caregiver burden in heart failure family caregivers. *Western Journal of Nursing Research*, *30*(8), 943–959.
<https://doi.org/10.1177/0193945908319990>
- Sawatzky, R., Ratner, P. A., & Chiu, L. (2005). A meta-analysis of the relationship between spirituality and quality of life. *Social Indicators Research*, *72*(2), 153–188.
<https://doi.org/10.1007/s11205-004-5577-x>
- Scharff, D. P., Mathews, K. J., Jackson, P., Hoffsuemmer, J., Martin, E., & Edwards, D. (2010). More than Tuskegee: Understanding mistrust about research participation. *Journal of Health Care for the Poor and Underserved*, *21*(3), 879–897.
<https://doi.org/10.1353/hpu.0.0323>
- Schwarz, E. R., Kapur, V., Bionat, S., Rastogi, S., Gupta, R., & Rosanio, S. (2008). The prevalence and clinical relevance of sexual dysfunction in women and men with chronic heart failure. *International Journal of Impotence Research*, *20*(1), 85–91.
<https://doi.org/10.1038/sj.ijir.3901613>
- Senol, V. (2018). The relationship between sexual dysfunction and health-related quality of life in chronic renal failure patients. *Studies on Ethno-Medicine*, *12*(1), 40–48.
<https://doi.org/10.31901/24566772.2018/12.01.519>

- Seo, Y., Wang, J., Barnes, D., & Barshikar, S. (2021). Heart failure and disability in community-dwelling older adults in the United States—1999 to 2018: Data from National Health and Nutrition Examination Survey. *Clinical Nursing Research*, 571–578.
<https://doi.org/10.1177/10547738211061620>
- Serdar, C. C., Cihan, M., Yücel, D., & Serdar, M. A. (2021). Sample size, power and effect size revisited: Simplified and practical approaches in pre-clinical, clinical and laboratory studies. *Biochemia Medica*, 31(1), 27–53. <https://doi.org/10.11613/BM.2021.010502>
- Sethares, K. A., & Chin, E. (2021). Age and gender differences in physical heart failure symptom clusters. *Heart & Lung*, 50(6), 832–837.
<https://doi.org/10.1016/j.hrtlng.2021.07.001>
- Sewell, J. L., & Velayos, F. S. (2013). Systematic review: The role of race and socioeconomic factors on IBD healthcare delivery and effectiveness. *Inflammatory Bowel Diseases*, 19(3), 627–643. <https://doi.org/10.1002/ibd.22986>
- Sharma, A., Colvin-Adams, M., & Yancy, C. W. (2014). Heart failure in African Americans: Disparities can be overcome. *Cleveland Clinical Journal of Medicine*, 81(5), 301–311.
http://www.abccardio.org/wp-content/uploads/2016/11/Yancy_Colvin.pdf
- Sharma, V., Zehtabchi, S., Rojas, N., & Birkhahn, R. (2009). Ethnic variations in quality of life and depressive symptoms among Black Americans with acute decompensated heart failure. *Journal of the National Medical Association*, 101(10), 985–991.
[https://doi.org/10.1016/S0027-9684\(15\)31064-6](https://doi.org/10.1016/S0027-9684(15)31064-6)
- Shavers, V. L. (2007). Measurement of socioeconomic status in health disparities research. *Journal of the National Medical Association*, 99(9), 1013–1023.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2575866/>

- Shen, B.-J., Eisenberg, S. A., Maeda, U., Farrell, K. A., Schwarz, E. R., Penedo, F. J., Bauerlein, E. J., & Mallon, S. (2010). Depression and anxiety predict decline in physical health functioning in patients with heart failure. *Annals of Behavioral Medicine, 41*(3), 373–382. <https://doi.org/10.1007/s12160-010-9251-z>
- Shetty, P., Mane, A., Fulmali, S., & Uchit, G. (2018). Understanding masked depression: A clinical scenario. *Indian Journal of Psychiatry, 60*(1), 97–102. https://doi.org/10.4103/psychiatry.IndianJPsychiatry_272_17
- Siler, S., Arora, K., Doyon, K., & Fischer, S. M. (2021). Spirituality and the illness experience: Perspectives of African American older adults. *American Journal of Hospice & Palliative Medicine, 38*(6), 618–625. <https://doi.org/10.1177/1049909120988280>
- Silver, M. A. (2010). Depression and heart failure: An overview of what we know and don't know. *Cleveland Clinic Journal of Medicine, 77*(Suppl 3), S7–S11.
- Simmonds, G. (2008). African American participation in public health research. *ABNF Journal, 19*(2), 69–72.
- Slack, P., & Aziz, V. M. (2020). Sexuality and sexual dysfunctions in older people: A forgotten problem. *BJPsych Advances, 26*(3), 173–182. <https://doi.org/10.1192/bja.2019.80>
- Sleath et al., 1998**
- Sokoreli, I., Vries, J., Pauws, S., & Steyerberg, E. (2016). Depression and anxiety as predictors of mortality among heart failure patients: Systematic review and meta-analysis. *Heart Failure Reviews, 21*(1), 49–63. <https://doi.org/10.1007/s10741-015-9517-4>
- Son, Y.-J., & Won, M. H. (2018). Symptom clusters and their impacts on hospital readmission in patients with heart failure: A cross-sectional study. *Research and Theory for Nursing Practice, 32*(3), 311–327. <https://doi.org/10.1891/1541-6577.32.3.311>

- Spertus, J. A., Jones, P. G., Sandhu, A. T., & Arnold, S. V. (2020). Interpreting the Kansas City Cardiomyopathy Questionnaire in clinical trials and clinical care: *JACC* state-of-the-art review. *Journal of the American College of Cardiology*, *76*(20), 2379–2390.
<https://doi.org/10.1016/j.jacc.2020.09.542>
- Spruill, I. J. (2010). Enhancing recruitment of African-American families into genetic research: Lessons learned from Project SuGar. *Journal of Community Genetics*, *1*(3), 125–132.
<https://doi.org/10.1007/s12687-010-0017-x>
- Stamp, K. D., Dunbar, S. B. R., Clark, P. C., Reilly, C. M., Gary, R. A., Higgins, M., & Kaslow, N. (2014). Family context influences psychological outcomes of depressive symptoms and emotional quality of life in patients with heart failure. *Journal of Cardiovascular Nursing*, *29*(6), 517–527. <https://doi.org/10.1097/JCN.0000000000000097>
- Suarez, L., Beach, S. R., Moore, S. V., Mastromauro, C. A., Januzzi, J. L., Celano, C. M., Chang, T. E., & Huffman, J. C. (2015). Use of the Patient Health Questionnaire-9 and a detailed suicide evaluation in determining imminent suicidality in distressed patients with cardiac disease. *Psychosomatics*, *56*(2), 181–189.
<https://doi.org/10.1016/j.psych.2014.12.005>
- Sztajzel, J. (2015). Chronic heart failure and its consequences on the partner relationship. *Revue Médicale Suisse*, *11*(498), 2313–2315.
- Tandon, V. V., Stringer, B. B., Conner, C. C., Gabriel, A. A., Tripathi, B. B., Balakumaran, K. K., & Chen, K. K. (2020). An observation of racial and gender disparities in congestive heart failure admissions using the national inpatient sample. *Curēus*, *12*(10), Article e10914. <https://doi.org/10.7759/cureus.10914>

- Tansey, P. (2010). Counting the cost of heart failure to the patient, the nurse and the NHS. *British Journal of Nursing*, 19(22), 1396–1401.
<https://doi.org/10.12968/bjon.2010.19.22.1396>
- Tański, W., Dudek, K., Tomaszewicz, A., & Świątoniowska-Lonc, N. (2022). Sexual dysfunction and quality of life in patients with rheumatoid arthritis. *International Journal of Environmental Research and Public Health*, 19(5), Article 3088.
<https://doi.org/10.3390/ijerph19053088>
- Tesio, V., Molinaro, S., Castelli, L., Andriani, M., Fanelli, A. L., Marra, S., & Torta, R. (2014). Screening of depression in cardiac patients: Is the HADS an effective tool? *International Journal of Cardiology*, 172(2), e344–e345. <https://doi.org/10.1016/j.ijcard.2013.12.308>
- Thomas, K. L., Hernandez, A. F., Dai, D., Heidenreich, P., Fonarow, G. C., Peterson, E. D., & Yancy, C. W. (2011). Association of race/ethnicity with clinical risk factors, quality of care, and acute outcomes in patients hospitalized with heart failure. *American Heart Journal*, 161(4), 746–754. <https://doi.org/10.1016/j.ahj.2011.01.012>
- Trantham, L. C., Carpenter, W. R., DiMartino, L. D., White, B., Green, M., Teal, R., Corbie-Smith, G., & Godley, P. A. (2015). Perceptions of cancer clinical research among African American men in North Carolina. *Journal of the National Medical Association*, 107(1), 33–41. [https://doi.org/10.1016/S0027-9684\(15\)30007-9](https://doi.org/10.1016/S0027-9684(15)30007-9)
- Trivedi, R. B., Blumenthal, J. A., O'Connor, C., Adams, K., Hinderliter, A., Dupree, C., Johnson, K., & Sherwood, A. (2009). Coping styles in heart failure patients with depressive symptoms. *Journal of Psychosomatic Research*, 67(4), 339–346.
<https://doi.org/10.1016/j.jpsychores.2009.05.014>

- Tromp, J., Bamadhaj, S., Cleland, J. G. F., Angermann, C. E., Dahlstrom, U., Ouwerkerk, W., Tay, W. T., Dickstein, K., Ertl, G., Hassanein, M., Perrone, S. V., Ghadanfar, M., Schweizer, A., Obergfell, A., Lam, C. S. P., Filippatos, G., & Collins, S. P. (2020). Post-discharge prognosis of patients admitted to hospital for heart failure by world region, and national level of income and income disparity (REPORT-HF): A cohort study. *The Lancet Global Health*, 8(3), e411–e422. [https://doi.org/10.1016/S2214-109X\(20\)30004-8](https://doi.org/10.1016/S2214-109X(20)30004-8)
- Tsao, C. W., Aday, A. W., Almarzooq, Z. I., Alonso, A., Beaton, A. Z., Bittencourt, M. S., Boehme, A. K., Buxton, A. E., Carson, A. P., Commodore-Mensah, Y., Elkind, M. S. V., Evenson, K. R., Eze-Nliam, C., Ferguson, J. F., Generoso, G., Ho, J. E., Kalani, R., Khan, S. S., Kissela, B. M., ... Martin, S. S. (2022). Heart disease and stroke statistics—2022 update: A report from the American Heart Association. *Circulation*, 145(8), e153–e639. <https://doi.org/10.1161/CIR.0000000000001052>
- Tutoglu, A., Boyaci, A., Koca, I., Celen, E., & Korkmaz, N. (2014). Quality of life, depression, and sexual dysfunction in spouses of female patients with fibromyalgia. *Rheumatology International*, 34(8), 1079–1084. <https://doi.org/10.1007/s00296-014-2944-z>
- Tuzun, H., Aycan, S., & Ilhan, M. N. (2015). Impact of comorbidity and socioeconomic status on quality of life in patients with chronic diseases who attend primary health care centres. *Central European Journal of Public Health*, 23(3), 188–194. <https://doi.org/10.21101/cejph.a3990>
- Urbich, M., Globe, G., Pantiri, K., Heisen, M., Bennison, C., Wirtz, H. S., & Di Tanna, G. L. (2020). A systematic review of medical costs associated with heart failure in the USA (2014–2020). *Pharmacoeconomics*, 38(11), 1219–1236. <https://doi.org/10.1007/s40273-020-00952-0>

- Usher, B. M., & Cammarata, K. (2009). Heart failure and family caregiver burden: An update. *Progress in Cardiovascular Nursing*, 24(3), 113–114. <https://doi.org/10.1111/j.1751-7117.2009.00046.x>
- Vaccarino, V., Gahbauer, E., Kasl, S. V., Charpentier, P. A., Acampora, D., & Krumholz, H. M. (2002). Differences between African Americans and Whites in the outcome of heart failure: Evidence for a greater functional decline in African Americans. *American Heart Journal*, 143(6), 1058–1067. <https://doi.org/10.1067/mhj.2002.122123>
- Vaccarino, V., Kasl, S. V., Abramson, J., & Krumholz, H. M. (2001). Depressive symptoms and risk of functional decline and death in patients with heart failure. *Journal of the American College of Cardiology*, 38(1), 199–205. [https://doi.org/10.1016/S0735-1097\(01\)01334-1](https://doi.org/10.1016/S0735-1097(01)01334-1)
- van Driel, A. G., de Hosson, M. J. J., & Gamel, C. (2014). Sexuality of patients with chronic heart failure and their spouses and the need for information regarding sexuality. *European Journal of Cardiovascular Nursing*, 13(3), 227–234. <https://doi.org/10.1177/1474515113485521>
- Verma, A. K., Schulte, P. J., Bittner, V., Keteyian, S. J., Fleg, J. L., Piña, I. L., Swank, A. M., Fitz-Gerald, M., Ellis, S. J., Kraus, W. E., Whellan, D. J., O'Connor, C. M., & Mentz, R. J. (2016). Socioeconomic and partner status in chronic heart failure: Relationship to exercise capacity, quality of life, and clinical outcomes. *The American Heart Journal*, 183, 54–61. <https://doi.org/10.1016/j.ahj.2016.10.007>

Virani, S. S., Alonso, A., Aparicio, H. J., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Cheng, S., Delling, F. N., Elkind, M. S. V., Evenson, K. R., Ferguson, J. F., Gupta, D. K., Khan, S. S., Kissela, B. M., Knutson, K. L., Lee, C. D., Lewis, T. T., ... Tsao, C. W. (2021). Heart disease and stroke statistics—2021 update: A report from the American Heart Association. *Circulation*, *143*(8), e254–e743.

<https://doi.org/10.1161/CIR.0000000000000950>

Virani, S. S., Alonso, A., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Chang, A. R., Cheng, S., Delling, F. N., Djousse, L., Elkind, M. S. V., Ferguson, J. F., Fornage, M., Khan, S. S., Kissela, B. M., Knutson, K. L., Kwan, T. W., Lackland, D. T., ... Tsao, C. W. (2020). Heart Disease and Stroke Statistics—2020 update: A report from the American Heart Association. *Circulation*, *141*(9), e139–e151.

<https://doi.org/10.1161/CIR.0000000000000757>

von Haehling, S., Schefold, J. C., Lainscak, M., Doehner, W., & Anker, S. D. (2009).

Inflammatory biomarkers in heart failure revisited: Much more than innocent bystanders.

Heart Failure Clinics, *5*(4), 549–560. <https://doi.org/10.1016/j.hfc.2009.04.001>

Vorderstrasse, A., Lewinski, A., Melkus, G. D. E., & Johnson, C. (2016). Social support for diabetes self-management via eHealth interventions. *Current Diabetes Reports*, *16*(7),

Article 56. <https://doi.org/10.1007/s11892-016-0756-0>

Ward, B. W., & Schiller, J. S. (2013). Prevalence of multiple chronic conditions among US adults: Estimates from the National Health Interview Survey, 2010. *Preventing Chronic Disease*, *10*, Article E65.

<https://doi.org/10.5888/pcd10.120203>

Warnecke, R. B., Ferrans, C. E., Johnson, T. P., Chapa-Resendez, G., O'Rourke, D. P., Chávez, N., Dudas, S., Smith, E. D., Martínez Schallmoser, L., Hand, R. P., & Lad, T. (1996). Measuring quality of life in culturally diverse populations. *Journal of the National Cancer Institute Monographs*, 20(1996), 29–38.

Warren-Findlow, J., & Issel, L. M. (2010). Stress and coping in African American women with chronic heart disease: A cultural cognitive coping model. *Journal of Transcultural Nursing*, 21(1), 45–54. <https://doi.org/10.1177/1043659609348622>

Watson, 1995

Wei, S., Le, N., Zhu, J. W., Breathett, K., Greene, S. J., Mamas, M. A., Zannad, F., & Van Spall, H. G. C. (2022). Factors associated with racial and ethnic diversity among heart failure trial participants: A systematic bibliometric review. *Circulation: Heart Failure*, 15(3), Article e008685. <https://doi.org/10.1161/CIRCHEARTFAILURE.121.008685>

Weiss, B. D., Mays, M. Z., Martz, W., Castro, K. M., DeWalt, D. A., Pignone, M. P., Mockbee, J., & Hale, F. A. (2005). Quick assessment of literacy in primary care: The newest vital sign. *Annals of Family Medicine*, 3(6), 514–522. <https://doi.org/10.1370/afm.405>

Westlake, C., & Dracup, K. (2001). Role of spirituality in adjustment of patients with advanced heart failure. *Progress in Cardiovascular Nursing*, 16(3), 119–125.

<https://doi.org/10.1111/j.0889-7204.2001.00592.x>

Westlake, C., Dracup, K., Creaser, J., Livingston, N., Heywood, J. T., Huiskes, B. L., Fonarow, G., & Hamilton, M. (2002). Correlates of health-related quality of life in patients with heart failure. *Heart & Lung*, 31(2), 85–93. <https://doi.org/10.1067/mhl.2002.122839>

- Westlake, C., Dracup, K., Walden, J. A., & Fonarow, G. (1999). Sexuality of patients with advanced heart failure and their spouses or partners. *The Journal of Heart and Lung Transplantation, 18*(11), 1133–1138. [https://doi.org/10.1016/S1053-2498\(99\)00084-4](https://doi.org/10.1016/S1053-2498(99)00084-4)
- Westlake, C., Sethares, K., & Davidson, P. (2013). How can health literacy influence outcomes in heart failure patients? Mechanisms and interventions. *Current Heart Failure Reports, 10*(3), 232–243. <https://doi.org/10.1007/s11897-013-0147-7>
- Whittingham, K., Barnes, S., & Gardiner, C. (2013). Tools to measure quality of life and carer burden in informal carers of heart failure patients: A narrative review. *Palliative Medicine, 27*(7), 596–607. <https://doi.org/10.1177/0269216313477179>
- Wierenga, K. L., Dekker, R. L., Lennie, T. A., Chung, M. L., & Dracup, K. (2017). African American race is associated with poorer outcomes in heart failure patients. *Western Journal of Nursing Research, 39*(4), 524–538. <https://doi.org/10.1177/0193945916661277>
- Williams, D. R., Gonzalez, H. M., Neighbors, H., Nesse, R., Abelson, J. M., Sweetman, J., & Jackson, J. S. (2007). Prevalence and distribution of major depressive disorder in African Americans, Caribbean Blacks, and non-Hispanic Whites: Results from the National Survey of American Life. *Archives of General Psychiatry, 64*(3), 305–315. <https://doi.org/10.1001/archpsyc.64.3.305>
- Winkleby, M. A., Jatulis, D. E., Frank, E., & Fortmann, S. P. (1992). Socioeconomic status and health: How education, income, and occupation contribute to risk factors for cardiovascular disease. *American Journal of Public Health, 82*(6), 816–820. <https://doi.org/10.2105/AJPH.82.6.816>

- Wispelwey, B., Chang, C. Y., Brooks, K. C., Kakoza, R., Loscalzo, J., Lewis, E. F., Avutu, V., Baird, K., Bakshi, S., Balaban, D., Baumrin, E., Bayomy, O., Beamesderfer, J., Becker, N., Berman, A. N., Boden, K., Bonacci, R. A., Chiang, D., Clark, D., ... Yazdi, D. (2019). Identification of racial inequities in access to specialized inpatient heart failure care at an academic medical center. *Circulation: Heart Failure*, 12(11), Article e006214. <https://doi.org/10.1161/CIRCHEARTFAILURE.119.006214>
- Wu, J.-R., Lennie, T. A., Frazier, S. K., & Moser, D. K. (2016). Health-related quality of life, functional status, and cardiac event-free survival in patients with heart failure. *The Journal of Cardiovascular Nursing*, 31(3), 236–244. <https://doi.org/10.1097/JCN.0000000000000248>
- Wu, V. C.-C., Chang, S.-H., Kuo, C.-F., Liu, J.-R., Chen, S.-W., Yeh, Y.-H., Luo, S.-F., & See, L.-C. (2018). Suicide death rates in patients with cardiovascular diseases – A 15-year nationwide cohort study in Taiwan. *Journal of Affective Disorders*, 238, 187–193. <https://doi.org/10.1016/j.jad.2018.05.046>
- Wu, J. R., Lennie, T. A., Frazier, S. K., & Moser, D. K. (2016). Health-related quality of life, functional status and cardiac event-free survival in patients with heart failure. *The Journal of cardiovascular nursing*, 31(3), 236.
- Wu, Y., Levis, B., Sun, Y., He, C., Krishnan, A., Neupane, D., Bhandari, P. M., Negeri, Z., Benedetti, A., & Thombs, B. D. (2021). Accuracy of the Hospital Anxiety and Depression Scale Depression subscale (HADS-D) to screen for major depression: Systematic review and individual participant data meta-analysis. *BMJ*, 373(972). <https://doi.org/10.1136/bmj.n972>

- Wu, Y., Levis, B., Sun, Y., Krishnan, A., He, C., Riehm, K. E., Rice, D. B., Azar, M., Yan, X. W., Neupane, D., Bhandari, P. M., Imran, M., Chiovitti, M. J., Saadat, N., Boruff, J. T., Cuijpers, P., Gilbody, S., McMillan, D., Ioannidis, J. P. A., ... Thombs, B. D. (2020). Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale – Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. *Journal of Psychosomatic Research*, 129, Article 109892. <https://doi.org/10.1016/j.jpsychores.2019.109892>
- Xiong, G. L., Prybol, K., Boyle, S. H., Hall, R., Streilein, R. D., Steffens, D. C., Krishnan, R., Rogers, J. G., O'Connor, C. M., & Jiang, W. (2015). Inflammation markers and major depressive disorder in patients with chronic heart failure: Results from the Sertraline Against Depression and Heart Disease in Chronic Heart Failure study. *Psychosomatic Medicine*, 77(7), 808–815. <https://doi.org/10.1097/PSY.0000000000000216>
- Yazdanpanahi, Z., Nikkholgh, M., Akbarzadeh, M., & Pourahmad, S. (2018). Stress, anxiety, depression, and sexual dysfunction among postmenopausal women in Shiraz, Iran, 2015. *Journal of Family & Community Medicine*, 25(2), 82–87. https://doi.org/10.4103/jfcm.JFCM_117_17
- Yazici, R., Altintepe, L., Guney, I., Yeksan, M., Atalay, H., Turk, S., Tonbul, H. Z., & Selcuk, N. Y. (2009). Female sexual dysfunction in peritoneal dialysis and hemodialysis patients. *Renal Failure*, 31(5), 360–364. <https://doi.org/10.1080/08860220902883012>

- Yohannes, A. M., Willgoss, T. G., Baldwin, R. C., & Connolly, M. J. (2010). Depression and anxiety in chronic heart failure and chronic obstructive pulmonary disease: Prevalence, relevance, clinical implications and management principles. *International Journal of Geriatric Psychiatry*, 25(12), 1209–1221. <https://doi.org/10.1002/gps.2463>
- Zhang, X., Zou, H., Hou, D., He, D., & Fan, X. (2020). Functional status mediates the association of nutritional status with depressive symptoms in patients with heart failure. *Journal of Advanced Nursing*, 76(12), 3363–3371. <https://doi.org/10.1111/jan.14522>
- Ziaeeian, B., Kominski, G. F., Ong, M. K., Mays, V. M., Brook, R. H., & Fonarow, G. C. (2017). National differences in trends for heart failure hospitalizations by sex and race/ethnicity. *Circulation: Cardiovascular Quality and Outcomes*, 10(7). <https://doi.org/10.1161/CIRCOUTCOMES.116.003552>
- Zou, H., Chen, Y., Fang, W., Zhang, Y., & Fan, X. (2017). Identification of factors associated with self-care behaviors using the COM-B model in patients with chronic heart failure. *European Journal of Cardiovascular Nursing*, 16(6), 530–538.

APPENDICES

Table 10

Characteristics Under Study

Characteristics Under Study	Variables
Sociodemographic Characteristics	Race Age Gender Marital Status Education Employment Status
Clinical Characteristics	Disease Severity Comorbidities Functional Status
Psychological Characteristics	Depression Anxiety
Quality of Life	Quality of Life (QOL) Health Functioning QOL Social and Economic QOL Psychological/Spiritual QOL Family QOL

Table 11*Aims, Variables, and Measures*

Aim	Variables	Instruments/measures
1. Describe the differences and examine the relationships between or among sociodemographic, clinical, and psychological characteristics and QOL between Blacks and Whites with HF.	Sociodemographic characteristics <ul style="list-style-type: none"> • Race* • Age* • Gender* • Marital status* • Education* • Employment status* Clinical characteristics <ul style="list-style-type: none"> • Disease severity • Comorbidities • Functional status 	*Indicates a variable that is its own measure <i>Disease severity</i> <ul style="list-style-type: none"> ▪ Ejection fraction ▪ New York Heart Association (NYHA) Classification <i>Comorbidities</i> <ul style="list-style-type: none"> ▪ Charlson Comorbidity Index <i>Functional status – MOS 36-Item Short Form (SF-36)</i> <ul style="list-style-type: none"> ▪ Functional limitations • Mobility limitations
2. Examine the influence of sociodemographic clinical and psychological characteristics on QOL in Blacks and Whites with HF.	Psychological characteristics <ul style="list-style-type: none"> • Depression • Anxiety Quality of life	<i>Hospital Anxiety and Depression Scale (HADS)</i> <ul style="list-style-type: none"> ▪ Depression (HADS – Depression Subscale) ▪ Anxiety (HADS – Anxiety Subscale) <i>Quality of Life Index (QLI – Cardiac IV)</i> <ul style="list-style-type: none"> ▪ Health Functioning Subscale ▪ Social and Economic Subscale ▪ Psychological/Spiritual Subscale • Family Subscale

Table 12*Statistical Analysis Plan*

Aim	Variables/measures	Statistical tests
1. Describe the differences in sociodemographic, clinical, and psychological characteristics and QOL between Blacks and Whites with HF.	Sociodemographic characteristics <ul style="list-style-type: none"> • Race • Age • Gender • Marital status • Education • Employment status 	Mean/median <i>t</i> tests/ Chi-square tests
2. Examine the relationships between or among sociodemographic, clinical, and psychological characteristics and QOL in Blacks and Whites with HF.	Clinical characteristics <ul style="list-style-type: none"> • Ejection fraction • NYHA Class • Charlson Comorbidity Index • Functional status: MOS 36-Item Short Form (SF-36) 	Pearson/Spearman correlation
3. Examine the influence of sociodemographic clinical and psychological characteristics on QOL in Blacks and Whites with HF.	Psychological characteristics <ul style="list-style-type: none"> • Hospital Anxiety & Depression Scale (HADS) Quality of life <ul style="list-style-type: none"> • Quality of Life Index (QLI) 	Multiple regression model

Figure 4

Sampling Method

CONSORT 2010 Flow Diagram

